
LIST OF TABLES

Table No	DESCRIPTION	Page No
2.1	Details of chemicals like name, chemical formula, molecular weight, physical appearance and manufacturer name of the reagent used in present work	46
3.1	Percentage of shape distribution present in C0, C1, C2, C3 and C4 AgNPs samples.	70
3.2	Average particle size, activation energy and apparent reaction rate constants for Nip reduction with NaBH ₄ and Gly as reductant in presence of C0, C1, C2, C3, C4 AgNPs catalyst.	77
3.3	Comparison of turnover frequency (TOF) of different catalyst in transfer hydrogenation reaction of nitroarenes using glycerol as hydrogen source.	81
3.4	Turnover frequency (TOF) of in presence of catalyst and NaBH ₄ as reducing agent at 301 K.	82
3.5	Apparent reaction rate constant and turnover frequency (TOF) for thermal and photocatalytic Nip reduction with Gly as reducing agent at 306 K.	84
3.6	Apparent reaction rate constant, activation energy and turnover frequency (TOF) for thermal and photocatalytic MO degradation at 306 K in presence of all AgNPs catalyst.	89
4.1	Average particle size, apparent reaction rate constants and activation energy for Nip reduction with NaBH ₄ and Gly reducing agent in presence of both A1 and A2 c-AgNPs catalyst	108
4.2	Comparison of turnover frequencies (TOF) for <i>p</i> -nitrophenol reduction by NaBH ₄	109
4.3	The photocatalytic apparent reaction rate constant, photocatalytic turnover frequency (PTOF) and TOF for Nip reduction with Gly as reducing agent at 306 K	112

4.4	Apparent reaction rate constant, activation energy and turnover frequency (TOF) for thermal and photocatalytic MO degradation at 306 K in presence of both c-AgNPs catalyst	115
5.1	Average particle size, apparent reaction rate constants and activation energy for Nip reduction with NaBH ₄ and Gly as reducing agent in presence of both B1 and B2 CuNPs catalyst	132
5.2	Comparison of turnover frequencies (TOF) for <i>p</i> -nitrophenol reduction by NaBH ₄ in presence of CuNPs catalyst	133
5.3	Displays the photocatalytic apparent reaction rate constant, photocatalytic turnover frequency (PTOF) and TOF for Nip reduction with Gly as reducing agent at 306 K	135
5.4	Apparent reaction rate constant, activation energy and turnover frequency (TOF) for thermal and photocatalytic MO degradation at 306 K in presence of both CuNPs catalyst	138
6.2.1	Catalytic activity parameters k_{app} , E_a and TOF values for different catalysts	155
6.2.2	Comparison of k_{app} values for both thermal and photocatalytic Nip reduction by Gly in presence of S2, S3 and S4 Ag-Cu BNPs catalyst	158
6.2.3	Apparent reaction rate constant and turnover frequency (TOF) for thermal and photocatalytic MO degradation at 306 K in presence of S1, S2, S3, S4 and S5 catalyst are tabulated here. Activation energies for the thermally catalyzed reaction are also mentioned in column 4 of the Table	161
6.3.1	Average particle size, apparent reaction rate constants and activation energy for Nip reduction with NaBH ₄ and Gly as reducing agent in presence of both S6 and S7BNPs catalyst	174
6.3.2	Displays the photocatalytic apparent reaction rate constant photocatalytic turnover frequency (PTOF) and TOF for Nip reduction with Gly as reducing agent at 306 K.	175
6.3.3	Apparent reaction rate constant, activation energy and turnover frequency (TOF) for thermal and photocatalytic MO degradation at	178

306 K in presence of both S6 and S7 BNPs catalyst

6.4.1	Average particle size, apparent reaction rate constants and activation energy for Nip reduction with NaBH ₄ and Gly as reducing agent in presence of both S8 and S9 BNPs catalyst	188
6.4.2	Displays the photocatalytic apparent reaction rate constant, photocatalytic turn over frequency (PTOF) and TOF for Nip reduction with Gly as reducing agent at 306 K	190
6.4.3	Apparent reaction rate constant, activation energy and turnover frequency (TOF) for thermal and photocatalytic MO degradation at 306 K in presence of both S8 and S9 BNPs catalyst	193
7.1	Average particle sizes, apparent reaction rate constant, activation energy and turnover frequency (TOF) for thermal Nip reduction with NaBH ₄ at 306 K in presence of G1, G2, G3, G4 and G5 catalyst	212
7.2	Apparent reaction rate constant, activation energy and turnover frequency (TOF) for thermal and photocatalytic MO degradation at 306 K in presence of G1, G2, G3, G4 and G5 catalyst	215
8.1	Comparison of Turn over frequencies for Nip reduction with NaBH ₄	220
8.2	Comparison of thermal and photocatalytic Turn over frequencies for Nip reduction with Gly	222
8.3	Comparison of thermal and photocatalytic Turn over frequencies for oxidative MO degradation	223