LIST OF TABLES

| Table No. | Table Title | Pg. No. |
|-----------|---|---------|
| 2.1 | Reported preclinical observations made with diverse types of <i>Curcuma longa</i> extracts | 17 |
| 2.2 | Some reported pharmacological activities of Curcumin | 19 |
| 2.3 | Marketed major formulations containing Curcuma longa extracts | 21 |
| 2.4 | Marketed major formulations containing curcumin | 25 |
| 4.1 | Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on body weight difference of male mice on day 5 | 105 |
| 4.2 | Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on body weight difference of male mice on day 7 | 105 |
| 4.3 | Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on body weight difference of male mice on day 10 | 106 |
| 4.4 | Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on body weight difference of female mice on day 5 | 106 |
| 4.5 | Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on body weight difference of female mice on day 7 | 107 |
| 4.6 | Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on body weight difference of female mice on day 10 | 107 |
| 4.7 | Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on tail suspension test in male mice | 108 |
| 4.8 | Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on tail suspension test in female mice | 108 |
| 4.9 | Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on pentobarbital induced sleep test in male mice | 109 |
| 4.10 | Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on pentobarbital induced sleep test in female mice | 109 |
| 4.11 | Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on glucose, insulin and corticosterone level in non-stressed and stressed male rats | 110 |

| 4.12 | Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on organ weight of stressed male rats | 110 |
|------|---|-----|
| 4.13 | Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on hot plate reaction time of saline and formalin injected male rats | 111 |
| 4.14 | Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on ratio of HDL and LDL of normal drinking water and fructose water consuming male rats | 111 |
| 4.15 | Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on organ weight of normal drinking water and fructose water consuming male rats | 112 |
| 4.16 | Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on immobility time period of normal and stressed diabetic rats in forced swim test. | 112 |
| 4.17 | Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on stomach ulceration index in normal and stressed diabetic rats | 113 |