List of Tables

Page No.

3.1	Table for release rate constant (k), correlation coefficient (r^2) and diffusion	82
	release exponent (n) obtained using different mathematical models for	
	drug loaded PU and its indicated nanocomposites.	

4.1 Table for release rate constant (k), correlation coefficient (r^2) and diffusion **109** release exponent (n) obtained using different mathematical models for drug loaded PU and its indicated nanocomposites.

List of Schemes

2.1	Reaction scheme for polyurethane synthesis.	Page No. 46
2.2	Reaction scheme for polyurethane / graphene nanocomposites synthesis.	47
2.3	Reaction scheme for synthesis of chemically tagged nanocomposites.(b) (iⅈ)- schematic representation of physically mixed (PUG-P) and chemically grafted (PUG-C) nanocomposites.	50
2.4	Reaction scheme for synthesis of diamine modified graphene nanocomposites.	51
2.5	Modification chemistry of graphene oxide and grafting of modified graphene with prepolymer chain.	52
3.3	(e) Schematic representation of stacking showing the interaction site without and with graphene layers in pristine PU and nanocomposites.	69
3.10	(c) Schematic model of drug release kinetics. The red sphere and green plates indicates the drug and graphene sheet, respectively. The arrows indicate the probable path of drug diffusion in pure PU and nanocomposites demonstrating sustained delivery in nanocomposites as compared to pure PU.	82
4.11	(c) Schematic model of drug release kinetics. The black line represents the polymer chains whereas plates indicate the graphene sheets. Arrow shows the probable path for drug diffusion.	109
5 1 1	(c) Schematic model of drug release kinetics. The green spheres and disc	132

5.11 (c) Schematic model of drug release kinetics. The green spheres and disc 132 represents the drug molecule and modified graphene respectively. Disc size increase with increasing the modifier chain length and arrow represents the probable path for drug release through the matrix.