

## *LIST OF FIGURES*

---

	<b>Page No.</b>
<b>Figure 1.1.</b> Schematic illustration of top-down and bottom-up approaches for the preparation of NMNPs.	<b>1</b>
<b>Figure 1.2.</b> Visual images of AgNPs and their corresponding UV-vis spectra.	<b>4</b>
<b>Figure 1.3.</b> TEM image of AgNPs.	<b>4</b>
<b>Figure 1.4.</b> Visual images of PdNPs and their corresponding UV-vis spectra.	<b>6</b>
<b>Figure 1.5.</b> TEM image of PdNPs.	<b>6</b>
<b>Figure 1.6.</b> Several possible representative architectures of BMNPs including (a) an alloy, (b) core-shell, (c) cluster-on-cluster, (d) sub-shell, and (e) intermetallic.	<b>9</b>
<b>Figure 1.7.</b> UV-Vis spectra of bimetallic nanoparticles of Au-Pd.	<b>9</b>
<b>Figure 2.1.</b> UV-Vis absorption spectra for the interaction between increasing concentrations of 3-APTMS into constant methanolic suspension of GPTMS.	<b>37</b>
<b>Figure 2.2.</b> UV-Vis absorption spectra for the interaction between increasing concentrations of GPTMS into constant methanolic suspension of 3-APTMS.	<b>37</b>
<b>Figure 2.3.</b> UV-Vis absorption spectra for the interaction between increasing concentrations of 3-APTMS into constant methanolic suspension of GPTMS containing Ag <sup>+</sup> . Inset shows the respective recording between 380–500 nm.	<b>38</b>
<b>Figure 2.4.</b> UV-Vis absorption spectra for the interaction between increasing concentrations of GPTMS into constant methanolic suspension of 3-APTMS containing Ag <sup>+</sup> . Inset shows the respective recording between 380–500 nm.	<b>38</b>
<b>Figure 2.5.</b> Visual photographs of AgNPs made by using composition of 3-APTMS, GPTMS and AgNO <sub>3</sub> as shown in Table 2.1 for respective samples.	<b>39</b>
<b>Figure 2.6.</b> Variation in absorbance as function of wavelength on the subsequent additions of 3-APTMS (i) and 25 mM HAuCl <sub>4</sub> (ii).	<b>39</b>
<b>Figure 2.7.</b> Variation in absorbance as function of wavelength on the subsequent additions of 3-APTMS (i) and 25 mM AgNO <sub>3</sub> (ii).	<b>40</b>

<b>Figure 2.8.</b>	The change in absorbance after immediate addition of GPTMS in 3-APTMS capped gold ions (i) followed by subsequent recording of change in absorbance as a function of time: after 1 h (ii) and 6 h (iii).	<b>40</b>
<b>Figure 2.9.</b>	The change in absorbance after immediate addition of GPTMS in 3-APTMS capped silver ions (i) followed by subsequent recording of change in absorbance as a function of time: after 1 h (ii) and 6h (iii).	<b>41</b>
<b>Figure 2.10.</b>	(a) TEM image of AgNPs made by 0.05 M 3-APTMS and 4 M GPTMS; (b) particle size distribution.	<b>41</b>
<b>Figure 2.11.</b>	(a) TEM image of AgNPs made by 0.5 M 3-APTMS and 4 M GPTMS; (b) particle size distribution.	<b>42</b>
<b>Figure 2.12.</b>	(a) TEM image of AgNPs made by 0.5 M 3-APTMS and 2 M GPTMS; (b) particle size distribution.	<b>42</b>
<b>Figure 2.13.</b>	Image shows the EDX of AgNPs	<b>42</b>
<b>Figure 2.14.</b>	UV-Vis absorption spectra of AgNP <sub>1</sub> and AgNP <sub>2</sub> in methanol and butanol. The curve as shown with red line represents the variation in absorbance in different solvents for AgNP <sub>1</sub> ( $\lambda_{\text{max}}$ 402 and 406 nm) and the curves with black line represent for AgNP <sub>2</sub> ( $\lambda_{\text{max}}$ 410 and 414 nm).	<b>43</b>
<b>Figure 2.15.</b>	UV-Vis absorption spectra of AgNP <sub>1</sub> and AgNP <sub>2</sub> in acetonitrile and dichloromethane. The curve as shown with red line represents the variation in absorbance in different solvents for AgNP <sub>1</sub> ( $\lambda_{\text{max}}$ 404 and 410 nm) and the curves with black line represent for AgNP <sub>2</sub> ( $\lambda_{\text{max}}$ 416 and 419 nm).	<b>43</b>
<b>Figure 2.16.</b>	(A) Visual photographs of AgNPs made by using composition of 3-APTMS, GPTMS and AgNO <sub>3</sub> as shown in Table 2.1. (Sample I–IV) and their dispersibility in water. (B) Shows the UV-Vis spectra of the corresponding sols in water.	<b>44</b>
<b>Figure 2.17.</b>	(A) Visual photographs of AgNPs made by using composition of 3-APTMS, GPTMS and AgNO <sub>3</sub> as shown in Table 2.1. (Sample I–IV) and their dispersibility in toluene. (B) Shows the UV-Vis spectra of the corresponding sols in toluene.	<b>44</b>
<b>Figure 2.18.</b>	(A) Visual photographs of AgNPs made by using composition of 3-APTMS, GPTMS and AgNO <sub>3</sub> as shown in Table 2.1. (Sample V–VIII) and their dispersibility in water. (B) Shows the UV-Vis spectra of the corresponding sols in water.	<b>45</b>

<b>Figure 2.19.</b>	(A) Visual photographs of AgNPs made by using composition of 3-APTMS, GPTMS and AgNO <sub>3</sub> as shown in Table 2.1. (Sample V-VIII) and their dispersibility in toluene. (B) Shows the UV-Vis spectra of the corresponding sols in toluene.	<b>45</b>
<b>Figure 2.20.</b>	Visual photographs of AgNPs made by using composition of 3-APTMS, GPTMS and AgNO <sub>3</sub> as shown in Table 2.1. (Sample IX–XII) and their dispersibility in water. (B) Shows the UV-Vis spectra of the corresponding sols in water.	<b>46</b>
<b>Figure 2.21.</b>	Visual photographs of AgNPs made by using composition of 3-APTMS, GPTMS and AgNO <sub>3</sub> as shown in Table 2.1. (Sample IX–XII) and their dispersibility in toluene. (B) Shows the UV-Vis spectra of the corresponding sols in toluene.	<b>46</b>
<b>Figure 2.22.</b>	The visual photographs of AgNPs containing constant concentrations of AgNO <sub>3</sub> (0.01 M) and Cyclohexanone (1.9 M) followed with varying concentrations of 3-APTMS between 0.1x10 <sup>-3</sup> M to 2 M (Table 2.2, Sr. no. a to j).	<b>50</b>
<b>Figure 2.23.</b>	The visual photographs of AgNPs containing constant concentrations of AgNO <sub>3</sub> (0.01 M) and 3-APTMS (0.5 M) followed with varying concentrations of Cyclohexanone between 0.3 M to 3.8 (Table 2.3, Sr. no. i to x).	<b>50</b>
<b>Figure 2.24.</b>	Real time synthesis of AgNPs using 0.5 M 3-APTMS and 0.8 M concentration of formaldehyde.	<b>51</b>
<b>Figure 2.25.</b>	Real time synthesis of AgNPs using 2 M 3-APTMS and 0.8 M concentration of formaldehyde.	<b>51</b>
<b>Figure 2.26.</b>	Formation of AgNPs as a function of 3-APTMS concentrations (Table 2.4, Sr. no. a to j) keeping constant concentrations of Formaldehyde (0.8 M) and 0.01 M AgNO <sub>3</sub> .	<b>52</b>
<b>Figure 2.27.</b>	Effect of Formaldehyde concentrations on the formation of AgNPs [(a) 0.4 M, (b) 0.8 M and (c) 1.2 M] keeping constant concentration of 3-APTMS (0.5 M).	<b>52</b>
<b>Figure 2.28.</b>	FTIR spectrum of AgNPs.	<b>53</b>
<b>Figure 2.29.</b>	TEM (a) and selected area electron diffraction pattern (b) of AgNPs consist of 0.25 M APTMS and fixed concentration of Cyclohexanone.	<b>53</b>
<b>Figure 2.30.</b>	TEM (a) and selected area electron diffraction pattern (b) of AgNPs consist of 0.5 M APTMS and fixed	<b>54</b>

	concentration of Cyclohexanone.	
<b>Figure 2.31.</b>	TEM (a) and selected area electron diffraction pattern (b) of AgNPs consist of 0.5 M APTMS and fixed concentration of Formaldehyde.	<b>54</b>
<b>Figure 2.32.</b>	UV-vis absorption spectra of increasing concentrations of AgNPs made from cyclohexanone and 3-APTMS in water and methanol; Respective insets show the dependence of absorption maxima ( $\lambda_{\max}$ ) on AgNPs concentration	<b>55</b>
<b>Figure 2.33.</b>	UV-vis absorption spectra of increasing concentrations of AgNPs made from cyclohexanone and 3-APTMS in acetonitrile and toluene; Respective insets show the dependence of absorption maxima ( $\lambda_{\max}$ ) on AgNPs concentration.	<b>55</b>
<b>Figure 2.34.</b>	UV-vis absorption spectra of increasing concentrations of AgNPs made from cyclohexanone and 3-APTMS in dichloromethane; Respective inset show the dependence of absorption maxima ( $\lambda_{\max}$ ) on AgNPs concentration.	<b>56</b>
<b>Figure 2.35.</b>	UV-vis absorption spectra of increasing concentrations of AgNPs made by using formaldehyde and 3-APTMS in water and methanol; Respective insets show the dependence of absorption maxima ( $\lambda_{\max}$ ) on AgNPs concentration.	<b>56</b>
<b>Figure 2.36.</b>	UV-vis absorption spectra of increasing concentrations of AgNPs made by using formaldehyde and 3-APTMS in acetonitrile and butanol; Respective insets show the dependence of absorption maxima ( $\lambda_{\max}$ ) on AgNPs concentration.	<b>57</b>
<b>Figure 2.37.</b>	Images of methanolic and aqueous suspension of GPTMS. Methanolic suspension of GPTMS (A) shows single layer formed while aqueous suspension of GPTMS (B) results two separate layers.	<b>58</b>
<b>Figure 2.38.</b>	UV-Vis spectra of increasing concentrations of AgNP in (A) butanol, (B) methanol, (C) toluene, (D) dichloromethane and (E) acetonitrile. Respective insets show the dependence of absorption maxima ( $\lambda_{\max}$ ) on relative concentration.	<b>63</b>
<b>Figure 2.39.</b>	Recording shows the dependence of absorption maxima ( $\lambda_{\max}$ ) on refractive index of solvents.	<b>63</b>
<b>Figure 3.1.</b>	Variation in the UV-vis spectra of methanolic solution of $K_2PdCl_4$ in absence and the presence of 3-APTMS (A and B) respectively.	<b>75</b>

<b>Figure 3.2.</b>	Variation in the UV-vis spectra of methanolic solution of $K_2PdCl_4$ in absence and the presence of TEOS (A and B) respectively.	<b>75</b>
<b>Figure 3.3.</b>	Effect of 3-APTMS concentrations ranging between $0.001 \times 10^{-3}$ to $50 \times 10^{-3} M$ (images 1–8) on the synthesis of PdNPs from 1.9 M cyclohexanone and 0.005 M $K_2PdCl_4$ as shown in Table 3.2.	<b>76</b>
<b>Figure 3.4.</b>	Effect of cyclohexanone concentrations ranging between 0.3 to 3.8 M (i–x) on the synthesis of PdNPs from $0.001 \times 10^{-3} M$ 3-APTMS and 0.005 M $K_2PdCl_4$ as shown in Table 3.3.	<b>76</b>
<b>Figure 3.5.</b>	Effect of cyclohexanone concentrations ranging between 0.3 to 3.8 M (i–x) on the synthesis of PdNPs from $0.01 \times 10^{-3} M$ 3-APTMS and 0.005 M $K_2PdCl_4$ as shown in Table 3.3.	<b>77</b>
<b>Figure 3.6.</b>	Effect of cyclohexanone concentrations ranging between 0.3 to 3.8 M (i–x) on the synthesis of PdNPs from $0.1 \times 10^{-3} M$ 3-APTMS and 0.005 M $K_2PdCl_4$ as shown in Table 3.3.	<b>77</b>
<b>Figure 3.7.</b>	SAED pattern of PdNPs made by using 1.9 M cyclohexanone in absence of 3-APTMS.	<b>78</b>
<b>Figure 3.8.</b>	SAED pattern of PdNPs made by using 1.9 M cyclohexanone in presence of $0.001 \times 10^{-3} M$ 3-APTMS.	<b>78</b>
<b>Figure 3.9.</b>	SAED pattern of PdNPs made by using 1.9 M cyclohexanone in presence of $0.1 \times 10^{-3} M$ 3-APTMS.	<b>79</b>
<b>Figure 3.10.</b>	(a) TEM image of PdNPs made by using 1.9 M cyclohexanone in absence of 3-APTMS, (b) particle size distribution.	<b>80</b>
<b>Figure 3.11.</b>	(a) TEM image of PdNPs made by using 1.9 M cyclohexanone in presence of $0.001 \times 10^{-3} M$ 3-APTMS, (b) particle size distribution.	<b>80</b>
<b>Figure 3.12.</b>	(a) TEM image of PdNPs made by using 1.9 M cyclohexanone in presence of $0.1 \times 10^{-3} M$ 3-APTMS, (b) particle size distribution.	<b>80</b>
<b>Figure 3.13.</b>	Visual photographs of PdNPs made by using $0.001 \times 10^{-3} M$ 3-APTMS, 1.9 M Cyclohexanone and 0.005 M $K_2PdCl_4$ as shown in Table 3.5 (A) and their dispersibility in water, methanol, acetonitrile and toluene.	<b>81</b>
<b>Figure 3.14.</b>	Visual photographs of PdNPs made by using $0.01 \times 10^{-3} M$ 3-APTMS, 1.9 M Cyclohexanone and 0.005 M $K_2PdCl_4$ as shown in Table 3.5 (B) and their dispersibility in water, methanol, acetonitrile and	<b>81</b>

	toluene.	
<b>Figure 3.15.</b>	Visual photographs of PdNPs made by using $0.1 \times 10^{-3}$ M 3-APTMS, 1.9 M Cyclohexanone and 0.005 M $K_2PdCl_4$ as shown in Table 3.5 (A) and their dispersibility in water, methanol, acetonitrile and toluene.	<b>82</b>
<b>Figure 3.16.</b>	The visual photographs of PdNPs (upper portion). The UV-Vis spectra of the corresponding PdNPs (lower portion).	<b>84</b>
<b>Figure 3.17.</b>	(a) TEM image of PdNPs made by using constant concentration of THF-HPO in presence of 1 M 3-APTMS, (b) particle size distribution.	<b>85</b>
<b>Figure 3.18.</b>	(a) TEM image of PdNPs made by using constant concentration of THF-HPO in presence of 0.5 M 3-APTMS, (b) particle size distribution.	<b>85</b>
<b>Figure 3.19.</b>	Visual photographs of PdNPs made by using 3-APTMS (0.25 M), THF-HPO (11.4 mg) and $K_2PdCl_4$ (0.003 M) as shown in Table 3.4 and their dispersibility in water, methanol, acetonitrile and toluene.	<b>86</b>
<b>Figure 3.20.</b>	Visual photographs of PdNPs made by using 3-APTMS (0.5 M), THF-HPO (11.4 mg) and $K_2PdCl_4$ (0.003 M) as shown in Table 3.4 and their dispersibility in water, methanol, acetonitrile and toluene.	<b>86</b>
<b>Figure 3.21.</b>	Visual photographs of PdNPs made by using 3-APTMS (1 M), THF-HPO (11.4 mg) and $K_2PdCl_4$ (0.003 M) as shown in Table 3.4 and their dispersibility in water, methanol, acetonitrile and toluene.	<b>87</b>
<b>Figure 4.1.</b>	The visual photographs of Au-PdNPs and Pd-AuNPs.	<b>99</b>
<b>Figure 4.2.</b>	UV-Vis spectra recorded on the simultaneous formation of Pd-Au from $K_2PdCl_4$ (80%) and $HAuCl_4$ (20%): the variation of absorbance as the function of wavelength on subsequent addition of $K_2PdCl_4$ (a), $HAuCl_4$ (b), 3-APTMS (c), THF-HPO (d), curve-e shows the absorbance recorded after 1 h from subsequent addition and curve-f shows the spectra recorded after simultaneous formation of Pd-Au.	<b>99</b>
<b>Figure 4.3.</b>	UV-Vis spectra recorded on the simultaneous formation of Au-Pd from $K_2PdCl_4$ (20%) and $HAuCl_4$ (80%): the variation of absorbance as the function of wavelength on subsequent addition of $K_2PdCl_4$ (a), $HAuCl_4$ (b), 3-APTMS (c), THF-HPO (d), curve-e shows the absorbance recorded after 1 h from subsequent addition and curve-f shows the spectra recorded after simultaneous formation of Au-Pd.	<b>100</b>

<b>Figure 4.4.</b>	UV-Vis spectra recorded on the sequential formation of Pd-Au from $K_2PdCl_4$ (80%) and $HAuCl_4$ (20%): the variation of absorbance of $K_2PdCl_4$ as a function of wavelength (a) followed by subsequent addition of 3-APTMS (b), THF-HPO (c), curve-d shows the spectra recorded after 1 h, curve-e shows the absorption after subsequent addition of $HAuCl_4$ followed by addition of 3-APTMS (f), curve-g shows the spectra recorded after sequential formation of Pd-Au.	<b>100</b>
<b>Figure 4.5.</b>	UV-Vis spectra recorded on the sequential formation of Au-Pd from $K_2PdCl_4$ (20%) and $HAuCl_4$ (80%): the variation of absorbance of $K_2PdCl_4$ as a function of wavelength (a) followed by subsequent addition of 3-APTMS (b), THF-HPO (c), curve-d shows the spectra recorded after 1 h, curve-e shows the absorption after subsequent addition of $HAuCl_4$ followed by addition of 3-APTMS (f), curve-g shows the spectra recorded after sequential formation of Au-Pd.	<b>101</b>
<b>Figure 4.6.</b>	SAED pattern of simultaneously synthesized Pd-Au.	<b>101</b>
<b>Figure 4.7.</b>	SAED pattern of simultaneously synthesized Au-Pd.	<b>102</b>
<b>Figure 4.8.</b>	SAED pattern of sequentially synthesized Pd-Au.	<b>102</b>
<b>Figure 4.9.</b>	SAED pattern of sequentially synthesized Au-Pd.	<b>103</b>
<b>Figure 4.10.</b>	TEM image of simultaneously synthesized Pd-Au.	<b>103</b>
<b>Figure 4.11.</b>	TEM image of simultaneously synthesized Au-Pd.	<b>104</b>
<b>Figure 4.12.</b>	TEM image of sequentially synthesized Pd-Au.	<b>104</b>
<b>Figure 4.13.</b>	TEM image of sequentially synthesized Au-Pd.	<b>105</b>
<b>Figure 4.14.</b>	The visual photographs of Ag-Au and Au-Ag.	<b>106</b>
<b>Figure 4.15.</b>	The UV-Vis spectra of the corresponding Ag-Au and Au-Ag and the insets shows the UV-Vis spectra of monometallic AgNPs and AuNPs.	<b>106</b>
<b>Figure 4.16.</b>	(a) TEM image of Au-Ag, (b) particle size distribution.	<b>107</b>
<b>Figure 4.17.</b>	SAED pattern of Au-Ag.	<b>107</b>
<b>Figure 4.18.</b>	UV-Vis spectra of (a) AgNP, (b) PBNP and (c) PBNP-AgNP; the inset shows the photographs of the corresponding system.	<b>108</b>
<b>Figure 4.19.</b>	(a) TEM image of PBNP-AgNP, (b) particle size distribution.	<b>109</b>
<b>Figure 4.20.</b>	SAED pattern of PBNP-AgNP.	<b>109</b>

<b>Figure 4.21.</b>	<b>(a)</b> TEM image of PBNP-AgNP (I), <b>(b)</b> particle size distribution.	<b>110</b>
<b>Figure 5A.1.</b>	Photographs showing the production of coloured product upon the addition of PBNP-Au-AgNPs to <i>o</i> -dianisidine in phosphate buffer pH (7.0).	<b>118</b>
<b>Figure 5A.2.</b>	Typical UV-Vis spectra of <i>o</i> -dianisidine-H <sub>2</sub> O <sub>2</sub> -PBNP-Au-AgNP reaction system.	<b>119</b>
<b>Figure 5A.3.</b>	Time-dependent absorbance changes at 430 nm in the presence of PBNP-Au-AgNP nanoparticles.	<b>119</b>
<b>Figure 5A.4.</b>	The photographs showing colour evaluation upon addition of <i>o</i> -dianisidine by varying concentrations of H <sub>2</sub> O <sub>2</sub> and constant PdNPs.	<b>120</b>
<b>Figure 5A.5.</b>	Represent the UV-Vis spectra of <i>o</i> -dianisidine oxidation products on the addition of PdNP <sub>1</sub> with various concentrations of H <sub>2</sub> O <sub>2</sub> (0.1, 0.2, 0.4, 0.8, 1.61, 3.23, 6.46, 25.84, 51.67x 10 <sup>-3</sup> M).	<b>121</b>
<b>Figure 5A.6.</b>	Represent the UV-vis spectra of <i>o</i> -dianisidine oxidation products on the addition of PdNP <sub>2</sub> with various concentrations of H <sub>2</sub> O <sub>2</sub> (0.1, 0.2, 0.4, 0.8, 1.61, 3.23, 6.46, 25.84, 51.67x 10 <sup>-3</sup> M).	<b>121</b>
<b>Figure 5A.7.</b>	Represent the UV-vis spectra of <i>o</i> -dianisidine oxidation products on the addition of PdNP <sub>3</sub> with various concentrations of H <sub>2</sub> O <sub>2</sub> (0.1, 0.2, 0.4, 0.8, 1.61, 3.23, 6.46, 25.84, 51.67x 10 <sup>-3</sup> M).	<b>122</b>
<b>Figure 5A.8.</b>	Represent the UV-vis spectra of <i>o</i> -dianisidine oxidation products on the addition of PdNP <sub>4</sub> with various concentrations of H <sub>2</sub> O <sub>2</sub> (0.1, 0.2, 0.4, 0.8, 1.61, 3.23, 6.46, 25.84, 51.67x 10 <sup>-3</sup> M).	<b>122</b>
<b>Figure 5A.9.</b>	<b>(a)</b> Kinetic analysis of PBNP-Au-Ag with H <sub>2</sub> O <sub>2</sub> as substrate; and <b>(b)</b> kinetic analysis of <i>o</i> -dianisidine as substrate.	<b>124</b>
<b>Figure 5A.10.</b>	Kinetic analysis of PBNP-AgNP (I) with H <sub>2</sub> O <sub>2</sub> as substrate.	<b>125</b>
<b>Figure 5A.11.</b>	<b>(a)</b> Kinetic analysis of PBNP-AgNP <sub>2</sub> and <b>(b)</b> PBNP-AgNP <sub>1</sub> with H <sub>2</sub> O <sub>2</sub> as substrate.	<b>125</b>
<b>Figure 5A.12.</b>	Steady-state kinetic assay of PdNP <sub>1</sub> (a) and PdNP <sub>2</sub> (b).	<b>127</b>
<b>Figure 5A.13.</b>	Steady-state kinetic assay of PdNP <sub>3</sub> (a) and PdNP <sub>4</sub> (b).	<b>127</b>
<b>Figure 5B.1.</b>	The electrochemical behavior of nanocomposite PBNP-AgNP (I) between -0.2 to 1 V vs. Ag/AgCl in 0.1 M KNO <sub>3</sub> at scan rate of 0.01 V s <sup>-1</sup> .	<b>134</b>
<b>Figure 5B.2.</b>	Cyclic Voltammograms of PBNPs (a) and PBNP-AgNP	<b>135</b>



	(I) (b), between -0.2 to 0.5 V vs. Ag/AgCl in 0.1 M KNO <sub>3</sub> at scan rate of 0.01 V s <sup>-1</sup> .	
<b>Figure 5B.3.</b>	Cyclic Voltammograms of PBNPs (a) and PBNP-AgNP (I) in absence (1) and the presence (2) of 5 mM H <sub>2</sub> O <sub>2</sub> at the scan rate of 0.01 V s <sup>-1</sup> in 0.1 M phosphate buffer (pH 7.0) containing 0.5 M KCl.	<b>135</b>
<b>Figure 5B.4.</b>	UV-Vis spectra showing the interaction between constant concentration of Fc-COOH (1) and 3-APTMS (2) in Phosphate Buffer (A) and Methanol (B).	<b>137</b>
<b>Figure 5B.5.</b>	UV-Vis spectra showing the interaction between constant concentration of Fc-COOH (1) and 3-APTMS (2) in Acetonitrile (A) and Toluene (B).	<b>137</b>
<b>Figure 5B.6.</b>	Cyclic voltammograms of Fc-COOH in 0.1 M Phosphate Buffer (pH-7.0) at various scan rates (Vs <sup>-1</sup> ); 1 = 0.01, 2 = 0.015, 3 = 0.035, 4 = 0.05, 5 = 0.07, 6 = 0.1, 7 = 0.15, 8 = 0.25, 9 = 0.35 and 10 = 0.5;	<b>138</b>
<b>Figure 5B.7.</b>	Cyclic voltammograms of FcCOOH in absence (1) and the presence of 10 mM 3-APTMS (2) and 40 mM 3-APTMS (3) at 0.01 Vs <sup>-1</sup> in 0.1 M Phosphate Buffer (pH-7.0).	<b>138</b>
<b>Figure 5B.8.</b>	Cyclic voltammograms of Fc-COOH-PdNP <sub>1</sub> (a) in 0.1 M KCl at various scan rates between 0.01Vs <sup>-1</sup> to 0.5 Vs <sup>-1</sup> ; (b) show corresponding plot of peak current density versus scan rate respectively.	<b>139</b>
<b>Figure 5B.9.</b>	Cyclic voltammograms of Fc- COOH-PdNP <sub>2</sub> (a) in 0.1 M KCl at various scan rates between 0.01Vs <sup>-1</sup> to 0.5 Vs <sup>-1</sup> ; (b) show corresponding plot of peak current density versus scan rate respectively.	<b>139</b>
<b>Figure 5B.10.</b>	Cyclic voltammograms of Fc- COOH-PdNP <sub>2</sub> (a) in 0.1 M KCl at various scan rates between 0.01Vs <sup>-1</sup> to 0.5 Vs <sup>-1</sup> ; (b) show corresponding plot of peak current density versus scan rate respectively.	<b>140</b>
<b>Figure 5B.11.</b>	Cyclic voltammograms of (a) Fc-COOH, (b) Fc-COOH-PdNP <sub>5</sub> and Fc-COOH-PdNP <sub>6</sub> systems in absence and the presence of 1 mM ascorbic acid at the scan rate of 0.01Vs <sup>-1</sup> in 0.1 M phosphate buffer (pH 7.0) containing 0.5 M KCl.	<b>141</b>
<b>Figure 5B.12.</b>	Amperometric response of Fc-COOH (a), Fc-COOH-PdNP <sub>5</sub> (b) and Fc-COOH-PdNP <sub>6</sub> , modified electrodes in 0.1 M phosphate buffer (pH 7.0) containing 0.5 M KCl on the addition of varying concentrations of ascorbic acid at 250 mV vs Ag/AgCl.	<b>142</b>