

---

---

## LIST OF FIGURES

---

---

	<b>Page No.</b>
<b>Chapter 1</b>	<b>Introduction and Literature Review</b>
Figure 1.1	Crystal structure of Fe <sub>3</sub> O <sub>4</sub> . 4
Figure 1.2	The single domain size ( $D_{crit}$ ) and superparamagnetic limit ( $D_{sp}$ ) for various ferro- or ferrimagnetic materials at room temperature. 5
Figure 1.4.1	Schematic diagram of the experimental setup for magnetic hyperthermia application. 12
<b>Chapter 3</b>	<b>Synthesis and Characterization of Zr- or Hf-substituted Magnetite Nanoparticles.</b>
Figure 3.2.1	XRD patterns for Zr <sub>x</sub> Fe <sub>3-x</sub> O <sub>4</sub> samples (a) $x=0.01$ , (b) $x=0.02$ , (c) $x=0.04$ , (d) $x=0.06$ and (e) $x=0.09$ respectively. 33
Figure 3.2.2	TEM micrographs for Zr <sub>x</sub> Fe <sub>3-x</sub> O <sub>4</sub> samples (a) $x=0.01$ and (b) $x=0.06$ . The insets show the corresponding SAD patterns for the samples. 33
Figure 3.2.3	The variation of $M_s$ , $M_r$ and $H_c$ for Zr <sub>x</sub> Fe <sub>3-x</sub> O <sub>4</sub> ( $0.01 \leq x \leq 1.0$ ) samples at room temperature and fields up to $\pm 2$ T. 35
Figure 3.2.4	The $M$ vs. $T$ curves at 50 mT for the Zr <sub>x</sub> Fe <sub>3-x</sub> O <sub>4</sub> ( $0.01 \leq x \leq 0.6$ ) samples. 35
Figure 3.2.5	The temperature vs. time curves for all the ferrofluids based on Zr <sub>x</sub> Fe <sub>3-x</sub> O <sub>4</sub> ( $0.01 \leq x \leq 1.0$ ) nanoparticles in an AC field with an amplitude of 25 mT and frequency of 112 kHz. 36

Figure 3.2.6	The temperatures achieved during magnetic hyperthermia vs. Zr concentration at different field strengths and frequencies for $Zr_xFe_{3-x}O_4$ , ( $0.01 \leq x \leq 1.0$ ) samples.	38
Figure 3.2.7	The temperature vs. time curve under a field with amplitude of 23 mT and a frequency of 173 kHz with different concentrations of the MNPs (e.g. 5, 10, 20 and 40 mg mL <sup>-1</sup> ) for samples (a) $x = 0.01$ and (b) $x = 0.5$ .	39
Figure 3.2.8	The SAR values for the $Zr_xFe_{3-x}O_4$ , $0.01 \leq x \leq 1.0$ samples at different field strengths and frequencies.	41
Figure 3.3.1	XRD patterns of $Hf_xFe_{3-x}O_4$ ( $x = 0.01, 0.06, 0.2, 0.4, 0.6$ and $0.8$ ) samples recorded at room temperature.	42
Figure 3.3.2	Variation in the lattice parameters of $Hf_xFe_{3-x}O_4$ ( $0.01 \leq x \leq 0.8$ ) with Hf-substitution.	43
Figure 3.3.3	TEM micrograph in bright field for a) $Hf_{0.06}Fe_{2.95}O_4$ and b) $Hf_{0.2}Fe_{2.8}O_4$ samples. Inset shows SAD pattern.	44
Figure 3.3.4	XPS spectra for (a) Fe 2p and (b) Hf 4d core level spectra for $Hf_{0.6}Fe_{2.4}O_4$ whereas (c) Fe 2p and (d) Hf 4d core level spectra for $Hf_{0.8}Fe_{2.2}O_4$ sample.	45
Figure 3.3.5	M vs. H curves for $Hf_xFe_{3-x}O_4$ ( $0.06 \leq x \leq 0.8$ ) samples at 300 K and $\pm 1.8$ T.	46
Figure 3.3.6	Variation of $M_S$ , $M_r$ and $H_C$ values with Hf concentration for $Hf_xFe_{3-x}O_4$ ( $0.01 \leq x \leq 0.8$ ) samples.	46
Figure 3.3.7	Magnetization vs. Temperature curves (field cooled data) for $Hf_xFe_{3-x}O_4$ (for $x = 0.06, 0.2$ and $0.8$ ) at 10 mT .	48

Figure 3.3.8	Room temperature Mössbauer spectra of $\text{Hf}_x\text{Fe}_{3-x}\text{O}_4$ ( $x = 0.06, 0.4$ and $0.6$ ) samples.	49
Figure 3.3.9	Variation in (a) hyperfine field (BHF), (b) isomer shift (d), and (c) quadrupole splitting (D) with Hf substitutions in $\text{Hf}_x\text{Fe}_{3-x}\text{O}_4$ samples (for $x= 0.06, 0.4$ and $0.6$ ).	51
Figure 3.3.10	Temperature vs. time curves for $\text{Hf}_x\text{Fe}_{3-x}\text{O}_4$ ( $0.06 \leq x \leq 0.8$ ) samples at a Field of amplitude 23 mT and a frequency 173 kHz.	54
Figure 3.3.11	Stable temperature vs. Hf-concentration plot for $\text{Hf}_x\text{Fe}_{3-x}\text{O}_4$ ( $0.01 \leq x \leq 0.8$ ) samples at different Field and frequency. Points lying in 42 - 46 °C are indicated by arrow for clarity.	55
Figure 3.3.12	The SAR values vs. Hf substitution for $\text{Hf}_x\text{Fe}_{3-x}\text{O}_4$ ( $0.01 < x \leq 0.8$ ) samples at different frequencies and fields	56

**Chapter4      Synthesis and Characterization of Al-substituted  
Magnetite Nanoparticles.**

Figure 4.2.1	(a) XRD patterns for $\text{Al}_x\text{Fe}_{3-x}\text{O}_4$ ( $x = 0.01, 0.07, 0.1, 0.5, 0.7$ and $1.0$ ) samples and (b) lattice parameter variation with Al content.	60
Figure 4.2.2	The variation of crystallite size with Al content for $\text{Al}_x\text{Fe}_{3-x}\text{O}_4$ ( $x = 0.01, 0.07, 0.1, 0.5, 0.7$ and $1.0$ ) samples.	61
Figure 4.2.3	TEM micrograph and histogram of particle size distribution for Al substituted (a, b) $\text{Al}_{0.07}\text{Fe}_{2.93}\text{O}_4$ sample (Inset shows corresponding SAD pattern) and (c) TEM micrograph for Al substituted $\text{Al}_{0.7}\text{Fe}_{2.3}\text{O}_4$ sample (Inset shows corresponding SAD pattern).	61

Figure 4.2.4	XPS spectra of $\text{Al}_{0.7}\text{Fe}_{2.3}\text{O}_4$ sample (a) Al 2p (b) Fe 2p and(c) O1s core level.	62
Figure 4.2.5	(a) Magnetization <i>vs.</i> magnetic field curves at 300 K and $\pm 2.0$ T and (b) variation of $M_S$ , $M_r$ and $H_C$ values for $\text{Al}_x\text{Fe}_{3-x}\text{O}_4$ ( $0.01 \leq x \leq 1.0$ ) samples.	63
Figure 4.2.6	(a) Temperature <i>vs.</i> time curves at a field of amplitude of 9 mT and a frequency of 640 kHz and (b) Temperature obtained during magnetic hyperthermia <i>vs.</i> Al-concentration curves at different fields and frequencies for $\text{Al}_x\text{Fe}_{3-x}\text{O}_4$ ( $0.01 \leq x \leq 1.0$ ) samples.	65
Figure 4.2.7	The SAR values <i>vs.</i> Al substitution of $\text{Al}_x\text{Fe}_{3-x}\text{O}_4$ ( $0.01 \leq x \leq 1.0$ ) samples at different frequencies and fields.	66
<b>Chapter 5</b>	<b>Synthesis and Characterization of Zn-substituted Magnetite Nanoparticles.</b>	
Figure 5.2.1	XRD patterns of $\text{Zn}_x\text{Fe}_{3-x}\text{O}_4$ ( $x = 0.01, 0.07, 0.2, 0.4$ and $0.8$ ) nanoparticles.	72
Figure 5.2.2	Variation with increased Zn content in (a) lattice parameter, (b) crystallite size.	72
Figure 5.2.3	TEM micrograph and histogram of particle size distribution for Zn substituted (a, b) $\text{Zn}_{0.2}\text{Fe}_{2.8}\text{O}_4$ and (c, d) $\text{Zn}_{0.8}\text{Fe}_{2.2}\text{O}_4$ samples (Inset shows corresponding SAD pattern).	73

Figure 5.2.4	XPS spectra of $Zn_{0.4}Fe_{2.6}O_4$ sample (a) Fe 2p (b) Zn2p and(c) O1s core level spectra.	74
Figure 5.2.5	(a) Magnetization vs. magnetic field curves at 300 K and $\pm 2.0$ T (b) variation of $M_S$ , $M_r$ and $H_C$ values and (c) $M$ vs. $T$ curves at 10 mT for $Zn_xFe_{3-x}O_4$ ( $0.01 \leq x \leq 0.8$ ) samples.	76
Figure 5.2.6	Mössbauer spectra of $Zn_xFe_{3-x}O_4$ samples ( $x = 0.01, 0.07, 0.4$ and $0.8$ ).	77
Figure 5.2.7	Variation in (a) isomer shift (b) Quadrupole splitting and (c) hyperfine field with Zn substitutions in $Zn_xFe_{3-x}O_4$ ( $x = 0.01, 0.07, 0.4$ and $0.8$ ) samples.	79
Figure 5.2.8	(a) Temperature vs. time curves at a field of amplitude of 11 mT and a frequency of 478 kHz and (b) Temperature obtained during magnetic hyperthermia vs. Zn-concentration curves at different fields and frequencies for $Zn_xFe_{3-x}O_4$ ( $0.01 \leq x \leq 0.8$ ) samples.	82
Figure 5.2.9	The SAR values vs. Zn substitution of $Zn_xFe_{3-x}O_4$ ( $0.01 \leq x \leq 0.8$ ) samples at different frequencies and fields.	83

**Chapter 6      Synthesis and Characterization of Li-substituted Magnetite Nanoparticles.**

Figure 6.2.1	X-ray diffraction patterns for $\text{Li}_x\text{Fe}_{3-x}\text{O}_4$ ( $x = 0.06, 0.1,$ and $0.3$ ) samples.	87
Figure 6.2.2	Variation with increased Li content in (a) lattice parameter, (b) crystallite size.	88
Figure 6.2.3	TEM bright field image of (a) $\text{Li}_{0.06}\text{Fe}_{2.94}\text{O}_4$ and (b) $\text{Li}_{0.3}\text{Fe}_{2.7}\text{O}_4$ samples (Inset shows corresponding SAD pattern).	89
Figure 6.2.4	XPS spectra of $\text{Li}_{0.3}\text{Fe}_{2.7}\text{O}_4$ sample (a) Fe 2p (b) Li 1s and(c) O1s core level spectra.	90
Figure 6.2.5	Room temperature magnetization vs. field curves for $\text{Li}_x\text{Fe}_{3-x}\text{O}_4$ ( $x = 0.06, 0.1$ and $0.3$ ) samples.	91
Figure 6.2.6	Mössbauer spectra of $\text{Li}_x\text{Fe}_{3-x}\text{O}_4$ samples ( $x = 0.06, 0.1$ and $0.3$ ).	92
Figure 6.2.7	Variation in (a) hyperfine field, (b) isomer shift and (c) quadrupole splitting with Li substitutions in $\text{Li}_x\text{Fe}_{3-x}\text{O}_4$ samples ( $x = 0.06, 0.1$ and $0.3$ ).	93

- Figure 6.2.8 (a)  $T$  vs.  $t$  curves at a field of amplitude of 24 mT and a frequency of 521 kHz and (b)  $T_S$  values obtained during magnetic hyperthermia at different fields and frequencies for  $\text{Li}_x\text{Fe}_{3-x}\text{O}_4$  ( $x= 0.06, 0.1$  and  $0.3$ ) samples. 96
- Figure 6.2.9 The SAR values vs. Li concentration of  $\text{Li}_x\text{Fe}_{3-x}\text{O}_4$  ( $x= 0.06, 0.1$  and  $0.3$ ) samples at different field and frequencies. 97