
LIST OF TABLES

Chapter 2	Experimental Details	
Table 2.1	Specifications for the materials utilized.	23
Chapter 3	Synthesis and Characterization of Zr- or Hf-substituted Magnetite Nanoparticles.	
Table 3.3.2	The values of hyperfine field (BHF), isomer shift (δ), quadrupole splitting (Δ), linewidth (I) and relative areas (R_A) in percentage of tetrahedral A (Fe^{3+}) and octahedral B sites of (Fe^{3+} , Fe^{2+}) ions for $\text{Hf}_x\text{Fe}_{3-x}\text{O}_4$ ($x = 0.06, 0.4$ and 0.6) samples derived from Mössbauer spectra recorded at room temperature.	51
Chapter 5	Synthesis and Characterization of Zn-substituted Magnetite Nanoparticles.	
Table 5.2.2	The values of hyperfine field (BHF), isomer shift (δ), quadrupole splitting (Δ), linewidth (I) and relative areas (R_A) in percentage of tetrahedral A (Fe^{3+} , Fe^{3+}) and octahedral B sites of (Fe^{3+} , Fe^{2+}) ions for $\text{Zn}_x\text{Fe}_{3-x}\text{O}_4$ ($x = 0.01, 0.07, 0.4$ and 0.8) samples derived from Mössbauer spectra recorded at room temperature.	80
Chapter 6	Synthesis and Characterization of Li-substituted Magnetite Nanoparticles.	

Table 6.2.1	The values of hyperfine field (B_{HF}), isomer shift (δ), quadrupole splitting (Δ), inewidth (Γ) and relative areas (R_A) in percentage of tetrahedral A (Fe^{3+}) and octahedral B sites of (Fe^{3+} , Fe^{2+}) ions for $\text{Li}_x\text{Fe}_{3-x}\text{O}_4$ ($x = 0.06, 0.1$ and 0.3) samples derived from Mössbauer spectra recorded at room temperature.	94
Table 7.1	Temperature achieved by MNPs at different combination of field and frequency during MHT	101