## LIST OF TABLES

Table No.	<b>Description</b> Pag	e No.
2.1	Specification of MODIS surface reflectance product (MOD09A1)	38
2.2	Band specification of Landsat-5 satellite data	39
2.3	Band specification of Landsat-8 satellite data	40
2.4	Equations used for the calculation of spectral indices	42
2.5	Land surface emissivity determination using NDVI	45
3.1	Details about the satellite data acquired for the study	52
3.2	Equations used for calculation of the selected indices	55
3.3	The slope, intercept and R <sup>2</sup> determined for intercaliberation from	59
	MODIS to Landsat equivalent indices for Bikaner and Hyderabad	
3.4	The slope, intercept and R <sup>2</sup> determined for intercaliberation from	60
	MODIS to Landsat equivalent indices for Vadodara and Varanasi	
3.5	LST error statistics of original MODIS LST with reference Landsat LST	61
3.6	LST error statistics of Bikaner and Hyderabad for downscaled LST	62
	using different indices	
3.7	LST error statistics of Vadodara and Varanasi for downscaled LST	63
	using different indices	
3.8	LST error statistics of Bikaner and Hyderabad for different	65
	downscaling methods	
3.9	LST error statistics of Vadodara and Varanasi for different	65
	downscaling methods	
3.10	Similarity assessment of downscaled MODIS LST with original	66
	MODIS LST for Bikaner and Hyderabad	
3.11	Similarity assessment of downscaled MODIS LST with original	67
	MODIS LST for Vadodara and Varanasi	
4.1	Details about the satellite data acquired for the study	78
4.2	Equations used for calculation of the SI	80

4.3	RMSE and the correlation coefficient (R) obtained from the	86
	downscaled MODIS-LST and the reference-Landsat-LST for	
	Bikaner and Hyderabad cities. The combinations of spectral indices	
	with best results were reported	
4.4	RMSE and the correlation coefficient (R) obtained from the	87
	downscaled MODIS-LST and the reference-Landsat-LST for	
	Vadodara and Varanasi cities. The combinations of spectral indices	
	with best results were reported	
4.5	The ratio of the RMSE to the SD determined from the downscaled	88
	MODIS-LST image and the reference-Landsat-LST image for best	
	combination. The SD for each reference-Landsat-LST was also	
	reported	
5.1	Datasets used in the study	101
5.2	The correlation of diurnal LST with combination of variables in	113
	Ahmedabad and Gandhinagar obtained from Landsat data for	
	post-monsoon season $(31/10/2015)$ for day time and $01/11/2015$	
	for night time)	
5.3	The correlation of diurnal LST with combination of variables in	116
	Ahmedabad for different seasons i.e. summer, post-monsoon and winter	
5.4	. The correlation of diurnal LST with combination of variables in	117
	Gandhinagar for different seasons i.e. summer, post-monsoon and winter	
6.1	Satellite data specification	126
6.2	Land Cover Contribution Index (LCCI) for each land covers for	136
	the years 1989, 1997, 2008 and 2018	
7.1	Satellite data specification	145
7.2	Description of five thermal levels	147
7.3	Area within regions showing SUHI effect in winter season for	151
	the years 2001, 2007, 2013 and 2019	
7.4	Area within regions showing SUHI effect in summer season for	152
	the years 2001, 2007, 2013 and 2019	

7.5	Area within regions showings SUHI effect in post-monsoon	153
	season for years 2001, 2007, 2013 and 2019	
7.6	Mean UHI value of regions 4 and 5 for winter, summer and	154
	post-monsoon season of the years 2001, 2007, 2013 and 2019	