

List of Tables

Table No.	Description	Page No.
2.1	Phases of healing of a full thickness wound	16
2.2	Factors affecting wound healing	17
2.3	Brief description of the most commonly used methods for nanofiber production	22
2.4	Approximate amino acid composition of a typical Gelatin	42
4.1	List of chemicals utilized during study	49
4.2	List of major equipment utilized during study	50
4.3	List of software utilized during study	50
4.4	Excipient composition of different PCL-based electrospinning solution	58
4.5	Electrospinning parameters for PCL-based nanofibers	59
4.6	Excipient composition of different PCL-GE-based electrospinning solution	60
4.7	Electrospinning parameters for PCL-GE-based nanofibers	60
4.8	Excipient composition of different PLGA-GE-based electrospinning solution	62
4.9	Electrospinning parameters for PLGA-GE-based nanofibers	62
5.1	λ_{\max} value, concentration range, regression equation and coefficient, and molar absorptivity of ciprofloxacin hydrochloride and quercetin	77
5.2	Accuracy of developed method	78
5.3	Intraday and inter day precision of developed method	79
5.4	Detection and quantification limit of drugs	79
5.5	Solubility of ciprofloxacin hydrochloride and quercetin in phosphate buffer (7.4)	80
5.6	Morphology, porosity and entrapment efficiency of different PCL based nanofibers.	81
5.7	<i>In-vitro</i> drug release data of PCL based nanofiber membrane in phosphate buffer (pH 7.4).	89
5.8	Diameter of <i>S. aureus</i> inhibition zone on agar plate after incubation with different PCL based nanofiber membranes.	90
5.9	Morphology, porosity and entrapment efficiency of different PCL-GE based nanofibers.	102
5.10	<i>In-vitro</i> drug release data of PCL-GE based nanofiber membrane in phosphate buffer (pH 7.4)	109
5.11	Diameter of <i>S. aureus</i> inhibition zone on agar plate after incubation with different PCL-GE based nanofiber membranes.	111
5.12	Morphology, porosity and entrapment efficiency of different PLGA-GE based nanofibers.	120

Table No.	Description	Page No.
5.13	<i>In-vitro</i> drug release data of PLGA-GE based nanofiber membrane in phosphate buffer (pH 7.4)	127
5.14	Diameter of <i>S. aureus</i> inhibition zone on agar plate after incubation with different PLGA-GE based nanofiber membranes.	129
6.1	Comparative analysis of <i>in-vitro</i> and <i>in-vivo</i> observations for three different types of nanofiber membranes	144