

LIST OF FIGURES

Fig. 2.1: Pathophysiology of ischemic stroke and mechanism of neuronal cell death	10
Fig. 2.2: Ischemic core and penumbra in a TTC stained coronal brain section of rat	11
Fig. 2.3: Types of cerebral ischemia. BCCAO: Bilateral Common Carotid Artery Occlusion; MCAO: Middle Cerebral Artery Occlusion.	14
Fig. 2.4: Arterial system of rat brain	16
Fig. 2.5: Bilateral common carotid artery occlusion	17
Fig. 2.6: Chemical structure of CGA (5-O-caffeoylquinic acid; 5-CQA)	20
Fig. 3.1 NMDARs mediated excitotoxicity resulting neuronal cell death.	25
Fig. 3.2 Rounded ribbon structure of NMDA amino-terminal domains (ATD) having two subunits: GluN1b (light sea green) and GluN2B (cornflower blue), in complex with Ifenprodil	26
Fig. 3.3 Scheme of the two-step oxygenation of arginine catalyzed by NOSes	27
Fig. 3.4 Neurodegeneration through MMP-2 and MMP-9 activation.	30
Figure: 3.5 Rounded ribbon heme domain structures of three isoforms of human nitric oxide synthase (Chain B) downloaded from PDB and used for the present study.	32
Figure 3.6: MMP-2 and MMP-9 catalytic domain with inhibitors.	33
Fig. 3.7: Binding analysis of Ifenprodil and chlorogenic acid at the heterodimer interface of the NMDAR's ATD.	47
Fig. 3.8 Binding analysis of hydroxamic acid and chlorogenic acid at the S1'-pocket of the catalytic domain of MMP-2	50

Fig. 3.9 Binding analysis of reverse hydroxamic acid and chlorogenic acid at the S1'-pocket of the catalytic domain of MMP-9	52
Figure: 3.10 Hydrogen bonding and hydrophobic interactions of Chlorogenic Acid with (A) nNOS (B) eNOS and (C) iNOS.	53
Fig. 4.1: Typical chromatogram of (A) Blank plasma and (B) Chlorogenic acid (Lowest QC concentration, 0.05 µg/ml) spiked in blank plasma at 327nm and 2.5 ml/min flow rate.	63
Fig. 4.2 Mean plasma concentration–time plot of CGA after IN and IV administration of 10 mg/kg doses in rats.	65
Fig. 4.3 Brain Penetration data of CGA following intravenous and intranasal administration at single doses (10mg/kg); (A) Brain, (B) CSF, (C) Systemic Plasma and (D) C_b/C_p	67
Fig. 4.4: HPLC peaks of brain sample after 30 minutes of CGA administration via IV and IN route	69
Fig. 4.5 Summary of PK and brain uptake analysis of CGA at single dose (10 mg/kg)	70
Fig. 5.1: Blank Hippocampi of normal rat spiked with Nitrate and Nitrite (5 µg/ml)	78
Fig. 5.2: HPLC chromatograms of nitrate in (A) Cortex; (B) Cerebellum; (C) Hippocampi and (D) CSF of normal rat.	82
Fig. 6.1: Animal preparation for bio-impedance and blood-flow measurement	89
Fig. 6.2: Equivalent circuit model of rat brain (a) Hayden Model; (b) modified Hayden Model	92
Fig. 6.3: Regional cerebral blood flow and bio-impedance of rat brain	95

Fig 6.4: Concentrations of Sodium, Chloride, Calcium and Potassium of whole rat brain in normal, occlusion and reperfusion conditions	97
Fig 6.5: Impedance of aCSF and aCSF solutions containing 2X and 4X concentration of sodium, calcium and potassium ions at 100, 500, 1k, 5k and 10k Hz frequencies	98
Fig.6.6: Bio-impedance model parameters for rat brain in different conditions. (a) Intracellular Resistance, R_i and Extracellular Resistance, R_e . (b) CPE coefficient (Q) and Amplitude of CPE exponent (n)	99
Fig. 6.7: Bio-impedance magnitude and phase plot of rat brain.	99
Fig. 7.1: Heart rate, SBP, and rCBF of normal and hypotension induced rat	111
Fig 7.2. Visualization and quantification of cerebral infarction using TTC staining of rat's brain	112
Fig. 7.3 Visualization and quantification of Evans blue (EB) extravasation in rat's brain	113
Fig. 7.4 Morphological and cellular changes in cortex visualized using H&E staining	113
Fig. 7.5 Fluorescent microscopic image of Annexin VFITC/PI stained apoptotic brain cells	114
Fig. 8.1: Workflow diagram of experimental design	121
Fig. 8.2: Dose optimization parameters. (a) Cerebral infarction area percentage, (b) Evans blue extravasation and (c) Brain water content percentage at different doses of CGA (5, 10 and 15 mg/kg b.w.)	129
Fig. 8.3: Immunohistochemistry of iNOS, TNF- α , and caspase-3 in the cortex of rat with 10X magnifications after staining	129
Fig. 8.4: Hematoxylin and eosin (H&E) staining of cortex.	130

Fig. 8.5: Biochemical parameters. (a) Calcium level, (b) Nitrate level and (c) 131
Glutamate level in cortex, cerebellum, hippocampi, and CSF