

REFERENCES

- [1] Mathers CD, Loncar D. Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Med* 2006. doi:10.1371/journal.pmed.0030442.
- [2] Lee JM, Grabb MC, Zipfel GJ, Choi DW. Brain tissue responses to ischemia. *J Clin Invest* 2000. doi:10.1172/JCI11003.
- [3] Hao L, Zou Z, Tian H, Zhang Y, Zhou H, Liu L. Stem cell-based therapies for ischemic stroke. *Biomed Res Int* 2014. doi:10.1155/2014/468748.
- [4] Granger DN, Kvietys PR. Reperfusion injury and reactive oxygen species: The evolution of a concept. *Redox Biol* 2015. doi:10.1016/j.redox.2015.08.020.
- [5] Deb P, Sharma S, Hassan KM. Pathophysiologic mechanisms of acute ischemic stroke: An overview with emphasis on therapeutic significance beyond thrombolysis. *Pathophysiology* 2010. doi:10.1016/j.pathophys.2009.12.001.
- [6] Liu S, Feng X, Jin R, Li G. Tissue plasminogen activator-based nanothrombolysis for ischemic stroke. *Expert Opin Drug Deliv* 2018. doi:10.1080/17425247.2018.1384464.
- [7] Hacke W, Kaste M, Bluhmki E, Brozman M, Dávalos A, Guidetti D, et al. Thrombolysis with Alteplase 3 to 4.5 Hours after Acute Ischemic Stroke. *N Engl J Med* 2008. doi:10.1056/NEJMoa0804656.
- [8] Miller DJ, Simpson JR, Silver B, Silver B. Safety of Thrombolysis in Acute Ischemic Stroke: A Review of Complications, Risk Factors, and Newer Technologies. *The Neurohospitalist* 2011. doi:10.1177/1941875211408731.
- [9] Xu X, Warrington AE, Bieber AJ, Rodriguez M. Enhancing CNS Repair in neurological disease: Challenges arising from neurodegeneration and rewiring of the network. *CNS Drugs* 2011. doi:10.2165/11587830-00000000-00000.
- [10] Hu H, Sun XO, Tian F, Zhang H, Liu Q, Tan W. Neuroprotective effects of isosteviol sodium injection on acute focal cerebral ischemia in rats. *Oxid Med Cell Longev* 2016. doi:10.1155/2016/1379162.
- [11] Wang L, Liu H, Zhang L, Wang G, Zhang M, Yu Y. Neuroprotection of dexmedetomidine against cerebral ischemia-reperfusion injury in rats: Involved in inhibition of NF- κ B and inflammation response. *Biomol Ther* 2017. doi:10.4062/biomolther.2015.180.
- [12] Amantea D, Tassorelli C, Russo R, Petrelli F, Morrone LA, Bagetta G, et al. Neuroprotection by leptin in a rat model of permanent cerebral ischemia: Effects on STAT3 phosphorylation in discrete cells of the brain. *Cell Death Dis* 2011. doi:10.1038/cddis.2011.125.
- [13] Lapchak PA. Emerging Therapies: Pleiotropic Multi-target Drugs to Treat Stroke Victims. *Transl Stroke Res* 2011. doi:10.1007/s12975-011-0074-4.
- [14] Gu J, Gui Y, Chen L, Yuan G, Lu HZ, Xu X. Use of Natural Products as Chemical Library for Drug Discovery and Network Pharmacology. *PLoS One* 2013. doi:10.1371/journal.pone.0062839.
- [15] More SV, Koppula S, Kim IS, Kumar H, Kim BW, Choi DK. The role of bioactive compounds on the promotion of neurite outgrowth. *Molecules* 2012. doi:10.3390/molecules17066754.
- [16] Aguiar S, Borowski T. Neuropharmacological Review of the Nootropic Herb *Bacopa monnieri*. *Rejuvenation Res* 2013. doi:10.1089/rej.2013.1431.
- [17] Singh N, Bhalla M, de Jager P, Gilca M. An overview on Ashwagandha: A Rasayana

- (Rejuvenator) of Ayurveda. African J Tradit Complement Altern Med 2011. doi:10.4314/ajtcam.v8i5S.9.
- [18] Kavitha C, Thangamani C. Amazing bean Mucuna pruriens: A comprehensive review. J Med Plants Res 2014. doi:10.5897/JMPR2013.5036.
- [19] Lokanathan Y, Omar N, Ahmad Puz NN, Saim A, Hj Idrus R. Recent updates in neuroprotective and neuroregenerative potential of Centella asiatica. Malaysian J Med Sci 2016. doi:10.1037//0012-1649.39.1.114.
- [20] Lima JDP, Farah A, King B, De Paulis T, Martin PR. Distribution of Major Chlorogenic Acids and Related Compounds in Brazilian Green and Toasted Ilex paraguariensis (Maté) Leaves. J Agric Food Chem 2016. doi:10.1021/acs.jafc.6b00276.
- [21] Clifford MN, Jaganath IB, Ludwig IA, Crozier A. Chlorogenic acids and the acyl-quinic acids: Discovery, biosynthesis, bioavailability and bioactivity. Nat Prod Rep 2017. doi:10.1039/c7np00030h.
- [22] Manjunath MJ, Muralidhara. Standardized extract of Withania somnifera (Ashwagandha) markedly offsets rotenone-induced locomotor deficits, oxidative impairments and neurotoxicity in Drosophila melanogaster. J Food Sci Technol 2015. doi:10.1007/s13197-013-1219-0.
- [23] Singh SS, Rai SN, Birla H, Zahra W, Kumar G, Gedda MR, et al. Effect of Chlorogenic Acid Supplementation in MPTP-Intoxicated Mouse . Front Pharmacol 2018;9:757. doi:10.3389/fphar.2018.00757.
- [24] Cropley V, Croft R, Silber B, Neale C, Scholey A, Stough C, et al. Does coffee enriched with chlorogenic acids improve mood and cognition after acute administration in healthy elderly? A pilot study. Psychopharmacology (Berl) 2012. doi:10.1007/s00213-011-2395-0.
- [25] Johnston KL, Clifford MN, Morgan LM. Coffee acutely modifies gastrointestinal hormone secretion and glucose tolerance in humans: Glycemic effects of chlorogenic acid and caffeine. Am J Clin Nutr 2003. doi:10.1021/jf980737w.
- [26] Ahn EH, Kim DW, Shin MJ, Kwon SW, Kim YN, Kim D-S, et al. Chlorogenic Acid Improves Neuroprotective Effect of PEP-1-Ribosomal Protein S3 Against Ischemic Insult. Exp Neurobiol 2011. doi:10.5607/en.2011.20.4.169.
- [27] Nabavi SF, Tejada S, Setzer WN, Gortzi O, Sureda A, Braidy N, et al. Chlorogenic Acid and Mental Diseases: From Chemistry to Medicine. Curr Neuropharmacol 2017. doi:10.2174/1570159X14666160325120625.
- [28] Heitman E, Ingram DK. Cognitive and neuroprotective effects of chlorogenic acid. Nutr Neurosci 2016. doi:10.1179/1476830514Y.0000000146.
- [29] Ito H, Sun X-L, Watanabe M, Okamoto M, Hatano T. Chlorogenic Acid and Its Metabolite m -Coumaric Acid Evoke Neurite Outgrowth in Hippocampal Neuronal Cells. Biosci Biotechnol Biochem 2008. doi:10.1271/bbb.70670.
- [30] Sato Y, Itagaki S, Kurokawa T, Ogura J, Kobayashi M, Hirano T, et al. In vitro and in vivo antioxidant properties of chlorogenic acid and caffeic acid. Int J Pharm 2011. doi:10.1016/j.ijpharm.2010.09.035.
- [31] Hwang SJ, Kim Y-W, Park Y, Lee H-J, Kim K-W. Anti-inflammatory effects of chlorogenic acid in lipopolysaccharide-stimulated RAW 264.7 cells. Inflamm Res 2014. doi:10.1007/s00011-013-0674-4.
- [32] Shen W, Qi R, Zhang J, Wang Z, Wang H, Hu C, et al. Chlorogenic acid inhibits LPS-

- induced microglial activation and improves survival of dopaminergic neurons. *Brain Res Bull* 2012. doi:10.1016/j.brainresbull.2012.04.010.
- [33] Bouayed J, Rammal H, Dicko A, Younos C, Soulimani R. Chlorogenic acid, a polyphenol from *Prunus domestica* (Mirabelle), with coupled anxiolytic and antioxidant effects. *J Neurol Sci* 2007. doi:10.1016/j.jns.2007.06.028.
- [34] Han J, Miyamae Y, Shigemori H, Isoda H. Neuroprotective effect of 3,5-di-O-caffeoquinic acid on SH-SY5Y cells and senescence-accelerated-prone mice 8 through the up-regulation of phosphoglycerate kinase-1. *Neuroscience* 2010. doi:10.1016/j.neuroscience.2010.05.049.
- [35] Lardeau A, Poquet L. Phenolic acid metabolites derived from coffee consumption are unlikely to cross the blood-brain barrier. *J Pharm Biomed Anal* 2013. doi:10.1016/j.jpba.2012.12.016.
- [36] Raval AP, Liu C, Hu BR. Rat model of global cerebral ischemia: The two-vessel occlusion (2VO) model of forebrain ischemia. *Anim. Model. Acute Neurol. Inj.*, 2009. doi:10.1007/978-1-60327-185-1_7.
- [37] McBean DE, Kelly PAT. Rodent models of global cerebral ischemia: A comparison of two-vessel occlusion and four-vessel occlusion. *Gen Pharmacol* 1998. doi:10.1016/S0306-3623(97)00284-X.
- [38] Bayat M, Haghani M. Acute bilateral common carotid arteries occlusion (2VO) alone could not be a proper method for induction of ischemia in rats. *Biomed Pharmacother* 2017. doi:10.1016/j.biopha.2017.11.102.
- [39] Sanderson TH, Wider JM. 2-Vessel Occlusion/Hypotension: A Rat Model of Global Brain Ischemia. *J Vis Exp* 2013. doi:10.3791/50173.
- [40] Yang L, Zhang G, Song J, Dai M, Xu C, Dong X, et al. Ex-vivo characterization of bioimpedance spectroscopy of normal, ischemic and hemorrhagic rabbit brain tissue at frequencies from 10 Hz to 1 MHz. *Sensors (Switzerland)* 2016. doi:10.3390/s16111942.
- [41] Dowrick T, Blochet C, Holder D. In vivo bioimpedance changes during haemorrhagic and ischaemic stroke in rats: Towards 3D stroke imaging using electrical impedance tomography. *Physiol Meas* 2016. doi:10.1088/0967-3334/37/6/765.
- [42] Lingwood BE, Dunster KR, Colditz PB, Ward LC. Noninvasive measurement of cerebral bioimpedance for detection of cerebral edema in the neonatal piglet. *Brain Res* 2002. doi:10.1016/S0006-8993(02)02744-0.
- [43] Choi S, Kang SW, Lee GJ, Choi SK, Chae SJ, Park HK, et al. Real-time ischemic condition monitoring in normoglycemic and hyperglycemic rats. *Physiol Meas* 2010. doi:10.1088/0967-3334/31/3/011.
- [44] Song J, Chen R, Yang L, Zhang G, Li W, Zhao Z, et al. Electrical Impedance Changes at Different Phases of Cerebral Edema in Rats with Ischemic Brain Injury. *Biomed Res Int* 2018. doi:10.1155/2018/9765174.
- [45] Castillo J, Rama R, Dávalos A. Nitric oxide-related brain damage in acute ischemic stroke. *Stroke* 2000. doi:10.1161/01.STR.31.4.852.
- [46] Csonka C, Páli T, Bencsik P, Görbe A, Ferdinand P, Csont T. Measurement of NO in biological samples. *Br J Pharmacol* 2015. doi:10.1111/bph.12832.
- [47] Ćurčić-Jovanović M, Đukić M, Vasiljević I, Ninković M, Jovanović M. Determination of nitrate by the IE-HPLC-UV method in the brain tissues of Wistar rats poisoned with paraquat. *J Serb Chem Soc* 2007;72:347–56.

- [48] Kelm M. Nitric oxide metabolism and breakdown. *Biochim Biophys Acta - Bioenerg* 1999. doi:10.1016/S0005-2728(99)00020-1.
- [49] World Health Organization. WHO - The top 10 causes of death. 24 Maggio 2018. doi:10.1007/978-3-540-87910-7_13.
- [50] Capriotti T, Murphy T. Ischemic Stroke. *Home Healthc Now* 2016. doi:10.1097/NHH.0000000000000387.
- [51] Jauch EC. Ischemic Stroke: Practice Essentials, Background, Anatomy. *Medscape* 2015.
- [52] Andersen KK, Olsen TS, Dehlendorff C, Kammersgaard LP. Hemorrhagic and ischemic strokes compared: Stroke severity, mortality, and risk factors. *Stroke* 2009. doi:10.1161/STROKEAHA.108.540112.
- [53] Xing C, Arai K, Lo EH, Hommel M. Pathophysiologic cascades in ischemic stroke. *Int J Stroke* 2012. doi:10.1111/j.1747-4949.2012.00839.x.
- [54] Cross JL, Meloni BP, Bakker AJ, Lee S, Knuckey NW. Modes of neuronal calcium entry and homeostasis following cerebral ischemia. *Int J Alzheimers Dis* 2010. doi:10.4061/2010/316862.
- [55] Bramlett HM, Dietrich WD. Pathophysiology of cerebral ischemia and brain trauma: Similarities and differences. *J Cereb Blood Flow Metab* 2004. doi:10.1097/01.WCB.0000111614.19196.04.
- [56] Carvajal FJ, Mattison HA, Cerpa W. Role of NMDA Receptor-Mediated Glutamatergic Signaling in Chronic and Acute Neuropathologies. *Neural Plast* 2016. doi:10.1155/2016/2701526.
- [57] Vanderklish PW, Bahr BA. The pathogenic activation of calpain: A marker and mediator of cellular toxicity and disease states. *Int J Exp Pathol* 2000. doi:10.1046/j.1365-2613.2000.00169.x.
- [58] Di Meo S, Reed TT, Venditti P, Victor VM. Role of ROS and RNS Sources in Physiological and Pathological Conditions. *Oxid Med Cell Longev* 2016. doi:10.1155/2016/1245049.
- [59] Broughton BRS, Reutens DC, Sobey CG. Apoptotic mechanisms after cerebral ischemia. *Stroke* 2009. doi:10.1161/STROKEAHA.108.531632.
- [60] Christophe BR, Mehta SH, Garton ALA, Sisti J, Connolly ES. Current and future perspectives on the treatment of cerebral ischemia. *Expert Opin Pharmacother* 2017. doi:10.1080/14656566.2017.1309022.
- [61] Lees KR, Bluhmki E, von Kummer R, Brott TG, Toni D, Grotta JC, et al. Time to treatment with intravenous alteplase and outcome in stroke: an updated pooled analysis of ECASS, ATLANTIS, NINDS, and EPITHET trials. *Lancet* 2010. doi:10.1016/S0140-6736(10)60491-6.
- [62] Evans MRB, White P, Cowley P, Werring DJ. Revolution in acute ischaemic stroke care: A practical guide to mechanical thrombectomy. *Pract Neurol* 2017. doi:10.1136/practneurol-2017-001685.
- [63] Lloyd M. A randomized trial of intraarterial treatment for acute ischemic stroke. *J Emerg Med* 2015. doi:10.1016/j.jemermed.2015.02.026.
- [64] Gupta R. Thrombolysis for Acute Ischemic Stroke. *Prim. Cerebrovasc. Dis.* Second Ed., 2017. doi:10.1016/B978-0-12-803058-5.00136-3.
- [65] Rabinstein AA. Treatment of Acute Ischemic Stroke. *Contin Lifelong Learn Neurol* 2017. doi:10.1212/CON.0000000000000420.

- [66] Hass WK, Easton JD, Adams HP, Pryse-Phillips W, Molony BA, Anderson S, et al. A Randomized Trial Comparing Ticlopidine Hydrochloride with Aspirin for the Prevention of Stroke in High-Risk Patients. *N Engl J Med* 1989. doi:10.1056/NEJM198908243210804.
- [67] Sandercock PAG, Counsell C, Kane EJ. Anticoagulants for acute ischaemic stroke. *Cochrane Database Syst Rev* 2015. doi:10.1002/14651858.CD000024.pub4.
- [68] Chimowitz MI, Lynn MJ, Howlett-Smith H, Stern BJ, Hertzberg VS, Frankel MR, et al. Comparison of Warfarin and Aspirin for Symptomatic Intracranial Arterial Stenosis. *N Engl J Med* 2005. doi:10.1056/NEJMoa043033.
- [69] Berge E, Sandercock P. Anticoagulants versus antiplatelet agents for acute ischemic stroke. *Stroke* 2003. doi:10.1161/01.STR.0000074551.42683.98.
- [70] Gagliardi RJ. Neuroprotection, excitotoxicity and NMDA antagonists. *Arq Neuropsiquiatr* 2000. doi:10.1590/S0004-282X2000000300030.
- [71] Hong SC, Goto Y, Lanzino G, Soleau S, Kassell NF, Lee KS. Neuroprotection with a calpain inhibitor in a model of focal cerebral ischemia. *Stroke* 1994. doi:10.1161/01.STR.25.3.663.
- [72] Chaturvedi M, Kaczmarek L. MMP-9 inhibition: A therapeutic strategy in ischemic stroke. *Mol Neurobiol* 2014. doi:10.1007/s12035-013-8538-z.
- [73] Chen W, Hartman R, Ayer R, Marcantonio S, Kamper J, Tang J, et al. Matrix metalloproteinases inhibition provides neuroprotection against hypoxia-ischemia in the developing brain. *J Neurochem* 2009. doi:10.1111/j.1471-4159.2009.06362.x.
- [74] Van Den Tweel ERW, Peeters-Scholte CMPCD, Van Bel F, Heijnen CJ, Groenendaal F. Inhibition of nNOS and iNOS following hypoxia-ischaemia improves long-term outcome but does not influence the inflammatory response in the neonatal rat brain. *Dev. Neurosci.*, 2002. doi:10.1159/000069044.
- [75] Parmentier S, Böhme G a, Lerouet D, Damour D, Stutzmann JM, Margaill I, et al. Selective inhibition of inducible nitric oxide synthase prevents ischaemic brain injury. *Br J Pharmacol* 1999. doi:10.1038/sj.bjp.0702549.
- [76] Favie LMA, Cox AR, van den Hoogen A, Nijboer CHA, Peeters-Scholte CMPCD, van Bel F, et al. Nitric Oxide Synthase Inhibition as a Neuroprotective Strategy Following Hypoxic-Ischemic Encephalopathy: Evidence From Animal Studies. *Front Neurol* 2018. doi:10.3389/fneur.2018.00258.
- [77] Kaundal RK, Shah KK, Sharma SS. Neuroprotective effects of NU1025, a PARP inhibitor in cerebral ischemia are mediated through reduction in NAD depletion and DNA fragmentation. *Life Sci* 2006. doi:10.1016/j.lfs.2006.07.034.
- [78] Yuan J. Neuroprotective strategies targeting apoptotic and necrotic cell death for stroke. *Apoptosis* 2009. doi:10.1007/s10495-008-0304-8.
- [79] Desai A, Singh N, Raghbir R. Neuroprotective potential of the NF- κ B inhibitor peptide IKK-NBD in cerebral ischemia-reperfusion injury. *Neurochem Int* 2010. doi:10.1016/j.neuint.2010.09.006.
- [80] Wang X, Feuerstein GZ, Xu L, Wang H, Schumacher W a, Ogletree ML, et al. Inhibition of tumor necrosis factor-alpha-converting enzyme by a selective antagonist protects brain from focal ischemic injury in rats. *Mol Pharmacol* 2004. doi:10.1124/mol.65.4.890.
- [81] Fluri F, Schuhmann MK, Kleinschnitz C. Animal models of ischemic stroke and their application in clinical research. *Drug Des Devel Ther* 2015.

- doi:10.2147/DDDT.S56071.
- [82] Stankowski JN, Gupta R. Therapeutic targets for neuroprotection in acute ischemic stroke: lost in translation? *Antioxid Redox Signal* 2011. doi:10.1089/ars.2010.3292.
- [83] Ducruet AF, Grobelny BT, Zacharia BE, Hickman ZL, Yeh ML, Connolly ES. Pharmacotherapy of cerebral ischemia. *Expert Opin Pharmacother* 2009. doi:10.1517/14656560903055095.
- [84] Traystman RJ. Animal models of focal and global cerebral ischemia. *ILAR J* 2003. doi:10.1093/ilar.44.2.85.
- [85] O'Neill MJ, Clemens JA. Rodent Models of Global Cerebral Ischemia. *Curr. Protoc. Neurosci.*, 2001. doi:10.1002/0471142301.ns0905s12.
- [86] Sicard KM, Fisher M. Animal models of focal brain ischemia. *Exp Transl Stroke Med* 2009. doi:10.1186/2040-7378-1-7.
- [87] Chen J, Xu XM, Xu ZC, Zhang JH. Animal models of acute neurological injuries. 2009. doi:10.1007/978-1-60327-185-1.
- [88] Sekhon MS, Ainslie PN, Griesdale DE. Clinical pathophysiology of hypoxic ischemic brain injury after cardiac arrest: A “two-hit” model. *Crit Care* 2017. doi:10.1186/s13054-017-1670-9.
- [89] Geocadin RG, Koenig MA, Jia X, Stevens RD, Peberdy MA. Management of Brain Injury After Resuscitation From Cardiac Arrest. *Neurol Clin* 2008. doi:10.1016/j.ncl.2008.03.015.
- [90] Bacigaluppi M, Comi G, Hermann DM. Animal models of ischemic stroke. Part two: modeling cerebral ischemia. *Open Neurol J* 2010. doi:10.2174/1874205X01004020034.
- [91] Hendrikse J, van Raamt AF, van der Graaf Y, Mali WPTM, van der Grond J. Distribution of Cerebral Blood Flow in the Circle of Willis. *Radiology* 2005. doi:10.1148/radiol.2351031799.
- [92] Iwasaki Y, Ito S, Suzuki M, Nagahori T, Yamamoto T, Konno H. Forebrain ischemia induced by temporary bilateral common carotid occlusion in normotensive rats. *J Neurol Sci* 1989. doi:10.1016/0022-510X(89)90098-1.
- [93] Yanpalloewar SU, Hota D, Rai S, Kumar M, Acharya SB. Nimodipine attenuates biochemical, behavioral and histopathological alterations induced by acute transient and long-term bilateral common carotid occlusion in rats. *Pharmacol Res* 2004. doi:10.1016/j.phrs.2003.08.005.
- [94] Jia H, Zhang XM, Zhang BA, Liu Y, Li JM. Dendritic morphology of neurons in medial prefrontal cortex and hippocampus in 2VO rats. *Neurol Sci* 2012. doi:10.1007/s10072-011-0898-4.
- [95] Bayat M, Sharifi MD, Haghani M, Shabani M. Enriched environment improves synaptic plasticity and cognitive deficiency in chronic cerebral hypoperfused rats. *Brain Res Bull* 2015. doi:10.1016/j.brainresbull.2015.10.001.
- [96] Khoshnam SE, Sarkaki A, Khorsandi L, Winlow W, Badavi M, Moghaddam HF, et al. Vanillic acid attenuates effects of transient bilateral common carotid occlusion and reperfusion in rats. *Biomed Pharmacother* 2017. doi:10.1016/j.biopha.2017.10.052.
- [97] Faezi M, Bigdeli MR. Prolonged and Intermittent Bilateral Common Carotid Artery Occlusion Induces Brain Lipidome Changes in a Rat Stroke Model. *Thrita* 2016;5:e28771. doi:10.5812/thrita.28771.
- [98] Smith M-L, Auer RN, Siesjö BK. The density and distribution of ischemic brain injury

- in the rat following 2–10 min of forebrain ischemia. *Acta Neuropathol* 1984. doi:10.1007/BF00690397.
- [99] Lionta E, Spyrou G, Vassilatis D, Cournia Z. Structure-Based Virtual Screening for Drug Discovery: Principles, Applications and Recent Advances. *Curr Top Med Chem* 2014. doi:10.2174/1568026614666140929124445.
- [100] Müller I. Guidelines for the successful generation of protein-ligand complex crystals. *Acta Crystallogr Sect D Struct Biol* 2017. doi:10.1107/S2059798316020271.
- [101] Taylor RD, Jewsbury PJ, Essex JW. A review of protein-small molecule docking methods. *J Comput Aided Mol Des* 2002. doi:10.1023/A:1020155510718.
- [102] Morris GM, Lim-Wilby M. Molecular docking. *Methods Mol Biol* 2008. doi:10.1007/978-1-59745-177-2_19.
- [103] Meng X-Y, Zhang H-X, Mezei M, Cui M. Molecular docking: a powerful approach for structure-based drug discovery. *Curr Comput Aided Drug Des* 2011. doi:10.1016/j.biotechadv.2011.08.021.Secreted.
- [104] Hill AD, Reilly PJ. Scoring functions for autodock. *Methods Mol Biol* 2015. doi:10.1007/978-1-4939-2343-4_27.
- [105] Garrett M, Morris D, Goodsell D, Pique M, Lindstrom W, Huey R, Forli S, Hart W, Halliday RB, Olson AJ. AutoDock Version 4.2. Citeseer 2012. doi:10.1101/354621.
- [106] Kulkarni SK, Dhir A. Withania somnifera: An Indian ginseng. *Prog Neuro-Psychopharmacology Biol Psychiatry* 2008. doi:10.1016/j.pnpbp.2007.09.011.
- [107] Pase MP, Kean J, Sarris J, Neale C, Scholey AB, Stough C. The Cognitive-Enhancing Effects of *Bacopa monnieri*: A Systematic Review of Randomized, Controlled Human Clinical Trials. *J Altern Complement Med* 2012. doi:10.1089/acm.2011.0367.
- [108] Sethiya NK, Mishra S. Review on ethnomedicinal uses and phyto-pharmacology of memory boosting herb *Convolvulus pluricaulis* Choisy. *Aust J Med Herbal* 2010.
- [109] Patay ÉB, Bencsik T, Papp N. Phytochemical overview and medicinal importance of Coffea species from the past until now. *Asian Pac J Trop Med* 2016. doi:10.1016/j.apjtm.2016.11.008.
- [110] Manyam BV, Dhanasekaran M, Hare TA. Neuroprotective effects of the antiparkinson drug *Mucuna pruriens*. *Phyther Res* 2004. doi:10.1002/ptr.1514.
- [111] Orhan IE. *Centella asiatica* (L.) Urban: From traditional medicine to modern medicine with neuroprotective potential. *Evidence-Based Complement Altern Med* 2012. doi:10.1155/2012/946259.
- [112] Lyle N, Bhattacharyya D, Sur TK, Munshi S, Paul S, Chatterjee S, et al. Stress modulating antioxidant effect of *Nardostachys jatamansi*. *Indian J Biochem Biophys* 2009.
- [113] Mutualik M, Mutualik M. *Tinospora cordifolia*: Role in depression, cognition and memory. *Aust J Med Herbal* 2011.
- [114] Suárez-Quiroz ML, Alonso Campos A, Valerio Alfaro G, González-Ríos O, Villeneuve P, Figueroa-Espinoza MC. Isolation of green coffee chlorogenic acids using activated carbon. *J Food Compos Anal* 2014. doi:10.1016/j.jfca.2013.10.005.
- [115] Farah A, Monteiro M, Donangelo CM, Lafay S. Chlorogenic Acids from Green Coffee Extract are Highly Bioavailable in Humans. *J Nutr* 2008. doi:10.3945/jn.108.095554.
- [116] Clifford MN. Chlorogenic acids and other cinnamates - Nature, occurrence and dietary burden. *J Sci Food Agric* 1999. doi:10.1002/(SICI)1097-

- 0010(19990301)79:3<362::AID-JSFA256>3.0.CO;2-D.
- [117] Meng S, Cao J, Feng Q, Peng J, Hu Y. Roles of chlorogenic acid on regulating glucose and lipids metabolism: A review. *Evidence-Based Complement Altern Med* 2013. doi:10.1155/2013/801457.
- [118] Shi H, Dong L, Jiang J, Zhao J, Zhao G, Dang X, et al. Chlorogenic acid reduces liver inflammation and fibrosis through inhibition of toll-like receptor 4 signaling pathway. *Toxicology* 2013. doi:10.1016/j.tox.2012.10.025.
- [119] Cho AS, Jeon SM, Kim MJ, Yeo J, Seo K Il, Choi MS, et al. Chlorogenic acid exhibits anti-obesity property and improves lipid metabolism in high-fat diet-induced-obese mice. *Food Chem Toxicol* 2010. doi:10.1016/j.fct.2010.01.003.
- [120] Ong KW, Hsu A, Tan BKH. Anti-diabetic and anti-lipidemic effects of chlorogenic acid are mediated by ampk activation. *Biochem Pharmacol* 2013. doi:10.1016/j.bcp.2013.02.008.
- [121] Zhao Y, Wang J, Ballevre O, Luo H, Zhang W. Antihypertensive effects and mechanisms of chlorogenic acids. *Hypertens Res* 2012. doi:10.1038/hr.2011.195.
- [122] Wang GF, Shi LP, Ren YD, Liu QF, Liu HF, Zhang RJ, et al. Anti-hepatitis B virus activity of chlorogenic acid, quinic acid and caffeic acid in vivo and in vitro. *Antiviral Res* 2009. doi:10.1016/j.antiviral.2009.05.002.
- [123] Xu R, Kang Q, Ren J, Li Z, Xu X. Antitumor molecular mechanism of chlorogenic acid on inducting genes GSK-3 β and APC and inhibiting gene β -catenin. *J Anal Methods Chem* 2013. doi:10.1155/2013/951319.
- [124] Kwon SH, Lee HK, Kim JA, Hong SI, Kim HC, Jo TH, et al. Neuroprotective effects of chlorogenic acid on scopolamine-induced amnesia via anti-acetylcholinesterase and anti-oxidative activities in mice. *Eur J Pharmacol* 2010. doi:10.1016/j.ejphar.2010.09.001.
- [125] Singh SS, Rai SN, Birla H, Zahra W, Kumar G, Gedda MR, Tiwari N, Patnaik R, Singh RK SS. Effect of Chlorogenic Acid Supplementation in MPTP-Intoxicated Mouse. *Front Pharmacol* 2018;9. doi:10.3389/fphar.2018.00757.
- [126] Kato M, Ochiai R, Kozuma K, Sato H, Katsuragi Y. Effect of Chlorogenic Acid Intake on Cognitive Function in the Elderly: A Pilot Study. *Evidence-Based Complement Altern Med* 2018. doi:10.1155/2018/8608497.
- [127] Oboh G, Agunloye OM, Akinyemi AJ, Ademiluyi AO, Adefegha SA. Comparative study on the inhibitory effect of caffeic and chlorogenic acids on key enzymes linked to Alzheimer's disease and some pro-oxidant induced oxidative stress in rats' brain-in vitro. *Neurochem Res* 2013. doi:10.1007/s11064-012-0935-6.
- [128] Miao M, Cao L, Li R, Fang X, Miao Y. Protective effect of chlorogenic acid on the focal cerebral ischemia reperfusion rat models. *Saudi Pharm J* 2017. doi:10.1016/j.jsps.2017.04.023.
- [129] Bioelectrical impedance analysis in body composition measurement. *Nutrition* 1996. doi:10.1016/S0899-9007(97)85179-9.
- [130] Khalil SF, Mohktar MS, Ibrahim F. The theory and fundamentals of bioimpedance analysis in clinical status monitoring and diagnosis of diseases. *Sensors (Switzerland)* 2014. doi:10.3390/s140610895.
- [131] Kyle UG, Bosaeus I, De Lorenzo AD, Deurenberg P, Elia M, Gómez JM, et al. Bioelectrical impedance analysis - Part I: Review of principles and methods. *Clin Nutr* 2004. doi:10.1016/j.clnu.2004.06.004.

- [132] Zhao X, Zhuang H, Yoon SC, Dong Y, Wang W, Zhao W. Electrical impedance spectroscopy for quality assessment of meat and fish: A review on basic principles measurement methods, and recent advances. *J Food Qual* 2017. doi:10.1155/2017/6370739.
- [133] Dean DA, Ramanathan T, Machado D, Sundararajan R. Electrical impedance spectroscopy study of biological tissues. *J Electrostat* 2008. doi:10.1016/j.elstat.2007.11.005.
- [134] Ivorra A, Genescà M, Sola A, Palacios L, Villa R, Hotter G, et al. Bioimpedance dispersion width as a parameter to monitor living tissues. *Physiol. Meas.*, 2005. doi:10.1088/0967-3334/26/2/016.
- [135] Blanke ML, VanDongen AM. Activation Mechanisms of the NMDA Receptor. 2008. doi:doi:10.1201/9781420044157.ch13.
- [136] Hardingham GE. Coupling of the NMDA receptor to neuroprotective and neurodestructive events. *Biochem Soc Trans* 2009. doi:10.1042/BST0371147.
- [137] Wyllie DJA, Livesey MR, Hardingham GE. Influence of GluN2 subunit identity on NMDA receptor function. *Neuropharmacology* 2013. doi:10.1016/j.neuropharm.2013.01.016.
- [138] Liu Y, Wong TP, Aarts M, Rooyakkers A, Liu L, Lai TW, et al. NMDA Receptor Subunits Have Differential Roles in Mediating Excitotoxic Neuronal Death Both In Vitro and In Vivo. *J Neurosci* 2007. doi:10.1523/JNEUROSCI.0116-07.2007.
- [139] Kumar G, Patnaik R. Exploring neuroprotective potential of *Withania somnifera* phytochemicals by inhibition of GluN2B-containing NMDA receptors: An in silico study. *Med Hypotheses* 2016;92. doi:10.1016/j.mehy.2016.04.034.
- [140] Milnerwood AJ, Gladding CM, Pouladi MA, Kaufman AM, Hines RM, Boyd JD, et al. Early Increase in Extrasynaptic NMDA Receptor Signaling and Expression Contributes to Phenotype Onset in Huntington's Disease Mice. *Neuron* 2010. doi:10.1016/j.neuron.2010.01.008.
- [141] Ahmed I, Bose SK, Pavese N, Ramlackhansingh A, Turkheimer F, Hotton G, et al. Glutamate NMDA receptor dysregulation in Parkinson's disease with dyskinesias. *Brain* 2011. doi:10.1093/brain/awr028.
- [142] Danysz W, Parsons CG. Alzheimer's disease, β -amyloid, glutamate, NMDA receptors and memantine - Searching for the connections. *Br J Pharmacol* 2012. doi:10.1111/j.1476-5381.2012.02057.x.
- [143] Shohami E, Biegon A. Novel Approach to the Role of NMDA Receptors in Traumatic Brain Injury. *CNS Neurol Disord - Drug Targets* 2014. doi:10.2174/18715273113126660196.
- [144] Rossi S, Studer V, Moscatelli A, Motta C, Coghe G, Fenu G, et al. Opposite Roles of NMDA Receptors in Relapsing and Primary Progressive Multiple Sclerosis. *PLoS One* 2013. doi:10.1371/journal.pone.0067357.
- [145] Ghasemi M, Schachter SC. The NMDA receptor complex as a therapeutic target in epilepsy: A review. *Epilepsy Behav* 2011. doi:10.1016/j.yebeh.2011.07.024.
- [146] Ozyurt E, Graham DI, Woodruff GN, McCulloch J. Protective effect of the glutamate antagonist, MK-801 in focal cerebral ischemic in the cat. *J Cereb Blood Flow Metab* 1988. doi:10.1038/jcbfm.1988.18.
- [147] Spalloni A, Nutini M, Longone P. Role of the N-methyl-d-aspartate receptors complex in amyotrophic lateral sclerosis. *Biochim Biophys Acta - Mol Basis Dis* 2013.

doi:10.1016/j.bbadis.2012.11.013.

- [148] Venkatramana DK PB, DK V, M M, R S, Kar A. Identification of Novel Drug Leads for Receptors Implicated in Migraine from Traditional Ayurvedic Herbs Using in silico and in vitro Methods. *J Neurol Disord* 2014. doi:10.4172/2329-6895.1000185.
- [149] Hansen KB, Furukawa H, Traynelis SF. Control of Assembly and Function of Glutamate Receptors by the Amino-Terminal Domain. *Mol Pharmacol* 2010. doi:10.1124/mol.110.067157.
- [150] Koller M, Urwyler S. Novel *N*-methyl-*d*-aspartate receptor antagonists: a review of compounds patented since 2006. *Expert Opin Ther Pat* 2010. doi:10.1517/13543776.2010.533656.
- [151] Williams K. Ifenprodil discriminates subtypes of the N-methyl-D-aspartate receptor: selectivity and mechanisms at recombinant heteromeric receptors. *Mol Pharmacol* 1993. doi:7901753.
- [152] Gallagher MJ, Huang H, Pritchett DB, Lynch DR. Interactions between ifenprodil and the NR2B subunit of the N-methyl-D-aspartate receptor. *J Biol Chem* 1996. doi:10.1074/jbc.271.16.9603.
- [153] Karakas E, Simorowski N, Furukawa H. Subunit Arrangement and Phenylethanolamine Binding in GluN1/GluN2B NMDA Receptors. *Nature* 2011;475:249–53. doi:10.1038/nature10180.
- [154] Ikonomidou C, Turski L. Why did NMDA receptor antagonists fail clinical trials for stroke and traumatic brain injury? *Lancet Neurol* 2002. doi:10.1016/S1474-4422(02)00164-3.
- [155] Calabrese V, Mancuso C, Calvani M, Rizzarelli E, Butterfield DA, Giuffrida Stella AM. Nitric oxide in the central nervous system: Neuroprotection versus neurotoxicity. *Nat Rev Neurosci* 2007. doi:10.1038/nrn2214.
- [156] Ying TG, Roman LJ, Martásek P, Panda SP, Ishimura Y, Masters BSS. Oxygen metabolism by endothelial nitric-oxide synthase. *J Biol Chem* 2007. doi:10.1074/jbc.M704890200.
- [157] Förstermann U, Boissel JP, Kleinert H. Expressional control of the “constitutive” isoforms of nitric oxide synthase (NOS I and NOS III). *FASEB J* 1998. doi:9657518.
- [158] Griffith OW, Stuehr DJ. Nitric Oxide Synthases: Properties and Catalytic Mechanism. *Annu Rev Physiol* 1995. doi:10.1146/annurev.ph.57.030195.003423.
- [159] Víteček J, Lojek A, Valacchi G, Kubala L. Arginine-based inhibitors of nitric oxide synthase: Therapeutic potential and challenges. *Mediators Inflamm* 2012. doi:10.1155/2012/318087.
- [160] Lüth HJ, Münch G, Arendt T. Aberrant expression of NOS isoforms in Alzheimer’s disease is structurally related to nitrotyrosine formation. *Brain Res* 2002. doi:10.1016/S0006-8993(02)03280-8.
- [161] Eliasson MJ, Huang Z, Ferrante RJ, Sasamata M, Molliver ME, Snyder SH, et al. Neuronal nitric oxide synthase activation and peroxynitrite formation in ischemic stroke linked to neural damage. *J Neurosci* 1999. doi:10.4319/lo.2006.51.5.2131.
- [162] Tieu K, Ischiropoulos H, Przedborski S. Nitric oxide and reactive oxygen species in Parkinson’s disease. *IUBMB Life* 2003. doi:10.1080/1521654032000114320.
- [163] Gruber HJ, Bernecker C, Lechner A, Weiss S, Wallner-Blazek M, Meinitzer A, et al. Increased nitric oxide stress is associated with migraine. *Cephalgia* 2010. doi:10.1111/j.1468-2982.2009.01964.x.

- [164] Garry PS, Ezra M, Rowland MJ, Westbrook J, Pattinson KTS. The role of the nitric oxide pathway in brain injury and its treatment - From bench to bedside. *Exp Neurol* 2015. doi:10.1016/j.expneurol.2014.10.017.
- [165] Kopincová J, Púzserová A, Bernátová I. L-NAME in the cardiovascular system - Nitric oxide synthase activator? *Pharmacol Reports* 2012. doi:10.1016/S1734-1140(12)70846-0.
- [166] Iadecola C, Zhang F, Xu X. Inhibition of inducible nitric oxide synthase ameliorates cerebral ischemic damage. *Am J Physiol* 1995.
- [167] Brkic M, Balusu S, Libert C, Vandenbroucke RE. Friends or Foes: Matrix Metalloproteinases and Their Multifaceted Roles in Neurodegenerative Diseases. *Mediators Inflamm* 2015. doi:10.1155/2015/620581.
- [168] Klein T, Bischoff R. Physiology and pathophysiology of matrix metalloproteases. *Amino Acids* 2011. doi:10.1007/s00726-010-0689-x.
- [169] Deryugina EI, Ratnikov B, Monosov E, Postnova TI, DiScipio R, Smith JW, et al. MT1-MMP initiates activation of pro-MMP-2 and integrin $\alpha v\beta 3$ promotes maturation of MMP-2 in breast carcinoma cells. *Exp Cell Res* 2001. doi:10.1006/excr.2000.5118.
- [170] Shapiro S, Khodalev O, Bitterman H, Auslender R, Lahat N. Different activation forms of MMP-2 oppositely affect the fate of endothelial cells. *Am J Physiol Cell Physiol* 2010. doi:10.1152/ajpcell.00305.2009.
- [171] Devarajan P, Johnston JJ, Ginsberg SS, Van Wart HE, Berliner N. Structure and expression of neutrophil gelatinase cDNA. Identity with type IV collagenase from HT1080 cells. *J Biol Chem* 1992.
- [172] Clark AW, Krekoski CA, Bou SS, Chapman KR, Edwards DR. Increased gelatinase A (MMP-2) and gelatinase B (MMP-9) activities in human brain after focal ischemia. *Neurosci Lett* 1997. doi:10.1016/S0304-3940(97)00859-8.
- [173] Rosenberg GA, Estrada EY, Dencoff JE. Matrix metalloproteinases and TIMPs are associated with blood-brain barrier opening after reperfusion in rat brain. *Stroke* 1998. doi:10.1161/01.STR.29.10.2189.
- [174] Cunningham LA, Wetzel M, Rosenberg GA. Multiple roles for MMPs and TIMPs in cerebral ischemia. *Glia* 2005. doi:10.1002/glia.20169.
- [175] Kurzepa J, Kurzepa J, Golab P, Czerska S, Bielewicz J. The significance of matrix metalloproteinase (MMP)-2 and MMP-9 in the ischemic stroke. *Int J Neurosci* 2014. doi:10.3109/00207454.2013.872102.
- [176] Power C, Henry S, Del Bigio MR, Larsen PH, Corbett D, Imai Y, et al. Intracerebral hemorrhage induces macrophage activation and matrix metalloproteinases. *Ann Neurol* 2003. doi:10.1002/ana.10553.
- [177] Lakhan SE, Kirchgessner A, Tepper D, Leonard A. Matrix metalloproteinases and blood-brain barrier disruption in acute ischemic stroke. *Front Neurol* 2013. doi:10.3389/fneur.2013.00032.
- [178] Rao B. Recent Developments in the Design of Specific Matrix Metalloproteinase Inhibitors aided by Structural and Computational Studies. *Curr Pharm Des* 2005. doi:10.2174/1381612053382115.
- [179] Williamson EM. Scientific Basis for Ayurvedic Therapies. *J Ethnopharmacol* 2004. doi:10.1016/j.jep.2004.01.005.
- [180] Raghavan A, Shah ZA. *Withania somnifera* Improves Ischemic Stroke Outcomes by Attenuating PARP1-AIF-Mediated Caspase-Independent Apoptosis. *Mol Neurobiol*

2015. doi:10.1007/s12035-014-8907-2.

- [181] Chaudhary G, Sharma U, Jagannathan NR, Gupta YK. Evaluation of *Withania somnifera* in a middle cerebral artery occlusion model of stroke in rats. *Clin Exp Pharmacol Physiol* 2003. doi:10.1046/j.1440-1681.2003.03849.x.
- [182] Sehgal N, Gupta A, Valli RK, Joshi SD, Mills JT, Hamel E, et al. *Withania somnifera* reverses Alzheimer's disease pathology by enhancing low-density lipoprotein receptor-related protein in liver. *Proc Natl Acad Sci* 2012. doi:10.1073/pnas.1112209109.
- [183] Kulkarni SK, Akula KK, Dhir A. Effect of *withania somnifera* dunal root extract against pentylenetetrazol seizure threshold in mice: Possible involvement of GABAergic system. *Indian J Exp Biol* 2008.
- [184] Kataria H, Wadhwa R, Kaul SC, Kaur G. Water extract from the leaves of *withania somnifera* protect ra differentiated c6 and IMR-32 cells against glutamate-induced excitotoxicity. *PLoS One* 2012. doi:10.1371/journal.pone.0037080.
- [185] Kumar P, Kumar A. Possible Neuroprotective Effect of *Withania somnifera* Root Extract Against 3-Nitropropionic Acid-Induced Behavioral, Biochemical, and Mitochondrial Dysfunction in an Animal Model of Huntington's Disease. *J Med Food* 2009. doi:10.1089/jmf.2008.0028.
- [186] RajaSankar S, Manivasagam T, Sankar V, Prakash S, Muthusamy R, Krishnamurti A, et al. *Withania somnifera* root extract improves catecholamines and physiological abnormalities seen in a Parkinson's disease model mouse. *J Ethnopharmacol* 2009. doi:10.1016/j.jep.2009.08.003.
- [187] Gavande K, Jain K, Jain B, Mehta R. Comprehensive Report on Phytochemistry and Pharmacological Prominence of *Withania somnifera*. 2015. doi:10.20510/ukjpb/3/i2/89342.
- [188] Matsuda H, Murakami T, Kishi A, Yoshikawa M. Structures of withanosides I, II, III, IV, V, VI, and VII, new withanolide glycosides, from the roots of Indian *Withania somnifera* DUNAL. and inhibitory activity for tachyphylaxis to clonidine in isolated guinea-pig ileum. *Bioorganic Med Chem* 2001. doi:10.1016/S0968-0896(01)00024-4.
- [189] Duke JA. Dr. Duke's phytochemical and ethnobotanical databases. *Choice Rev Online* 2001. doi:10.1111/j.1468-3083.2008.02890.x.
- [190] Kim S, Thiessen PA, Bolton EE, Chen J, Fu G, Gindulyte A, et al. PubChem substance and compound databases. *Nucleic Acids Res* 2016. doi:10.1093/nar/gkv951.
- [191] ProteinDataBank. RCSB Protein Data Bank - RCSB PDB RCSB Protein Data Bank - RCSB PDB. *Hum Serum Albumin* 2013. doi:10.1093/nar/28.1.235.
- [192] Feng Y, Likos JJ, Zhu L, Woodward H, Munie G, McDonald JJ, et al. Solution structure and backbone dynamics of the catalytic domain of matrix metalloproteinase-2 complexed with a hydroxamic acid inhibitor. *Biochim Biophys Acta - Proteins Proteomics* 2002. doi:10.1016/S0167-4838(02)00307-2.
- [193] Rowsell S, Hawtin P, Minshull CA, Jepson H, Brockbank SMV, Barratt DG, et al. Crystal structure of human MMP9 in complex with a reverse hydroxamate inhibitor. *J Mol Biol* 2002. doi:10.1016/S0022-2836(02)00262-0.
- [194] Li H, Jamal J, Plaza C, Pineda SH, Chreifi G, Jing Q, et al. Structures of human constitutive nitric oxide synthases. *Acta Crystallogr Sect D Biol Crystallogr* 2014. doi:10.1107/S1399004714017064.
- [195] Li H, Jamal J, Delker S, Plaza C, Ji H, Jing Q, et al. The mobility of a conserved

- tyrosine residue controls isoform-dependent enzyme-inhibitor interactions in nitric oxide synthases. *Biochemistry* 2014. doi:10.1021/bi500561h.
- [196] S.K.Lee, I.H.Lee, H.J.Kim, G.S.Chang, J.E.Chung KTN. The PreADME Approach: Web-based program for rapid prediction of physico-chemical, drug absorption and drug-like properties. *EuroQSAR 2002 Des. Drugs Crop Prot. Process. Probl. Solut.*, 2003.
- [197] Jayaram B, Singh T, Mukherjee G, Mathur A, Shekhar S, Shekhar V. Sanjeevini: a freely accessible web-server for target directed lead molecule discovery. *BMC Bioinformatics* 2012. doi:10.1186/1471-2105-13-S17-S7.
- [198] Dallakyan S, Olson AJ. Small-molecule library screening by docking with PyRx. *Methods Mol Biol* 2015. doi:10.1007/978-1-4939-2269-7_19.
- [199] Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, et al. UCSF Chimera—A Visualization System for Exploratory Research and Analysis. *J Comput Chem* 2004. doi:10.1002/jcc.20084.
- [200] Wang J, Wang W, Kollman PA, Case DA. Automatic atom type and bond type perception in molecular mechanical calculations. *J Mol Graph Model* 2006. doi:10.1016/j.jmgm.2005.12.005.
- [201] Sanner MF. Python: a programming language for software integration and development. *J Mol Graph Model* 1999. doi:10.1016/S1093-3263(99)99999-0.
- [202] Lipinski CA, Lombardo F, Dominy BW, Feeney PJ. Experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings. *Adv Drug Deliv Rev* 2012. doi:10.1016/j.addr.2012.09.019.
- [203] Lipinski CA. Lead- and drug-like compounds: The rule-of-five revolution. *Drug Discov Today Technol* 2004. doi:10.1016/j.ddtec.2004.11.007.
- [204] Blake J. Identification and Evaluation of Molecular Properties Related to Preclinical Optimization and Clinical Fate. *Med Chem (Los Angeles)* 2005. doi:10.2174/157340605774598081.
- [205] Leeson P. Drug discovery: Chemical beauty contest. *Nature* 2012. doi:10.1038/481455a.
- [206] Banks WA. Characteristics of compounds that cross the blood-brain barrier. *BMC Neurol.*, 2009. doi:10.1186/1471-2377-9-S1-S3.
- [207] Wessel MD, Jurs PC, Tolan JW, Muskal SM. Prediction of human intestinal absorption of drug compounds from molecular structure. *J Chem Inf Comput Sci* 1998. doi:10.1021/ci980029a.
- [208] Ames B, Gurney E, Miller J, Bartsch H. Carcinogens as frameshift mutagens: metabolites and derivatives of 2-acetylaminofluorene and other aromatic amine carcinogens. *Proc Natl Acad Sci U S A* 1972. doi:10.1073/pnas.69.11.3128.
- [209] Karakas E, Simorowski N, Furukawa H. Subunit arrangement and phenylethanolamine binding in GluN1/GluN2B NMDA receptors. *Nature* 2011. doi:10.1038/nature10180.
- [210] Gupta SP, Patil VM. Specificity of binding with matrix metalloproteinases. *EXS* 2012. doi:10.1007/978-3-0348-0364-9_2.
- [211] Wang Y, Yu SJ, Li YX, Luo HS. Expression and clinical significance of matrix metalloproteinase-17 and -25 in gastric cancer. *Oncol Lett* 2015. doi:10.3892/ol.2014.2747.
- [212] Reinhard SM, Razak K, Ethell IM. A delicate balance: role of MMP-9 in brain development and pathophysiology of neurodevelopmental disorders. *Front Cell*

Neurosci 2015. doi:10.3389/fncel.2015.00280.

- [213] Nanri K, Montécot C, Springhetti V, Seylaz J, Pinard E, Dietrich WD. The Selective Inhibitor of Neuronal Nitric Oxide Synthase, 7-Nitroindazole, Reduces the Delayed Neuronal Damage Due to Forebrain Ischemia in Rats • Editorial Comment. *Stroke* 1998. doi:10.1161/01.STR.29.6.1248.
- [214] Olthof MR, Hollman PCH, Katan MB. Chlorogenic Acid and Caffeic Acid Are Absorbed in Humans. *J Nutr* 2001. doi:10.1093/jn/131.1.66.
- [215] Higdon JV, Frei B. Coffee and Health: A Review of Recent Human Research Coffee and Health: A Review of Recent Human Research. *Crit Rev Food Sci Nutr* 2006. doi:10.1080/10408390500400009.
- [216] Gebara KS, Gasparotto-Junior A, Santiago PG, Cardoso CAL, De Souza LM, Morand C, et al. Daily Intake of Chlorogenic Acids from Consumption of Maté (*Ilex paraguariensis* A.St.-Hil.) Traditional Beverages. *J Agric Food Chem* 2017. doi:10.1021/acs.jafc.7b04093.
- [217] Huang SM, Chuang HC, Wu CH, Yen GC. Cytoprotective effects of phenolic acids on methylglyoxal-induced apoptosis in Neuro-2A cells. *Mol Nutr Food Res* 2008. doi:10.1002/mnfr.200700360.
- [218] Lee K, Lee J-S, Jang H-J, Kim S-M, Chang MS, Park SH, et al. Chlorogenic acid ameliorates brain damage and edema by inhibiting matrix metalloproteinase-2 and 9 in a rat model of focal cerebral ischemia. *Eur J Pharmacol* 2012. doi:10.1016/j.ejphar.2012.05.028.
- [219] Azuma K, Ippoushi K, Nakayama M, Ito H, Higashio H, Terao J. Absorption of chlorogenic acid and caffeic acid in rats after oral administration. *J Agric Food Chem* 2000. doi:10.1021/jf000483q.
- [220] Jung JW, Kim JM, Jeong JS, Son M, Lee HS, Lee MG, et al. Pharmacokinetics of chlorogenic acid and corydaline in DA-9701, a new botanical gastroprotective agent, in rats. *Xenobiotica* 2014. doi:10.3109/00498254.2013.874610.
- [221] Yuan B, Qiao M, Xu H, Wang L, Li F. Determination of chlorogenic acid in rat plasma by high performance chromatography after peritoneal administration of compound Daqingye injection. *Yakugaku Zasshi* 2006. doi:JST.JSTAGE/yakushi/126.811 [pii].
- [222] Guo Z, Li J. Chlorogenic acid prevents alcohol-induced brain damage in neonatal rat. *Transl Neurosci* 2017. doi:10.1515/tnsci-2017-0024.
- [223] Taram F, Winter AN, Linseman DA. Neuroprotection comparison of chlorogenic acid and its metabolites against mechanistically distinct cell death-inducing agents in cultured cerebellar granule neurons. *Brain Res* 2016. doi:10.1016/j.brainres.2016.07.028.
- [224] Graff CL, Pollack GM. Nasal drug administration: Potential for targeted central nervous system delivery. *J Pharm Sci* 2005. doi:10.1002/jps.20318.
- [225] Illum L. Transport of drugs from the nasal cavity to the central nervous system. *Eur J Pharm Sci* 2000. doi:10.1016/S0928-0987(00)00087-7.
- [226] B. Chauhan M, B. Chauhan N. Brain Uptake of Neurotherapeutics after Intranasal Delivery in Mice. *J Neurol Neurosurg* 2015. doi:10.19104/jnn.2015.91.
- [227] Yang H, Yuan B, Li L, Chen H, Li F. HPLC Determination and Pharmacokinetics of Chlorogenic Acid in Rabbit Plasma after An Oral Dose of *Flos Lonicerae* Extract. *J Chromatogr Sci* 2004. doi:10.1093/chromsci/42.4.173.

- [228] Shrivastava A, Gupta V. Methods for the determination of limit of detection and limit of quantitation of the analytical methods. *Chronicles Young Sci* 2011. doi:10.4103/2229-5186.79345.
- [229] Nirogi R, Kandikere V, Mudigonda K, Bhyrapuneni G, Muddana N, Saralaya R, et al. A simple and rapid method to collect the cerebrospinal fluid of rats and its application for the assessment of drug penetration into the central nervous system. *J Neurosci Methods* 2009. doi:10.1016/j.jneumeth.2008.12.001.
- [230] Hanson LR, Frey WH. Intranasal delivery bypasses the blood-brain barrier to target therapeutic agents to the central nervous system and treat neurodegenerative disease. *BMC Neurosci.*, 2008. doi:10.1186/1471-2202-9-S2-S5.
- [231] Kaur P, Kim K. Pharmacokinetics and brain uptake of diazepam after intravenous and intranasal administration in rats and rabbits. *Int J Pharm* 2008. doi:10.1016/j.ijpharm.2008.07.030.
- [232] Dhuria S V., Hanson LR, Frey WH. Intranasal delivery to the central nervous system: Mechanisms and experimental considerations. *J Pharm Sci* 2010. doi:10.1002/jps.21924.
- [233] Faraci FM, Brian JE. Nitric oxide and the cerebral circulation. *Stroke* 1994. doi:10.1161/01.STR.25.3.692.
- [234] Moncada S, Palmer RMJ, Higgs EA. Nitric oxide: physiology, pathophysiology, and pharmacology. *Pharmacol Rev* 1991. doi:10.1017/CBO9781107415324.004.
- [235] Mander P, Brown GC. Nitric oxide, hypoxia and brain inflammation. *Biochem Soc Trans* 2004. doi:10.1042/BST0321068.
- [236] Stuehr DJ. Mammalian nitric oxide synthases. *Biochim Biophys Acta - Bioenerg* 1999. doi:10.1016/S0005-2728(99)00016-X.
- [237] Kumar G, Mukherjee S, Patnaik R. Identification of Withanolide-M and Stigmasterol as Potent neuroprotectant and Dual inhibitor of Inducible/Neuronal Nitric Oxide Synthase by Structure-Based Virtual Screening. *J Biol Eng Res Rev* 2017;4:09-13.
- [238] Kumar G, Paliwal P, Patnaik N, Patnaik R. Withania somnifera phytochemicals confer neuroprotection by selective inhibition of nNos: An in silico study to search potent and selective inhibitors for human nNOS. *J Theor Comput Chem* 2017. doi:10.1142/S0219633617500420.
- [239] Sun J, Zhang X, Broderick M, Fein H. Measurement of Nitric Oxide Production in Biological Systems by Using Griess Reaction Assay. *Sensors* 2003. doi:10.3390/s30800276.
- [240] Misko TP, Schilling RJ, Salvemini D, Moore WM, Currie MG. A fluorometric assay for the measurement of nitrite in biological samples. *Anal Biochem* 1993. doi:10.1006/abio.1993.1449.
- [241] Braman RS, Hendrix S a. Nanogram nitrite and nitrate determination in environmental and biological materials by vanadium (III) reduction with chemiluminescence detection. *Anal Chem* 1989. doi:10.1021/ac00199a007.
- [242] Badea M, Amine A, Palleschi G, Moscone D, Volpe G, Curulli A. New electrochemical sensors for detection of nitrites and nitrates. *J Electroanal Chem* 2001. doi:10.1016/S0022-0728(01)00358-8.
- [243] Gao L, Barber-Singh J, Kotegoda S, Wirtshafer D, Shippy SA. Determination of nitrate and nitrite in rat brain perfusates by capillary electrophoresis. *Electrophoresis* 2004. doi:10.1002/elps.200305840.

- [244] Zhao J, Wang J, Yang Y, Lu Y. The determination of nitrate and nitrite in human urine and blood by high-performance liquid chromatography and cloud-point extraction. *J Chromatogr Sci* 2015. doi:10.1093/chromsci/bmu212.
- [245] Jedlickova V, Jedlicková V, Paluch Z, Alušk Š. Determination of nitrate and nitrite by high-performance liquid chromatography in human plasma. *J Chromatogr B* 2002. doi:[http://dx.doi.org/10.1016/S1570-0232\(02\)00405-1](http://dx.doi.org/10.1016/S1570-0232(02)00405-1).
- [246] Wang X, Lin F, Gao Y, Lei H. Bilateral common carotid artery occlusion induced brain lesions in rats: A longitudinal diffusion tensor imaging study. *Magn Reson Imaging* 2015. doi:10.1016/j.mri.2015.02.010.
- [247] Kumar G, Paliwal P, Mukherjee S, Patnaik N, Krishnamurthy S, Patnaik R. Pharmacokinetics and brain penetration study of chlorogenic acid in rats. *Xenobiotica* 2018. doi:10.1080/00498254.2018.1445882.
- [248] Bryan NS, Grisham MB. Methods to detect nitric oxide and its metabolites in biological samples. *Free Radic Biol Med* 2007. doi:10.1016/j.freeradbiomed.2007.04.026.
- [249] Shrivastava A, Chaturvedi U, Sonkar R, Khanna AK, Saxena JK, Bhatia G. Antioxidant effect of Azadirachta indica on high fat diet induced diabetic charles foster rats. *Appl Biochem Biotechnol* 2012. doi:10.1007/s12010-012-9681-0.
- [250] Bhattacharya P, Pandey AK, Paul S, Patnaik R. Alleviation of glutamate mediated neuronal insult by piroxicam in rodent model of focal cerebral ischemia: a possible mechanism of GABA agonism. *J Physiol Biochem* 2014. doi:10.1007/s13105-014-0358-8.
- [251] Chanda D, Shanker K, Pal A, Luqman S, Bawankule DU, Mani D, et al. Safety evaluation of Trikatu, a generic Ayurvedic medicine in Charles Foster rats. *J Toxicol Sci* 2009. doi:10.2131/jts.34.99.
- [252] Rizzo V, Montalbetti L, Rozza AL, Bolzani W, Porta C, Balduzzi G, et al. Nitrite/nitrate balance during photoinduced cerebral ischemia in the rat determined by high-performance liquid chromatography with UV and electrochemical detection. *J Chromatogr A* 1998;798:103–8. doi:[https://doi.org/10.1016/S0021-9673\(97\)01186-2](https://doi.org/10.1016/S0021-9673(97)01186-2).
- [253] Raichle ME. The pathophysiology of brain ischemia. *Ann Neurol* 1983. doi:10.1002/ana.410130103.
- [254] Rueda CB, Llorente-Folch I, Traba J, Amigo I, Gonzalez-Sanchez P, Contreras L, et al. Glutamate excitotoxicity and Ca²⁺-regulation of respiration: Role of the Ca²⁺-activated mitochondrial transporters (CaMCs). *Biochim Biophys Acta - Bioenerg* 2016. doi:10.1016/j.bbabi.2016.04.003.
- [255] Nishizawa Y. Glutamate release and neuronal damage in ischemia. *Life Sci* 2001. doi:10.1016/S0024-3205(01)01142-0.
- [256] Zündorf G, Reiser G. Calcium Dysregulation and Homeostasis of Neural Calcium in the Molecular Mechanisms of Neurodegenerative Diseases Provide Multiple Targets for Neuroprotection. *Antioxid Redox Signal* 2011. doi:10.1089/ars.2010.3359.
- [257] Weber JT. Altered calcium signaling following traumatic brain injury. *Front Pharmacol* 2012. doi:10.3389/fphar.2012.00060.
- [258] Iwasaki Y, Ito S, Suzuki M, Nagahori T, Yamamoto T, Konno H. Forebrain ischemia induced by temporary bilateral common carotid occlusion in normotensive rats. *J Neurol Sci* 1989. doi:10.1016/0022-510X(89)90098-1.
- [259] McDermott B, Porter E, Jones M, McGinley B, O'Halloran M. Symmetry difference

- electrical impedance tomography- A novel modality for anomaly detection. *Physiol Meas* 2018. doi:10.1088/1361-6579/aab656.
- [260] Oka K, Yamamoto M, Nonaka T, Tomonaga M. The significance of artificial cerebrospinal fluid as perfusate and endoneurosurgery. *Neurosurgery* 1996. doi:10.1227/00006123-199604000-00019.
- [261] Yao H, Nabika T. Excess salt increases infarct size produced by photothrombotic distal middle cerebral artery occlusion in spontaneously hypertensive rats. *PLoS One* 2014. doi:10.1371/journal.pone.0097109.
- [262] Bera TK. Bioelectrical Impedance and the Frequency Dependent Current Conduction Through Biological Tissues: A Short Review. *IOP Conf. Ser. Mater. Sci. Eng.*, 2018. doi:10.1088/1757-899X/331/1/012005.
- [263] Lempka SF, Miocinovic S, Johnson MD, Vitek JL, McIntyre CC. In vivo impedance spectroscopy of deep brain stimulation electrodes. *J Neural Eng* 2009. doi:10.1088/1741-2560/6/4/046001.
- [264] Wei XF, Grill WM. Impedance characteristics of deep brain stimulation electrodes in vitro and in vivo. *J Neural Eng* 2009. doi:10.1088/1741-2560/6/4/046008.
- [265] Holder DS. Detection of cerebral ischaemia in the anaesthetised rat by impedance measurement with scalp electrodes: Implications for non-invasive imaging of stroke by electrical impedance tomography. *Clin Phys Physiol Meas* 1992. doi:10.1088/0143-0815/13/1/006.
- [266] Eldarrat A, High A, Kale G. Influence of sodium chloride content in electrolyte solution on electrochemical impedance measurements of human dentin. *Dent Res J (Isfahan)* 2017. doi:10.4103/1735-3327.201130.
- [267] Yu Xm, Groveman Br, Fang Xq, Lin Sx. The Role Of Intracellular Sodium (Na) In The Regulation Of Calcium (Ca)-Mediated Signaling And Toxicity. *Health (Irvine Calif)* 2010. doi:10.4236/health.2010.21002.
- [268] Traynelis SF, Wollmuth LP, McBain CJ, Menniti FS, Vance KM, Ogden KK, et al. Glutamate Receptor Ion Channels: Structure, Regulation, and Function. *Pharmacol Rev* 2010. doi:10.1124/pr.109.002451.
- [269] Terry J. The major electrolytes: sodium, potassium, and chloride. *J Intraven Nurs* 1994.
- [270] Xin WK, Kwan CL, Zhao XH, Xu J, Ellen RP, McCulloch CA, et al. A functional interaction of sodium and calcium in the regulation of NMDA receptor activity by remote NMDA receptors. *J Neurosci* 2005. doi:10.1523/JNEUROSCI.3791-04.2005.
- [271] Canazza A, Minati L, Boffano C, Parati E, Binks S. Experimental models of brain ischemia: A review of techniques, magnetic resonance imaging, and investigational cell-based therapies. *Front Neurol* 2014. doi:10.3389/fneur.2014.00019.
- [272] Casals JB, Pieri NCG, Feitosa MLT, Ercolin ACM, Roballo KCS, Barreto RSN, et al. The use of animal models for stroke research: A review. *Comp Med* 2011. doi:10.1017/CBO9781107415324.004.
- [273] Fares H, DiNicolantonio JJ, O'Keefe JH, Lavie CJ. Amlodipine in hypertension: A first-line agent with efficacy for improving blood pressure and patient outcomes. *Open Hear* 2016. doi:10.1136/openhrt-2016-000473.
- [274] Rao NS, Oommen A, Bindumathi PL, Sharma V, Rao S, Moodahadu LS, et al. Efficacy and tolerability of fixed dose combination of metoprolol and amlodipine in Indian patients with essential hypertension. *J Midlife Health* 2013;4:160–6.

doi:10.4103/0976-7800.119000.

- [275] Ichiki H, Kuroiwa T, Taniguchi I, Okeda R. Delayed recovery of cortical auditory evoked potentials and blood flow precede cortical neuronal death after transient cerebral ischemia in gerbils. *J Med Dent Sci* 1998.
- [276] Schilichting CLR, Lima KCM, Cestari LA, Sekiyama JY, Silva FM, Milani H. Validation of a simple and inexpensive method for the quantitation of infarct in the rat brain. *Brazilian J Med Biol Res* 2004. doi:10.1590/S0100-879X2004000400008.
- [277] Türeyen K, Vemuganti R, Sailor KA, Dempsey RJ. Infarct volume quantification in mouse focal cerebral ischemia: A comparison of triphenyltetrazolium chloride and cresyl violet staining techniques. *J Neurosci Methods* 2004. doi:10.1016/j.jneumeth.2004.04.029.
- [278] Martin Y, Avendaño C, Piedras MJ, Krzyzanowska A. Evaluation of Evans Blue extravasation as a measure of peripheral inflammation 2010.
- [279] Cardiff RD, Miller CH, Munn RJ. Manual hematoxylin and eosin staining of mouse tissue sections. *Cold Spring Harb Protoc* 2014. doi:10.1101/pdb.prot073411.
- [280] Dhawan J, Benveniste H, Luo Z, Nawrocky M, Smith SD, Biegon A. A new look at glutamate and ischemia: NMDA agonist improves long-term functional outcome in a rat model of stroke. *Future Neurol* 2011. doi:10.2217/fnl.11.55.
- [281] Farbiszewski R, Bielawski K, Bielawska A, Sobaniec W. Spermine protects in vivo the antioxidant enzymes in transiently hypoperfused rat brain. *Acta Neurobiol Exp (Wars)* 1995.
- [282] Moore PK, Wallace P, Gaffen Z, Hart SL, Babbedge RC. Characterization of the novel nitric oxide synthase inhibitor 7- nitro indazole and related indazoles: antinociceptive and cardiovascular effects. *Br J Pharmacol* 1993. doi:10.1111/j.1476-5381.1993.tb13795.x.
- [283] Mishra V, Verma R, Singh N, Raghbir R. The neuroprotective effects of NMDAR antagonist, ifenprodil and ASIC1a inhibitor, flurbiprofen on post-ischemic cerebral injury. *Brain Res* 2011. doi:10.1016/j.brainres.2011.03.011.
- [284] Keep RF, Hua Y, Xi G. Brain Water Content: A Misunderstood Measurement? *Transl Stroke Res* 2012. doi:10.1007/s12975-012-0152-2.
- [285] Gorbatyuk OS, Li S, Sullivan LF, Chen W, Kondrikova G, Manfredsson FP, et al. The phosphorylation state of Ser-129 in human alpha-synuclein determines neurodegeneration in a rat model of Parkinson disease. *Proc Natl Acad Sci U S A* 2008. doi:10.1073/pnas.0711053105.
- [286] Wu X, Wang R, Jiang Q, Wang S, Yao Y, Shao L. Determination of amino acid neurotransmitters in rat hippocampi by HPLC-UV using NBD-F as a derivative. *Biomed Chromatogr* 2014. doi:10.1002/bmc.3062.
- [287] Ogata J, Fujishima M, Morotomi Y, Omae T. Cerebral infarction following bilateral carotid artery ligation in normotensive and spontaneously hypertensive rats: a pathological study. *Stroke* 1976. doi:10.1161/01.STR.7.1.54.
- [288] Sadoshima S, Fujishima M, Ogata J, Ibayashi S, Shiokawa O, Omae T. Disruption of blood-brain barrier following bilateral carotid artery occlusion in spontaneously hypertensive rats. A quantitative study. *Stroke* 1983. doi:10.1161/01.STR.14.6.876.
- [289] NAGAHORI T, NISHIJIMA M, ENDO S, TAKAKU A, IWASAKI Y. Ischemic brain damage induced by repeated brief occlusions of bilateral common carotid artery in rats. *Tohoku J Exp Med* 1994;172:253–62.

- [290] Mattila OS, Strbian D, Saksi J, Pikkarainen TO, Rantanen V, Tatlisumak T, et al. Cerebral mast cells mediate blood-brain barrier disruption in acute experimental ischemic stroke through perivascular gelatinase activation. *Stroke* 2011. doi:10.1161/STROKEAHA.111.632224.
- [291] Simard JM, Kent TA, Chen M, Tarasov K V., Gerzanich V. Brain oedema in focal ischaemia: molecular pathophysiology and theoretical implications. *Lancet Neurol* 2007. doi:10.1016/S1474-4422(07)70055-8.
- [292] Zhu Y, Yang G-Y, Ahlemeyer B, Pang L, Che X-M, Culmsee C, et al. Transforming Growth Factor- β 1 Increases Bad Phosphorylation and Protects Neurons Against Damage. *J Neurosci* 2002. doi:10.1523/JNEUROSCI.22-10-03898.2002.
- [293] Liu T, Clark RK, McDonnell PC, Young PR, White RF, Barone FC, et al. Tumor necrosis factor-alpha expression in ischemic neurons. *Stroke* 1994. doi:10.1161/01.STR.25.7.1481.
- [294] Suzuki M, Tabuchi M, Ikeda M, Tomita T. Concurrent formation of peroxynitrite with the expression of inducible nitric oxide synthase in the brain during middle cerebral artery occlusion and reperfusion in rats. *Brain Res* 2002. doi:10.1016/S0006-8993(02)03145-1.
- [295] Ferrer I, Friguls B, Dalfót E, Justicia C, Planas AM. Caspase-dependent and caspase-independent signalling of apoptosis in the penumbra following middle cerebral artery occlusion in the adult rat. *Neuropathol Appl Neurobiol* 2003. doi:10.1046/j.1365-2990.2003.00485.x.
- [296] O'Mahony D, Kendall MJ. Nitric oxide in acute ischaemic stroke: A target for neuroprotection. *J Neurol Neurosurg Psychiatry* 1999. doi:10.1136/jnnp.67.1.1.
- [297] Sun Y, Xu Y, Geng L. Caspase-3 inhibitor prevents the apoptosis of brain tissue in rats with acute cerebral infarction. *Exp Ther Med* 2015. doi:10.3892/etm.2015.2462.
- [298] Tuttolomondo A, Pecoraro R, Pinto A. Studies of selective TNF inhibitors in the treatment of brain injury from stroke and trauma: a review of the evidence to date. *Drug Des Devel Ther* 2014. doi:10.2147/DDDT.S67655.
- [299] Mărgăritescu O, Mogoantă L, Pirici I, Pirici D, Cernea D, Mărgăritescu C. Histopathological changes in acute ischemic stroke. *Rom J Morphol Embryol* 2009.
- [300] Dirnagl U, Iadecola C, Moskowitz MA. Pathobiology of ischaemic stroke: An integrated view. *Trends Neurosci* 1999. doi:10.1016/S0166-2236(99)01401-0.
- [301] Wu QJ, Tymianski M. Targeting nmda receptors in stroke: New hope in neuroprotection Tim Bliss. *Mol Brain* 2018. doi:10.1186/s13041-018-0357-8.
- [302] Mikami Y, Yamazawa T. Chlorogenic acid, a polyphenol in coffee, protects neurons against glutamate neurotoxicity. *Life Sci* 2015. doi:10.1016/j.lfs.2015.08.005.
- [303] Rebai O, Belkhir M, Sanchez-Gomez MV, Matute C, Fattouch S, Amri M. Differential Molecular Targets for Neuroprotective Effect of Chlorogenic Acid and its Related Compounds Against Glutamate Induced Excitotoxicity and Oxidative Stress in Rat Cortical Neurons. *Neurochem Res* 2017. doi:10.1007/s11064-017-2403-9.
- [304] Mohammadi MT, Shid-Moosavi SM, Dehghani GA. Contribution of nitric oxide synthase (NOS) in blood-brain barrier disruption during acute focal cerebral ischemia in normal rat. *Pathophysiology* 2012. doi:10.1016/j.pathophys.2011.07.003.
- [305] Mohammadi MT. Overproduction of nitric oxide intensifies brain infarction and cerebrovascular damage through reduction of claudin-5 and ZO-1 expression in striatum of ischemic brain. *Pathol Res Pract* 2016. doi:10.1016/j.prp.2015.12.009.

- [306] Zheng L, Ding J, Wang J, Zhou C, Zhang W. Effects and Mechanism of Action of Inducible Nitric Oxide Synthase on Apoptosis in a Rat Model of Cerebral Ischemia-Reperfusion Injury. *Anat Rec* 2016. doi:10.1002/ar.23295.
- [307] Rudnitskaya A, Török B, Török M. Molecular docking of enzyme inhibitors: A COMPUTATIONAL TOOL FOR STRUCTURE-BASED DRUG DESIGN. *Biochem Mol Biol Educ* 2010. doi:10.1002/bmb.20392.
- [308] Varma AK, Patil R, Das S, Stanley A, Yadav L, Sudhakar A. Optimized hydrophobic interactions and hydrogen bonding at the target-ligand interface leads the pathways of Drug-Designing. *PLoS One* 2010. doi:10.1371/journal.pone.0012029.
- [309] Li MH, Inoue K, Si HF, Xiong ZG. Calcium-permeable ion channels involved in glutamate receptor-independent ischemic brain injury. *Acta Pharmacol. Sin.*, 2011. doi:10.1038/aps.2011.47.