List of Figures

| Figure No. | Figure Caption | Page |
|-------------|---|------|
| | | No. |
| Figure 1.1 | Nitrogen containing some main class of organic | 1 |
| | compounds | |
| Figure 1.2 | Some biological active drugs containing amine functional | 3 |
| | group. | |
| Figure 1.3 | Some biological active drugs containing imine functional group. | 5 |
| Figure 1.4 | Structure (A & B) and classifications (C & D) of oximes | 7 |
| Figure 1.5 | Examples of some drugs containing oxime groups | 8 |
| Figure 1.6 | Examples of some drugs containing amide groups | 10 |
| Figure 1.7 | Few biologically active compounds containing pyrrole | 12 |
| Eigung 1 0 | moiety Penrocentative compounds containing puriding | 14 |
| Figure 1.8 | Representative compounds containing pyridine substructure. | 14 |
| Figure 1.9 | Representatives indoles | 17 |
| Figure 1.10 | Few biologically active compounds containing | 19 |
| | benzothiazole moiety | |
| Figure 1.11 | Few biologically active compounds containing | 21 |
| | benzimidazole moiety | |
| Figure 1.12 | Few biologically active compounds containing | 23 |
| F: 0.1 | benzoxazole moiety | 47 |
| Figure 2.1 | FT-IR of s-Fe ₃ O ₄ | 47 |
| Figure 2.2 | XRD pattern of s-Fe ₃ O ₄ | 48 |
| Figure 2.3 | SEM image of s-Fe ₃ O ₄ | 48 |
| Figure 2.4 | EDAX of s-Fe ₃ O ₄ . | 48 |
| Figure 2.5 | TEM image s-Fe ₃ O ₄ | 49 |
| Figure 2.6 | MPMS analysis; Magnetic moment versus magnetic field | 49 |
| | graph of Fe ₃ O ₄ and s- Fe ₃ O ₄ | |
| Figure 2.7 | Recyclability of catalyst upto 6 runs | 62 |
| Figure 2.8 | XRD pattern of reused s-Fe ₃ O ₄ | 62 |
| Figure 2.9 | FT-IR of reused s-Fe ₃ O ₄ | 63 |
| Figure 2.10 | Plausible mechanism for s-Fe ₃ O ₄ catalyzed synthesis of | 64 |
| T' 0.11 | imidazopyrimidine | 72 |
| Figure 2.11 | ¹ H NMR & ¹³ C NMR of 2-Amino-4-phenyl-1,4 dihydrobenzo[4,5]imidazo[1,2-a]pyrimidine-3- | 73 |

| | carbonitrile (4a) | |
|-------------|---|-----|
| Figure 2.12 | ¹ H NMR & ¹³ C NMR of 2-Amino-4-phenyl-1,4 dihydrobenzo[4,5]imidazo [1,2-a]pyrimidine-3-carbonitrile (4m) | 74 |
| Figure 3.1 | Pharmacologically active benzimidazole and benzothiazole compounds | 84 |
| Figure 3.2 | Proposed mechanism for the formation of 1,2-disubstituted benzimidazoles and 2-substituted benzimidazoles | 102 |
| Figure 3.3 | ¹ H & ¹³ C NMR of 1-benzyl-2-phenyl-1H-benzo[d]imidazole (3a) | 112 |
| Figure 3.4 | ¹ H & ¹³ C NMR of 1-benzyl-2-phenyl-1H-benzo[d]imidazole (4a) | 113 |
| Figure 4.1 | Medicinally important molecules containing the pyranopyrazoles group | 128 |
| Figure 4.2 | Possible reaction mechanism | 141 |
| Figure 4.3 | ¹ H & ¹³ C NMR of 6-amino-3-methyl-4-phenyl-1,4-dihydropyrano[2,3-c]pyrazole-5-carbonitrile (4a) | 147 |
| Figure 5.1 | Some example of bioactive substituted amides. | 158 |
| Figure 5.2 | A plausible reaction pathway for the conversion of aldehyde derivatives to the amide | 170 |
| Figure 5.3 | ¹ H NMR and ¹³ C NMR of benzamide (3a) | 176 |
| Figure 5.4 | ¹ H NMR and ¹³ C NMR of <i>N</i> -Phenylacetamide (3k) | 177 |
| Figure 5.5 | Hydroxyphenyl)acetamide(3m) | 178 |