



## **REFERENCES**



---

## References:

- Agarwal, S., Tyagi, I., Gupta, V.K., Ghasemi, N., Shahivand, M., Ghasemi, M., 2016. Kinetics, equilibrium studies and thermodynamics of methylene blue adsorption on *Ephedra strobilacea* saw dust and modified using phosphoric acid and zinc chloride. *Journal of Molecular Liquids*, 218, 208-218.
- Ajmani, A., Shahnaz, T., Subbiah, S., Narayanasamy, S., 2019. Hexavalent chromium adsorption on virgin, biochar, and chemically modified carbons prepared from *Phanera vahlii* fruit biomass: equilibrium, kinetics, and thermodynamics approach. *Environmental Science and Pollution Research*, 26, 32137–32150.
- Ali, I., Asim, M., Khan, T.A., 2012. Low cost adsorbents for the removal of organic pollutants from wastewater. *J. Environ. Manag.*, 113, 170–183.
- Aljeboree, A.M., Alkaim, A. F, Al-Dujaili, A.H., 2015. Adsorption isotherm, kinetic modeling and thermodynamics of crystal violet dye on coconut husk-based activated carbon. *Desalin. Water Treat.*, 53, 3656–3667.
- Al-Othman, Z.A., Ali, R., Naushad, M., 2012. Hexavalent chromium removal from aqueous medium by activated carbon prepared from peanut shell: adsorption kinetics, equilibrium and thermodynamic studies. *Chemical Engineering Journal*, 184, 238-247.
- Al-Othman, Z.A., Naushad, M., Ali, R., 2013. Kinetic, equilibrium isotherm and thermodynamic studies of Cr (VI) adsorption onto low-cost adsorbent developed from peanut shell activated with phosphoric acid. *Environmental Science and Pollution Research*, 20(5), 3351-3365.
- An, B., Liang, Q., Zhao, D., 2011. Removal of arsenic (V) from spent ion exchange brine using a new class of starch-bridged magnetite nanoparticles, *Water Res.*, 45, 1961–1972.
- Anderson, L.D., Kent, D.B., Davis, J.A., 1994. Batch experiments characterizing the reduction of chromium (VI) using sub-toxic material from a mildly reducing sand and gravel aquifer. *Environmental Science and Technology*, 28, 178–185.

- Anupam, K., Dutta, S., Bhattacharjee, C., Datta, S., 2011. Adsorptive removal of chromium (VI) from aqueous solution over powdered activated carbon: Optimization through response surface methodology. *Chemical Engineering Journal*, 173, 135–143.
- Argun, M.E., Dursun, S., Karatas, M., Gürü, M., 2008. Activation of pine cone using Fenton oxidation for Cd (II) and Pb (II) removal. *Bioresour. Technol.*, 99, 8691–8698.
- Baral, S. S., Das, S. N., Rath, P., 2006. Hexavalent chromium removal from aqueous solution by adsorption on treated saw dust. *Biochemical Engineering Journal*, 31, 216–222.
- Basu, S., Ghosh, G., Saha, S., 2018. Adsorption characteristics of phosphoric acid induced activation of bio-carbon: Equilibrium, kinetics, thermodynamics and batch adsorber design. *Process Safety and Environmental Protection*, 117, 125-142.
- Bayazit, S.S., Kerkez, O., 2014. Hexavalent chromium adsorption on superparamagnetic multi-wall carbon nanotubes and activated carbon composites. *Chemical Engineering Research and Design*, 92, 2725-2733.
- Baysal, M., Bilge, K., Yılmaz, B., Papila, M., Yürüm, Y. 2018. Preparation of high surface area activated carbon from waste-biomass of sunflower piths: Kinetics and equilibrium studies on the dye removal. *Journal of Environmental Chemical Engineering*, 6(2), 1702-1713.
- Baytar, O., Şahin, O., Saka, C., 2018. Sequential application of microwave and conventional heating methods for preparation of activated carbon from biomass and its methylene blue adsorption. *Applied Thermal Engineering*, 138, 542-551.
- Beakou, B.H., E.I. Hassani, K., Houssaini, M.A., Belbahloul, M., Oukani, E., Anouar, A., 2017. Novel activated carbon from *Manihot esculenta* Crantz for removal of Methylene Blue. *Sustainable Environment Research*, 27(5), 215-222.
- Bilal, M., Shah, J.A., Ashfaq, T., Gardazi, S.M.H., Tahir, A.A., Pervez, A., Haroon, H., Mahmood, Q., 2013. Waste biomass adsorbents for copper removal from industrial wastewater-A review. *J. Hazard. Mater.*, 263, 322–333.
- Bouguettoucha, A., Reffas, A., Chebli, D., Mekhalif, T., Amrane, A., 2016. Novel activated carbon prepared from an agricultural waste, *Stipa tenacissima*, based on  $ZnCl_2$

- activation-characterization and application to the removal of methylene blue. *Desalin. Water Treat.*, 57 (50), 24056-24069.
- Bulut, Y., Aydin, H., 2006. A kinetics and thermodynamics study of methylene blue adsorption on wheat shells. *Desalination*, 194, 259–267.
- Chaudhuri, M., Azizan, N.K.B., 2012. Adsorptive removal of chromium (VI) from aqueous solution by an agricultural waste-based activated carbon. *Water, Air, & Soil Pollution*, 223(4), 1765-1771.
- Chia, C.H., Gong, B., Joseph, S.D., Marjo, C.E., Munroe, P., Rich, A.M., 2012. Imaging of mineral-enriched biochar by FTIR, Raman and SEM-EDX. *Vibrational Spectroscopy*, 62, 248–257.
- Chowdhury, S., Mishra, R., Saha, P., Kushwaha, P., 2011. Adsorption thermodynamics, kinetics and isosteric heat of adsorption of malachite green onto chemically modified rice husk. *Desalination*, 265, 159–168.
- Clemens, S., Ma, F., 2016. Toxic heavy metal and metalloid accumulation in crop plants and foods. *Annu. Rev. Plant Biol.*, 67 (1), 489-512.
- Coman, V., Robotin, B., Ilea, P., 2013. Nickel recovery/removal from industrial wastes: a review. *Resour. Conserv. Recycl.*, 73, 229–238.
- Crini, G., 2006. Non-conventional low-cost adsorbents for dye removal: a review. *Bioresour. Technol.*, 97, 1061-1085.
- Cronje, K.J., Chetty, K., Carsky, M., Sahu, J.N, Meikap, B.C. 2011. Optimization of chromium (VI) sorption potential using developed activated carbon from sugarcane bagasse with chemical activation by zinc chloride. *Desalination*, 275, 276–284.
- Cruz, G., Pirila, M., Huuhtanen, M., Carrion, L., Alvarenga, E., Keiski, R.L. 2012. Production of activated carbon from Cocoa (*Theobroma cacao*) pod husk. *Civ Environ Eng.*, 2(2), 1–6.
- Danish, M., Ahmad, T., Majeed, S., Ahmad, M., Ziyang, L., Pin, Z., Iqbal, S.S. 2018. Use of banana trunk waste as activated carbon in scavenging methylene blue dye: kinetic, thermodynamic, and isotherm studies. *Bioresource Technology Reports*, 3, 127-137.

- Dardouri, S., Sghaier, J., 2017. A comparative study of adsorption and regeneration with different agricultural wastes as adsorbents for the removal of methylene blue from aqueous solution. *Chin. J. Chem. Eng.*, 25, 1282-1287.
- Das, D., Samal, D.P., Meikap, B.C., 2015. Preparation of activated carbon from green coconut shell and its characterization. *J. Chem. Eng. Process. Technol.*, 6(5), 1–7.
- Dawood, S., Sen, T.K., Phan, C., 2014. Synthesis and characterisation of novel-activated carbon from waste biomass pine cone and its application in the removal of Congo red dye from aqueous solution by adsorption. *Water Air Soil Pollut.*, 225(1), 1–16.
- Demirbas, A., 2008. Heavy metal adsorption onto agro-based waste materials: a review. *J. Hazard. Mater.*, 157, 220–229.
- Dhir, B., Sharmila, P., Saradhi, P.P., 2009. Potential of aquatic macrophytes for removing contaminants from the environment, *Critical Review. Environ. Sci. Technol.*, 39,754–781.
- Doke, K.M., Khan, E.M. 2017. Equilibrium, kinetic and diffusion mechanism of Cr (VI) adsorption onto activated carbon derived from wood apple shell. *Arabian Journal of Chemistry*, 10, S252-S260.
- Droussi, Z., D’orazio, V., Provenzano, M. R., Hafidi, M., Ouatmane, A., 2009. Study of the biodegradation and transformation of olive-mill residues during composting using FTIR spectroscopy and differential scanning calorimetry. *J. Hazard. Mater.*, 164 (2–3), 1281–1285.
- Droussi, Z., Dorazio, V., Provenzano, M.R., Hafidi, M., Ouatmane, A., 2009. Study of the biodegradation and transformation of olive-mill residues during composting using FTIR spectroscopy and differential scanning calorimetry. *J. of Hazard. Mater.*, 164(2–3), 1281–1285.
- El-Shafey, E., Ali, S.N., Al-Busafi, S., Al-Lawati, H.A., 2016. Preparation and characterization of surface functionalized activated carbons from date palm leaflets and application for methylene blue removal. *Journal of Environmental Chemical Engineering*, 4(3), 2713-2724.

- Enniya, I., Rghioui, L., Jourani, A., 2018. Adsorption of hexavalent chromium in aqueous solution on activated carbon prepared from apple peels. *Sustainable Chemistry and Pharmacy*, 7, 9-16.
- Ewecharoen, A. Thiravetyan, P., Wendel, E., Bertagnolli, H., 2009. Nickel adsorption by sodium polyacrylate-grafted activated carbon. *J. Hazard. Mater.*, 171, 335–339.
- Fan, L., Wan, W., Wang, X., Cai, J., Chen, F., Chen, W., Ji, L., Luo, H., Cheng, L. 2019. Adsorption Removal of Cr (VI) with activated carbon prepared by co-pyrolysis of rice straw and sewage sludge with ZnCl<sub>2</sub> activation. *Water, Air, & Soil Pollution*, 230(10), 233.
- Fernandez, M.E., Ledesma, B., Román, S., Bonelli, P.R., Cukierman, A.L. 2015. Development and characterization of activated hydrochars from orange peels as potential adsorbents for emerging organic contaminants. *Bioresour. Technol.*, 183, 221–228.
- Freundlich, H.M.F. 1906. Uber die adsorption in losungen, *J. Chem. Phys*, 57 ,385–470.
- Fu, F., Wang, Q., 2011. Removal of heavy metal ions from wastewaters: A review. *Journal of Environmental Management*, 92, 407-418.
- Girgis, B.S., Yunis, S.S., Soliman, A.M., 2002. Characteristics of activated carbon from peanut hulls in relation to conditions of preparation. *Mater. Lett.*, 57, 164–72.
- Gorzin, F., Ghoreyshi, A.A., 2013. Synthesis of a new low-cost activated carbon from activated sludge for the removal of Cr (VI) from aqueous solution: Equilibrium, kinetics, thermodynamics and desorption studies. *Korean Journal of Chemical Engineering*, 30(8), 1594-1602.
- Goswami, M., Borah, L., Mahanta, D., Phukan, P., 2014. Equilibrium modeling, kinetic and thermodynamic studies on the adsorption of Cr (VI) using activated carbon derived from matured tea leaves. *Journal of Porous Materials*, 21(6), 1025-1034.
- Gottipati, R., Mishra, S., 2016. Preparation of microporous activated carbon from Aegle Marmelos fruit shell and its application in removal of chromium (VI) from aqueous phase. *Journal of Industrial and Engineering Chemistry*, 36, 355-363.

- Gupta V.K., Agarwal S., Saleh T., 2011 Synthesis and characterization of alumina coated carbon nanotubes and their application for lead removal. *J. Hazard. Mater.*, 185, 17–23.
- Gupta, S., Babu, B.V., 2009. Utilization of waste product (tamarind seeds) for the removal of Cr(VI) from aqueous solutions: equilibrium, kinetics, and regeneration studies. *Journal of Environmental Management*, 90, 3013–3022.
- Hameed, B.H., Ahmad, A.L., Latiff, K.N.A., 2007. Adsorption of basic dye (methylene blue) onto activated carbon prepared from rattan sawdust. *Dyes Pigm.*, 75, 143–149.
- Han, X., Wang, H., Zhang, L., 2018. Efficient removal of methylene blue using nanoporous carbon from the waste biomass. *Water, Air, Soil Pollut.*, 229, 26.
- Han, W., Cheng, F. F, Z., Tang, B., Wu, S., 2016. Studies on the optimum conditions using acid- washed zero-valent iron/aluminum mixtures in permeable reactive barriers for the removal of different heavy metal ions from wastewater. *J. Hazard. Mater.*, 302, 437–446.
- Hashem, A., Akasha, R.A., Ghith, A., Hussein, D.A., 2007. Adsorbent based on agricultural wastes for heavy metal and dye removal: a review. *Energy Educ. Sci. Technol.*, 19, 69–86.
- Hashim, M.A., Mukhopadhyay, S., Sahu, J.N., Sengupta, B., 2011. Remediation technologies for heavy metal contaminated groundwater, *J. Environ. Manage.*, 92, 2355–2388.
- Holkar , C. R., Jadhav, A. J., Pinjari, D. V., Mahamuni, N. M., Pandit, A. B., 2016. A critical review on textile wastewater treatments: possible approaches, *J. Environ. Manage.*, 182, 351–366.
- Hosseini, S.S., Bringas, E., Tan, N.R., Ortiz, I., Ghahramani, M., Shahmirzadi, M.A. A., 2016. Recent progress in development of high performance polymeric membranes and materials for metal plating wastewater treatment: a review. *J. Water Process Eng.*, 9, 78–110.

- Hsua, N.H., Wang, S. L., Liao, Y.H., Huang, S.T., Tzou, Y.M., Huang, Y.M., 2009. Removal of hexavalent chromium from acidic aqueous solutions using rice straw-derived carbon. *J. Hazard. Mater.*, 171, 1066–1070.
- Imran, M., Crowley, D. E., Khalid, A., Hussain, S., Mumtaz, M. W., Arshad, M., 2015. Microbial biotechnology for decolorization of textile wastewaters. *Reviews in Environmental Science and Biotechnology*, 14(1), 73-92.
- Islam, M.A., Tan I.A.W., Benhouria, A., Asif, M., Hameed, B.H., 2015. Mesoporous and adsorptive properties of palm date seed activated carbon prepared via sequential hydrothermal carbonization and sodium hydroxide activation. *Chem. Eng. J.*, 270, 187-195.
- Islam, M.A., Sabar, S., Benhouria, A., 2017. Nanoporous activated carbon prepared from karanj (*Pongamia pinnata*) fruit hulls for methylene blue adsorption. *J. Taiwan Inst. Chem. Eng.*, 74, 96–104.
- Islam, M.A., Sabar, S., Benhouria, A., Khanday, W., Asif, M., Hameed, B. 2017. Nanoporous activated carbon prepared from karanj (*Pongamia pinnata*) fruit hulls for methylene blue adsorption. *Journal of the Taiwan Institute of Chemical Engineers*, 74, 96-104.
- Jain, M., Garg, V.K., Kadirvelu, K., 2010. Adsorption of hexavalent chromium from aqueous medium onto carbonaceous adsorbents prepared from waste biomass. *J. Environ. Manag.*, 91, 949–957.
- Jin, X.J., Yu, Z.M., Wu, Y., 2010. Preparation of activated carbon from lignin obtained by straw pulping by KOH and  $K_2CO_3$  chemical activation. *Cellul Chem Technol*, 46(1–2), 79–85.
- Jung, C., Heo, J., Han, J., Her, N., Lee, S. J., Oh, J., Ryu, J., Yoon, Y., 2013. Hexavalent chromium removal by various adsorbents: Powdered activated carbon, chitosan, and single/multi-walled carbon nanotubes. *Separation and Purification Technology*, 106, 63–71.
- Kanjanarong, J., Giri, B.S., Jaisi, D.P., Oliveira, F.R., Boonsawang, P., Khanal, S.K., 2017. Removal of hydrogen sulfide generated during anaerobic treatment of sulfate-



laden wastewater using biochar: evaluation of efficiency and mechanisms. *Bioresource Technology*, 234, 115–121.

Karacetin, G., Sivrikaya, S., Imamoglu, M. 2014. Adsorption of Methylene Blue from aqueous solutions by activated carbon prepared from hazelnut husk using zinc chloride. *J Anal. Appl. Pyrolysis.*, 110, 270–276.

Katheresan, V., Kansedo, J., Lau, S.Y., 2018. Efficiency of various recent wastewater dye removal methods: A review. *Journal of Environmental Chemical Engineering*, 6, 4676–4697.

Khan, M.I., Min, T.K., Azizli, K., Sufian, S., Man, Z., Ullah, H., 2015. Effective removal of methylene blue from water using phosphoric acid based geopolymers. *RSC Adv.*, 5, 61410-61420.

Khezami, L., Ould-Dris, A., Capart, R., 2007. Activated carbon from thermo-compressed wood and other ligno-cellulosic precursors. *Bioresources.*, 2(2), 193–209.

Koedrith, P., Kim, H.L., Weon, J., Seo, Y.R., 2013. Toxicogenomic approaches for understanding molecular mechanisms of heavy metal mutagenicity and carcinogenicity, *Int. J. Hyg. Environ. Health*. 216, 587–598.

Kula, I., Ugurlu, M., Karaoglu, H., 2008. Adsorption of Cd (II) ions from aqueous solutions using activated carbon prepared from olive stone by ZnCl<sub>2</sub> activation, *Bioresource Technology*, 99, 492–501.

Kumar, A., Jena, H.M., 2017. Adsorption of Cr (VI) from aqueous phase by high surface area activated carbon prepared by chemical activation with ZnCl<sub>2</sub>. *Process Safety and Environmental Protection*, 109, 63-71.

Kumar, M., Tamilarasan, R., 2013. Modeling studies for the removal of methylene blue from aqueous solution using *Acacia fumosa* seed shell activated carbon. *Journal of Environmental Chemical Engineering*, 1(4), 1108-1116.

Kumar, M., Tamilarasan, R., 2017. Kinetics, equilibrium data and modeling studies for the sorption of chromium by *Prosopis juliflora* bark carbon. *Arabian Journal of Chemistry*, 10, S1567-S1577.

- Kumar, P., Ramalingam, S., Sathishkumar, K., 2011. Removal of methylene blue dye from aqueous solution by activated carbon prepared from cashew nut shell as a new low-cost adsorbent. *Korean J. Chem. Eng.*, 28,149–155.
- Lam, S.S., Liew, R.K., Wong, Y.M., Yek, P.N.Y., Ma, N.L., Lee, C.L., Chase, H.A., 2017. Microwave-assisted pyrolysis with chemical activation, an innovative method to convert orange peel into activated carbon with improved properties as dye adsorbent. *J. Clean. Prod.*, 162, 1376–1387.
- Langmuir, 1918. The Adsorption of gases on plane surfaces of glass, mica and platinum. *J. Am. Chem. Soc.*, 40, 1361–1403.
- Lei Y., Su H., Tian R., 2016. Morphology evolution, formation mechanism and adsorption properties of hydro chars prepared by hydrothermal carbonization of corn stalk. *RSC Adv.*, 6, 107829–107835.
- Li, Z., Wang, G., Zhai, K., He, C., Li, Q., Guo, P., 2018. Methylene blue adsorption from aqueous solution by loofah sponge-based porous carbons. *Colloids Surf. A.*, 538, 28-35.
- Li, J., Lin, Q., Zhang, X., Yan, Y., 2009. Kinetic parameters and mechanisms of the batch biosorption of Cr(VI) and Cr(III) onto *Leersia hexandra* Swartz biomass. *J. Colloid Interface Sci.*, 333, 71–77.
- Liu, H., Liang, S., Gao, J., Ngo, H.H., Guo, W., Guo, Z., Wang, J., Li, Y., 2014. Enhancement of Cr (VI) removal by modifying activated carbon developed from *Zizania caduciflora* with tartaric acid during phosphoric acid activation. *Chemical Engineering Journal*, 246, 168-174.
- Ma, H., Li, J.B., Liu, W.W., Miao, M., Cheng, B.J., Zhu, S.W., 2015. Novel synthesis of a versatile magnetic adsorbent derived from corncob for dye removal. *Bioresour. Technol.*, 190, 13–20.
- Mahamad, M.N., Zaini, M.A.A., Zakaria, Z.A., 2015. Preparation and characterization of activated carbon from pineapple waste biomass for dye removal. *Int. Biodeterior. Biodegrad.*, 102, 274–280.

- Mahamad, M.N., Zaini, M.A.A., Zakaria, Z.A., 2015. Preparation and characterization of activated carbon from pineapple waste biomass for dye removal. *International Biodeterioration & Biodegradation*, 102, 274-280.
- Mahmoud, D.K., Salleh, M.A.M., Karim, Z.A.W.A, Idris, A., Abidin, Z.Z., 2012. Batch adsorption of basic dye using acid treated kenaf fibre char: equilibrium, kinetic and thermodynamic studies. *Chem. Eng. J.*, 181–182, 449–457.
- Malkoc, E., Nuhoglu, Y., Dundar, M., 2006. Adsorption of chromium (VI) on pomace-an olive oil industry waste: batch and column studies. *J. Hazard. Mater.*, 138, 142–151.
- Mamanía, A., Ramírezb, N., Deianaa, C., Giméneza, M., Sardella, F., 2019. Highly microporous sorbents from lignocellulosic biomass: Different activation routes and their application to dyes adsorption. *J. of Envir. Chem. Eng*, 7, 1031-1048.
- Maneechakr, P., Karnjanakom, S., 2017. Adsorption behaviour of Fe (II) and Cr (VI) on activated carbon: Surface chemistry, isotherm, kinetic and thermodynamic studies. *The Journal of Chemical Thermodynamics*, 106, 104-112.
- Miyah, Y., Lahrichi, A., Idrissi, M., 2018. Adsorption of methylene blue dye from aqueous solutions onto walnut shells powder: equilibrium and kinetic studies. *Surf. Interfaces*, 11, 74–81.
- Mohanty, K., Jha, M., Meikap, B.C., Biswas, M.N., 2005. Removal of chromium (VI) from dilute aqueous solutions by activated carbon developed from Terminalia arjuna nuts activated with zinc chloride. *Chemical Engineering Science*, 60, 3049–3059.
- Mohanty, K., Naidu, J.T., Meikap, B.C., Biswas, M.N., 2006. Removal of crystal violet from wastewater by activated carbons prepared from rice husk. *Ind. Eng. Chem. Res.*, 45, 5165–5171.
- Naeem, S., Baheti, V., Wiener, J., Marek, J., 2017. Removal of methylene blue from aqueous media using activated carbon web. *J. Textil. Inst.*, 108, 803-811.
- Nakajima, A., Baba, Y., 2004. Mechanism of hexavalent chromium adsorption by persimmon tannin gel. *Water Research.*, 38, 2859–2864.
- Nandi, B., Goswami, A., Purkait, M., 2009. Removal of cationic dyes from aqueous solutions by kaolin: kinetic and equilibrium studies. *Appl. Clay Sci.*, 42(3–4), 583–590.

- Nasrullah, A., Saad, B., Bhat, A., Khan, A.S., Danish, M., Isa, M.H., Naeem, A., 2019. Mangosteen peel waste as a sustainable precursor for high surface area mesoporous activated carbon: Characterization and application for methylene blue removal. *Journal of Cleaner Production*, 211, 1190-1200.
- Nemr, A.E., 2009. Potential of pomegranate husk carbon for Cr (VI) removal from wastewater: kinetic and isotherm studies. *Journal of Hazardous Material*, 161, 132–141.
- Niazi, L., Lashanizadegan, A., Sharififard, H., 2018. Chestnut oak shells activated carbon: Preparation, characterization and application for Cr (VI) removal from dilute aqueous solutions. *Journal of Cleaner Production*, 185, 554-561.
- Norouzi, S., Heidari, M., Alipour, V., Rahmanian, O., Fazlzadeh, M., Mohammadi-Moghadam, F., Nourmoradi, H., Goudarzi, B., Dindarloo, K., 2018. Preparation, characterization and Cr (VI) adsorption evaluation of NaOH-activated carbon produced from Date Press Cake; an agro-industrial waste. *Bioresource Technology*, 258, 48-56.
- Novais, R.M., Caetano, A.P., Seabra, M.P., Labrincha, J.A., Pullar, R.C. 2018. Extremely fast and efficient methylene blue adsorption using eco-friendly cork and paper waste-based activated carbon adsorbents. *Journal of cleaner production*, 197, 1137-1147.
- Ofomaja, A.E., Naidoo, E.B., 2011. Biosorption of copper from aqueous solution by chemically activated pine cone: a kinetic study. *Chem. Eng. J.*, 175, 260–70.
- Ogugbue, C.J., Sawidis, T., 2011. Bioremediation and detoxification of synthetic wastewater containing Triaryl methane dyes by *aeromonas hydrophila* isolated from industrial effluent. *Biotechnol. Res. Int.*, 2011, 967925.
- Onal, Y., Basar, A.C., Ozdemir, S. C., Erdogan, S., 2007. Textural development of sugar beet bagasse activated with  $ZnCl_2$ . *J. Hazard. Mat.*, 142, 138–143.
- Onal, Y., Basar, A.C., Ozdemir, S.C., Erdogan, S., 2007. Textural development of sugar beet bagasse activated with  $ZnCl_2$ . *J. Hazard. Mater.*, 142, 138–143.
- Onal, Y., Basar, A.C., Ozdemir, S.C., Erdogan, S., 2007. Textural development of sugar beet bagasse activated with  $ZnCl_2$ . *Journal of Hazardous Material*, 142, 138–143.

- Owlad, M., Aroua, M. K., Daud W. M. A. W., 2010. Hexavalent chromium adsorption on impregnated palm shell activated carbon with polyethyleneimine. *Bioresource Technology*, 101, 5098–5103.
- Pari, G., Darmawan, S., Prihandoko, B., 2014. Porous carbon spheres from hydrothermal carbonization and KOH activation on cassava and tapioca flour raw material. *Procedia. Environ. Procedia. Environ. Sci.*, 20, 342–351.
- Pathania, D., Sharma, S., Singh, P., 2017. Removal of methylene blue by adsorption onto activated carbon developed from *Ficus carica* bast. *Arabian. J. Chem.*, 10, S1445-S1451.
- Pedroza, M.M., Sousa, J.F., Vieira, G.E.G., Bezerra, M.B.D., 2014. Characterization of the products from the pyrolysis of sewage sludge in 1kg/h rotating cylinder reactor. *Journal of Analytical and Applied Pyrolysis*, 105, 108–115.
- Pezoti Jr, O., Cazetta, A.L., Souza, I.P., Bedin, K.C., Martins, A.C., Silva, T.L., Almeida, V.C., 2014. Adsorption studies of methylene blue onto ZnCl<sub>2</sub>-activated carbon produced from buriti shells (*Mauritia flexuosa* L.). *Journal of Industrial and Engineering Chemistry*, 20(6), 4401-4407.
- Ponnusami, V., Krithika, V., Madhuran, R., Srivastava, S.N., 2007. Biosorption of reactive dye using acid-treated rice husk: factorial design analysis. *J. Hazard. Mater.*, 142, 397–403.
- Qu, X., Alvarez, P.J.J., Li, Q., 2013. Applications of nanotechnology in water and wastewater treatment. *Water Res.*, 47 (12) (2013), 3931–3946.
- Rangabhashiyam, S., Selvaraju, N., 2015. Adsorptive remediation of hexavalent chromium from synthetic wastewater by a natural and ZnCl<sub>2</sub> activated *Sterculia guttata* shell. *Journal of Molecular Liquids*, 207, 39-49.
- Rangabhashiyam, S., Selvaraju, N., 2015 (b). Efficacy of unmodified and chemically modified *Swietenia mahagoni* shells for the removal of hexavalent chromium from simulated wastewater. *Journal of Molecular Liquids*, 209, 487–497.
- Rangabhashiyam, S., Selvaraju, N., 2015 (c). Evaluation of the biosorption potential of a novel *Caryota urens* inflorescence waste biomass for the removal of hexavalent

chromium from aqueous solutions. *Journal of the Taiwan Institute of Chemical Engineers*, 47, 59–70.

Rangabhashiyam, S., Selvaraju, N., 2015(a). Adsorptive remediation of hexavalent chromium from synthetic wastewater by a natural and ZnCl<sub>2</sub> activated *Sterculia guttata* shell. *J. of Mole. Liq.*, 207, 39–49.

Raval, N.P., Shah, P.U., Shah, N.K. 2016. Adsorptive removal of nickel (II) ions from aqueous environment: a review, *J. Environ. Manage.*, 179, 1-20.

Ravi, K., Deebika, B., Balu, K., 2005. Decolourization of aqueous dye solutions by a novel adsorbent: application of statistical designs and surface plots for the optimization and regression analysis, *J. Hazard. Mater.*, B122, 75–83.

Renuka, N., Sood, A., Ratha, S.K, Prasanna, R., Ahluwalia, A.S., 2013. Evaluation of microalgal consortia for treatment of primary treated sewage effluent and biomass production. *J. Appl. Phycol.* 25, 1529–1537.

Rodríguez-Couto, S., Osma, J.F., Toca-Herrera, J.L., 2009. Removal of synthetic dyes by an eco-friendly strategy. *Engineering in Life Sciences*, 9(2), 116-123

Sangon, S., Hunt, A.J., Attard, T.M., Mengchang, P., Ngernyen, Y., Supanchaiyamat, N., 2018. Valorisation of waste rice straw for the production of highly effective carbon based adsorbents for dyes removal. *J Clean. Prod.*, 172, 1128–1139.

Sharma, G., Naushad, M., Kumar, A., Rana, S., Sharma, S., Bhatnagar, A.J., Stadler, F., Ghfar, A.A., Khan, M.R., 2017. Efficient removal of coomassie brilliant blueR-250 dye using starch/poly(alginate-chitosan) nanohydrogel. *Process Saf. Environ. Prot.*, 109, 301–310.

Singh, K., Giri, B.S., Sahi, A., Geed, S.R., Kureel, M.K., Singh, S., Dubey, S.K., Rai, B.N., Kumar, S., Upadhyay, S.N., Singh, R.S., 2017. Biofiltration of xylene using wood charcoal as the biofilter media under transient and high loading conditions. *Bioresource Technology*, 242, 351–358.

Spagnoli, A.A., Giannakoudakis, D.A., Bashkova, S., 2017. Adsorption of methylene blue on cashew nut shell based carbons activated with zinc chloride: The role of surface and structural parameters. *Journal of Molecular Liquids*, 229, 465-471.

- Srivastava, N.K., Majumder, C.B., 2008. Novel biofiltration methods for the treatment of heavy metals from industrial wastewater. *J. Hazard. Mater.*, 151, 1–8.
- Suksabye, P., Thiravetyan, P., 2012. Cr (VI) adsorption from electroplating plating wastewater by chemically modified coir pith. *Journal of Environment Management*, 102, 1–8.
- Talha, M.A., Goswami, M., Giri, B.S., Sharma, A., Rai, B.N., Singh, R.S., 2018. Bioremediation of Congo red dye in immobilized batch and continuous packed bed bioreactor by *Brevibacillus parabrevis* using coconut shell bio-char. *Bioresource Technology*, 252, 37–43.
- Tan, I.A.W., Hameed, B.H., Ahmad, A.L., 2007. Equilibrium and kinetic studies on basic dye adsorption by oil palm fibre activated carbon. *Chem. Eng. J.*, 127, 111–119.
- Tan, C., Zeyu, Z., Sai, X., Hongtao, W., Wenjing, L., 2015. Adsorption behaviour comparison of trivalent and hexavalent chromium on biochar derived from municipal sludge. *Bioresource Technology*, 190, 388–394.
- Tan, I.A.W., Ahmad, A.L., Hameed, B.H., 2008. Adsorption of basic dye using activated carbon prepared from oil palm shell: batch and fixed bed studies. *Desalination*, 225, 13–28.
- Tan, I.A.W., Hameed, B.H., Ahmad, A.L., 2007. Equilibrium and kinetic studies on basic dye adsorption by oil palm fibre activated carbon. *Chem. Eng. J.*, 127, 111–119. *Environ. Sci. Pollut. Res.* 23 (20), 20087–20099.
- Tanong, K., Tran, L.H., Mercier, G., Blais, J.F., 2017. Recovery of Zn (II), Mn (II) Cd (II) and Ni (II) from the unsorted spent batteries using solvent extraction, electrodeposition and precipitation methods. *J. Clean. Prod.*, 148, 233–244.
- Tran, V.S., Ngo, H.H., Guo, W., Zhang, J., Liang, S., Ton-That, C., Zhang, X., 2015. Typical low cost biosorbents for adsorptive removal of specific organic pollutants from water. *Bioresour. Technol.*, 182, 353–363.
- Trivedi, N.S., Mandavgane, S.A., Kulkarni, B.D., 2016. Mustard plant ash: a source of micronutrient and an adsorbent for removal of 2, 4-dichlorophenoxyacetic acid. *Environmental Science and Pollution Research*, 23 (20), 20087–20099.

- Ugurlu, M., Karaoglu, H., 2008. Adsorption of Cd (II) ions from aqueous solutions using activated carbon prepared from olive stone by ZnCl<sub>2</sub> activation. *Bioresource Technology*, 99, 492–501.
- Uner, O. 2019. Hydrogen storage capacity and methylene blue adsorption performance of activated carbon produced from *Arundo donax*. *Materials Chemistry and Physics*, 237, 121858.
- Verma, A., Chakraborty, S., Basu, J.K., 2006. Adsorption study of hexavalent chromium using tamarind hull-based adsorbents. *Separation and Purification Technology*, 50, 336–341.
- Vunain, E., Mishra, A.K., Mamba, B.B., 2016. Dendrimers, mesoporous silicas and chitosan-based nanosorbents for the removal of heavy-metal ions: a review, *Int. J. Biol. Macromol.*, 86, 570–586.
- Wang, H., Xie, R., Zhang, J., Zhao, J. 2018. Preparation and characterization of distillers' grain based activated carbon as low cost methylene blue adsorbent: Mass transfer and equilibrium modeling. *Advanced Powder Technology*, 29(1), 27-35.
- Wang, S.Y., Tang, Y.K., Li, K., Mo, Y.Y., Li, H.F., Gu, Z.Q., 2014. Combined performance of biochar sorption and magnetic separation processes for treatment of chromium-contained electroplating waste water. *Bioresource Technology*, 174, 67–73.
- Williams, P.T., Reed, A.R., 2004. High grade activated carbon matting derived from the chemical activation and pyrolysis of natural fibre textile waste. *J. Anal. Appl. Pyrol.* 71, 971–86.
- Xu, Q., Qian, Q., Quek, A., Ai, N., Zeng, G., Wang, J., 2013. Hydrothermal carbonization of macro algae and the effects of experimental parameters on the properties of hydro chars. *ACS Sustain Chem Eng*, 1, 1092–1101.
- Yagub, M.T., Sen, T.K., Afroze, S., Ang, H.M., 2014. Dye and its removal from aqueous solution by adsorption: a review. *Adv. Colloid Interface Sci.*, 209, 172–184
- Yang, J., Yu, M., Chen, W., 2015. Adsorption of hexavalent chromium from aqueous solution by activated carbon prepared from longan seed: Kinetics, equilibrium and thermodynamics. *Journal of industrial and engineering chemistry*, 21, 414-422.



- Zare, E.N., Mansour Lakouraj, M., Ramezani, A., 2016. Efficient sorption of Pb (ii) from an aqueous solution using a poly (aniline-co-3-aminobenzoic acid)-based magnetic core-shell nanocomposite. *New J. Chem.* 40, 2521–2529.
- Zhang, Y., Song, X., Xu, Y., Shen, H., Kong, X., Xu, H., 2019. Utilization of wheat bran for producing activated carbon with high specific surface area via NaOH activation using industrial furnace. *Journal of cleaner production*, 210, 366-375.
- Zhang, B., Xiong, S., Xiao, B., Yu, D., Jia, X., 2011. Mechanism of wet sewage sludge pyrolysis in a tubular furnace. *International Journal of Hydrogen Energy*, 36(1), 355–363.
- Zhang, H., Tang, Y., Cai, D., Liu, X., Wang, X., Huang, Q., Yu, Z., 2010. Hexavalent chromium removal from aqueous solution by algal bloom residue derived activated carbon: equilibrium and kinetic studies. *J. Hazard. Mater.*, 181, 801–808.
- Zhang, J., Chen, H., Chen, Z., He, J., Shi, W., Liu, D., Chi, H., Fuyi, C., Wang, W., 2016. Microstructured macroporous adsorbent composed of polypyrrole modified natural corn cob-core sponge for Cr (VI) removal. *RSC Adv.* 64, 59292-59298.
- Zhou, Y., Zhou J.L.Y., Liu Y., 2019. Recent advances for dyes removal using novel adsorbents: A review. *Environmental Pollution*, 252, 352-365.
- Zou, Z., Tang, Y., Jiang, C., Zhang, J., 2015. Efficient adsorption of Cr (VI) on sunflower seed hull derived porous carbon. *Journal of Environmental Chemical Engineering*, 3(2), 898-905.