

List of Tables

<i>Table 2.1: Details of structural and fitting parameters as obtained from high-resolution x-ray diffraction. Compound, lattice constant, variable coordinate x in the 48f-site, R-factors (R_p, R_{wp} and R_e) and goodness of fit χ^2.....</i>	58
<i>Table 2.2: Details of structural and fitting parameters as obtained from high-resolution x-ray diffraction. Compound, lattice constant, variable coordinate x in the 48f-site, R-factors (R_p, R_{wp} and R_e) and goodness of fit χ^2.....</i>	59
<i>Table 3.1: Details of structural refinement as obtained from high-resolution x-ray diffraction. Compound, lattice constant, variable coordinate x in the 48f-site, R-factors (R_p, R_{wp} and R_e) and goodness of fit χ^2.....</i>	62
<i>Table 3.2: Value of activation energy E_a (meV) and characteristic relaxation time τ_0 (sec) for 90 K and 36 K relaxations deduced from- peak position T_m'' of temperature dependent ϵ'' and Lacroix-Bènè function fitted obtained parameters τ_0 (in the bracket.).....</i>	67
<i>Table 3.3: Obtained values of fitting parameters ϵ_s, ϵ_∞, β, and τ_0 referring to the dispersion data ϵ' of figure 6 for $\text{Ho}_2\text{Ti}_2\text{O}_7$ and $\text{Dy}_2\text{Ti}_2\text{O}_7$.....</i>	69
<i>Table 3.4: Observed values of bond length and bond angle of $\text{Ho}_2\text{Ti}_2\text{O}_7$, $\text{Dy}_2\text{Ti}_2\text{O}_7$, $\text{Ho}_2\text{Ti}_{1.85}\text{Fe}_{0.15}\text{O}_7$ and $\text{Ho}_{1.8}\text{Fe}_{0.2}\text{Ti}_2\text{O}_7$ obtained from Rietveld refinement.</i>	73
<i>Table 3.5: Details of crystallographic parameters and goodness of fit as obtained from Rietveld refinement of low temperature synchrotron x-ray diffraction $\text{Ho}_2\text{Ti}_2\text{O}_7$.</i>	79
<i>Table 3.6: Details of crystallographic parameters and goodness of fit as obtained from Rietveld refinement of low temperature synchrotron x-ray diffraction $\text{Dy}_2\text{Ti}_2\text{O}_7$.</i>	80
<i>Table 5.1: Composition, the length of unit cell volume, variable coordinate x in the 48f-site, reliable parameters (R_p, R_{wp} and R_{exp}) and goodness of fit χ^2.</i>	99
<i>Table 5.2: Values of bond length and bond angle as obtained by the Rietveld refinement for $\text{Dy}_{2-x}\text{Fe}_x\text{Ti}_2\text{O}_7$.</i>	99
<i>Table 6.1: Details of refinement as obtained from high-resolution x-ray diffraction. Compound, lattice constant, variable coordinate x of 48f-site, R-factors (R_p, R_{wp} and R_e) and goodness of fit χ^2.</i>	117