

LIST OF FIGURES

S. No.	Details	Page No.
Chapter 1		1
Figure 1.1	Prediction of International Energy Outlook till 2050	2
Figure 2.2	World energy Outlook (2019)	3
Figure 3.3	Emission profile of fossil fuel combustion	4
Figure 4.4	Global CO ₂ emission per region/country report (updated in 2019)	4
Figure 5.5	Cumulative Installed power capacity Mix (%) in India	8
Figure 6.6	An illustration of biofuel production methods	11
Figure 7.7	Stoichiometrically balanced transesterification reaction	14
Figure 8.8	A depiction of various feedstocks	15
Chapter 2		
Figure 2.1	Transesterification reaction protocol	41
Chapter 3		
Figure 3.1	Thermal curves of crude catalyst prepared by different synthesis routes	58
Figure 3.2	X- Ray diffractogram of calcined catalyst at 800° C (a), BLOA2 (b), BLOA1 (c), BLOC and (d), BLOS	60
Figure 3.3	SEM image of uncalcined catalyst of BLOA2 (a) and calcined catalysts at 800°C: BLOS (b), BLOC (c), BLOA1 (d), and BLOA2 (e)	63
Figure 3.4	Particle size analysis of the catalyst samples using SEM image.	64
Figure 3.5	EDXS histogram of calcined BLOA2 catalyst	64
Figure 3.6	FT-IR spectra of (a): BLOC, (b): BLOS, (c): BLOA1, (d): BLOA2	65
Figure 3.7	N ₂ adsorption-desorption isotherm profile of the catalyst samples	67
Figure 3.8	Effect of synthesis route on FAME conversion	68

Figure 3.9	Optimization plots (a) Effect of alcohol to oil molar ratio alongwith different catalyst dose (b) Effect of reaction temperature, (c) Effect of reaction time on MOME conversion, (d) Effect of stirring speed	70
Figure 3.10	(a) $-\ln(1-C_{ME})$ versus reaction time (min) plot at different temperatures, (b) Arrhenius plot $\ln k$ versus $1/T$ (K^{-1}) for transesterification of Mahua oil by BLOA2, (c) Eyring plot	73
Figure 3.11	Catalyst reusability test profile of BLOA2	75
Figure 3.12	FT-IR spectra for comparative study of fresh BLOA2 and used catalyst	75
Figure 3.13	An illustration of mechanism of transesterification reaction	76
Figure 3.14	GC-MS of methyl ester derived from Madhuca longifolia oil	78
Figure 3.15	1H -NMR spectra of methyl ester produced at optimized conditions of all process parameters	79
Figure 3.16	^{13}C - NMR spectra of methyl ester produced at optimized conditions of all process parameters	79
Figure 3.17	ATR FT-IR spectra for methyl ester prepared at most optimized condition of process variables	80
Chapter 4		
Figure 4.1	Thermal curves of crude catalyst	86
Figure 4.2	X- Ray diffractogram of perovskite sample calcined for different time intervals	88
Figure 4.3	X- Ray diffractogram of calcined catalysts with Ba/Ce stoichiometric ratio	89
Figure 4.4	SEM image of synthesized catalyst; (a): uncalcined perovskite sample, (b): Ba/Ce: 2/1,(c):Ba/Ce: 1/2, (d) Ba/Ce: 1/1(perovskite)	90
Figure 4.5	EDXS histogram of perovskite catalyst	91
Figure 4.6	FT-IR spectra of perovskite material calcined at different time interval	92
Figure 4.7	FT-IR spectra of calcined catalysts with different Ba/Ce stoichiometric ratios	93

Figure 4.8	a)N ₂ adsorption-desorption isotherm (BET) linear plot; (b) BJH adsorption cumulative pore volume; (c) optimization plot of calcination period for perovskite synthesis ;(d) plot inferring the metals stoichiometric ration on FAME conversion	94
Figure 4.9	Optimization plot showing the impact of various process variables on KOME conversion (a) Synergistic effect of catalyst dose with oil to methanol molar ratio;(b) Reaction temperature;(c) Reaction time;(d) Stirring speed	95
Figure 4.10	(a) -ln (1-C _{ME}) versus reaction time (min) plot at different temperatures using most optimized conditions, (b) Arrhenius plot lnk versus 1/T (K ⁻¹) for KOME synthesis from its crude oil,(c) Eyring plot	100
Figure 4.11	Reusability test of perovskite catalyst	102
Figure 4.12	FT-IR spectra for comparative study of fresh barium cerate and recycled catalyst	103
Figure 4.13	GC-MS profile of fatty acid methyl ester derived from <i>Milletia pinnatta</i> oil	104
Figure 4.14	(a) ¹ H- NMR spectra of fatty acid methyl ester obtained at most optimized condition of transesterification;(b) ¹³ C-NMR of methyl ester produced at optimized conditions of transesterification	106
Figure 4.15	ATR FT-IR spectra of fatty acid methyl ester derived from <i>Milletia pinnata</i> oil	107
Chapter 5		
Figure 5.1	TGA /DTA curve of uncalcined catalyst	112
Figure 5.2	X- Ray diffractogram of prepared catalyst calcined at 900°C for (a) 1 h, (b) 2 h, (c) 3 h, and (d) 4 h	114
Figure 5.3	SEM images of (a) uncalcined catalyst and calcined catalyst at 900°C for (b) 1 h, (c) 2 h, (d) 3 h, and (e) 4 h	115
Figure 5.4	Particle size analysis of the catalyst using SEM image	116
Figure 5.5	EDXS histogram of calcined catalyst at 900°C for 4h	116
Figure 5.6	(a) N ₂ adsorption-desorption profile of the catalyst; (b) BJH adsorption pore size distribution of the catalyst	117
Figure 5.7	X-Ray photoelectron spectra of the synthesized catalyst (a) Sr XPS 3d spectra, (b) La XPS 3d spectra, (c) O XPS 1s	119

	spectra of strontium lanthanum mixed metal oxide	
Figure 5.8	FT-IR spectra of calcined catalyst at different calcination time intervals	120
Figure 5.9	An illustration of effect of (a) calcination time duration, (b) catalyst dose and methanol to oil molar ratio, (c) reaction temperature, (d) reaction time, (e) agitation speed	122
Figure 5.10	(a) $-\ln(1-C_{ME})$ versus reaction time (min) plot for pseudo-first order kinetics,(b) Arrhenius plot,(c) Eyring plot	127
Figure 5.11	Catalyst reusability profile	128
Figure 5.12	FT-IR spectra for comparative study of (a) fresh and (b) used catalyst	129
Figure 5.13	GC-MS profile of fatty acid methyl ester	130
Figure 5.14	$^1\text{H-NMR}$ and $^{13}\text{C-NMR}$ spectra of methyl ester prepared at optimized condition of process variables	132
Figure 5.15	ATR FT-IR spectra of synthesized fatty acid methyl ester prepared at optimized condition of process variables	133
Chapter 6		
Figure 6.1	TGA-DTA of crude Sr-Ti mixed metals oxide as solid catalyst	141
Figure 6.2	X-Ray diffractogram of prepared Sr-Ti mixed metal oxide with different atomic ratio (1:2, 1:1, 2:1, 3:1, 4:1, and 5:1)	142
Figure 6.3	SEM images of prepared Sr-Ti mixed metal oxide with different atomic ratio (1:2, 1:1, 2:1, 3:1, 4:1, and 5:1)	144
Figure 6.4	(a) EDX spectra, (b) particle size distribution inferred from SEM image,(c) N_2 adsorption- desorption profile of the catalyst; (d) BJH adsorption pore size distribution of the catalyst	145
Figure 6.5	FT-IR spectra of prepared catalyst calcined at different temperatures	146
Figure 6.6	XPS analysis of catalyst i.e. Sr-Ti mixed metal oxide with atomic ratios (4:1)(a) Sr 3d spectra,(b) Ti 2p spectra,(c) O 1s spectra (d) wide scan full spectrum of catalyst	148
Figure 6.7	Optimization of Sr/Ti atomic ratios for highest FAME conversion	152

Figure 6.8	(a) Surface plot of FAME conversion vs methanol to oil molar ratio, catalyst dose, (b) Contour plot of FAME conversion vs methanol to oil molar ratio, catalyst dose (c) Surface plot of FAME conversion vs reaction time, methanol to oil molar ratio (d) Contour plot of FAME conversion vs reaction time, methanol to oil molar ratio (e) Surface plot of FAME conversion vs reaction time, catalyst dose (f) Contour plot of FAME conversion vs reaction time, catalyst dose	157
Figure 6.9	Optimization plot of process variables in transesterification reaction for methyl ester production using Sr-Ti mixed metal oxide with atomic ratio of 4:1	159
Figure 6.10	Endurance profile of catalyst efficiency over multiple recycling runs	160
Figure 6.11	FT-IR spectra of spent catalyst before and after regeneration process	161
Figure 6.12	Schematic illustration of reaction mechanism followed by catalyst	162
Figure 6.13	(a) $-\ln(1-C_{ME})$ versus reaction time (min) plot for Pseudo-first order kinetics, (b) Arrhenius plot, (c) Eyring plot	164
Figure 6.14	GC-MS profile for the composition of synthesized FAME	165
Figure 6.15	$^1\text{H-NMR}$ spectra of methyl ester produced at optimized conditions of all process parameters (catalyst dose, 1.0 wt%; methanol to oil molar ratio, 11:1; reaction temperature, 65°C; reaction time, 80min; agitation speed, 600 rpm)	166
Figure 6.16	ATR FT-IR spectrum of methyl ester prepared at optimum condition	167