

List of Tables

Table No.	Captions	Page No.
Table 3.1:	Experimentally observed and theoretically calculated atomic percentages for the Nd, Ba, Mn and Ti atoms present in $\text{Nd}_{0.7}\text{Ba}_{0.3}\text{Mn}_{1-x}\text{Ti}_x\text{O}_3$ ($0 \leq x \leq 0.30$) perovskite manganites obtained from EDS measurements.	85
Table 3.2:	Lattice parameters, unit cell volume, average bond-length and bond-angle and χ^2 for $\text{Nd}_{0.7}\text{Ba}_{0.3}\text{Mn}_{1-x}\text{Ti}_x\text{O}_3$ ($0 \leq x \leq 0.30$) manganites as determined from Rietveld structure refinement. Average crystallite size estimated using Debye-Scherrer formula (d_x) and particle size from SEM micrographs (d_{SEM}) are also listed.	92
Table 3.3:	The magnetic parameters for $\text{Nd}_{0.7}\text{Ba}_{0.3}\text{Mn}_{1-x}\text{Ti}_x\text{O}_3$ manganites with $0 \leq x \leq 0.30$. The Curie temperature T_C , Curie-Weiss temperature θ_{CW} , Curie constant C , experimental effective paramagnetic moment [$\mu_{\text{eff}}^{\text{exp}} (\mu_B)$], theoretical effective paramagnetic [$\mu_{\text{eff}}^{\text{theo}} (\mu_B)$], critical exponent for magnetic susceptibility (γ) and AFM fraction present in the sample.	99
Table 4.1:	Refined lattice parameters, unit cell volume and χ^2 for $P4mm$ and $I4/mcm$ phases of $\text{Nd}_{0.7}\text{Ba}_{0.3}\text{Mn}_{1-x}\text{Ti}_x\text{O}_3$ manganites with $x = 0.30, 0.40$ and 0.50 obtained from Rietveld structure refinement.	120
Table 4.2:	Curie temperature (T_C), Curie-Weiss temperature (θ_{CW}), Curie constant (C), theoretical and experimental values of the effective paramagnetic moments ($\mu_{\text{eff}}^{\text{theo}}$) ($\mu_{\text{eff}}^{\text{exp}}$), coercive field (H_C) and remnant magnetization (M_r) for $\text{Nd}_{0.7}\text{Ba}_{0.3}\text{Mn}_{1-x}\text{Ti}_x\text{O}_3$ with $x = 0.40$ and 0.50 .	126
Table 5.1:	Average values of particle & crystalline size, lattice strain, lattice constants and unit cell volume of NBMTO-10 manganite samples calcined at different temperatures.	140
Table 5.2:	Curie temperature T_C ; Curie-Weiss temperature θ_{CW} ; Curie constant C ; experimental $\mu_{\text{eff}}^{\text{exp}}$ and theoretical $\mu_{\text{eff}}^{\text{theo}}$ effective paramagnetic moments; for NBMTO-10 samples of various particle sizes.	150
Table 5.3:	Lattice parameters and unit cell volume for bulk sample (C13) of $\text{Nd}_{0.7}\text{Ba}_{0.3}\text{Mn}_{0.9}\text{Ti}_{0.1}\text{O}_3$ manganite at various temperatures.	160

Table 6.1:	Experimental and theoretical atomic percentages obtained from EDS spectrum for nano and bulk samples of LBMO-40 manganite.	169
Table 7.1:	Experimentally observed and theoretically calculated atomic percentages for the La, Ba, Mn and Ti atoms present in $\text{La}_{0.6}\text{Ba}_{0.4}\text{Mn}_{1-x}\text{Ti}_x\text{O}_3$ ($0.02 \leq x \leq 0.08$) manganites obtained from EDS measurements.	187
Table 7.2:	Particle size, lattice parameter, unit cell volume, bond-lengths ($d_{\text{Mn-O}}$ and $d_{\text{La-O}}$) and χ^2 for LBMTO manganites with $x = 0.02, 0.04, 0.06$ and 0.08 for $Pm\bar{3}m$ space group.	189
Table 7.3:	The magnetic parameters for LBMTO manganites with $x = 0.02, 0.04, 0.06$ and 0.08 . The Curie temperature T_C , Curie-Weiss temperature θ_{CW} , Curie constant C , the experimental effective paramagnetic moment $\mu_{\text{eff}}^{\text{exp}} (\mu_B)$, the theoretical effective paramagnetic $\mu_{\text{eff}}^{\text{theo}} (\mu_B)$, Remnant magnetization M_r , Coercive field H_C , experimental saturation moment $\mu_{\text{sat}}^{\text{exp}} (\mu_B)$ and theoretical saturation moment $\mu_{\text{sat}}^{\text{the}} (\mu_B)$.	200