## LIST OF FIGURES

Figure No.	Figure Description	Page No.
Figure 1.1	Proposed hypotheses	14
Figure 2.1.1	Proposed hypothesis of long term exposure to EMR causes anxiety-like symptom	18
Figure 2.1.2	Electromagnetic radiation exposure system and design	20
Figure 2.1.3	Schematic representation of the experimental design. '+' denotes experiment performed	21
Figure 2.1.4	Effect of EMR on elevated plus maze to estimate the anxiety- like behaviour in laboratory animals shows the changes in the percentage in open arm entries (A) time spent (B) and number of total arm entries (C) in elevated plus maze test respectively	30
Figure 2.1.5.1	The effect of long term exposure of EMR on open field test to evaluate anxiety-like behavior in rats caused behavioral changes (A) ambulation (B) rearing (C) grooming and (D) number of central square crossing in open field test	33
Figure 2.1.5.2	Second set of experiment on the effect of long term exposure of EMR on open field test to evaluate anxiety-like behavior in rats caused behavioral changes (A) ambulation (B) rearing (C) grooming and (D) number of central square crossing in open field test	33
Figure 2.1.6.1	Effect of EMR-900, 1800 and 2450 MHz on hole board test, (A) head dip, (B) sniffing (C) head dip/sniffing and (D) number of square crossed to evaluate anxiety-like behavior in rats	35
Figure 2.1.6.2	Second set of Experiment on Effect of EMR-900, 1800 and 2450 MHz on hole board test, (A) head dip, (B) sniffing (C) head dip/sniffing and (D) number of square crossed to evaluate anxiety-like behavior in rats	35
Figure 2.1.7	EMR-2450 MHz increased plasma corticosterone levels in experimental animals	36
Figure 2.1.8	Correlation between plasma corticosterone and percentage arm entries in EMR subjected rats	37

Figure 2.1.9	Effect of EMR on mitochondrial membrane potential (MMP) in amygdalar region of experimental rats	38
Figure 2.1.10	Effect of EMR-900, 1800 and 2450 MHz on (A) Catalase and (B) SOD activities in amygdala	40
Figure 2.1.11	Effect of EMR on protein expression of (A) CRH-2, GR and $\beta$ - actin, (B) represents the ratio of CRH-2/ $\beta$ -actin and (C) represents the ratio of GR/ $\beta$ -actin in amygdala of rats	41
Figure 2.1.12	Effect of EMR on the expression of cytoplasmic Bax, Bcl2 and their ratio in amygdalar tissue in brain	42
Figure 2.1.13	Effect of EMR on the expression of mitochondrial Bax, Bcl2 and their ratio in amygdalar tissue in brain	43
Figure 2.1.14	The effect of EMR- 900, 1800 and 2450 MHz on protein expression of (A) Cytochrome-C and (B) caspase-9 in amygdalar tissue	44
Figure 2.1.15	Effect of EMR on histopathology of amygdalar brain tissues (A) Control (B) EMR-900MHz (C) EMR-1800 MHz (D) EMR- 2450MHz and (B) Quantification value of neuronal damages	45-46
Figure 2.1.16	Effect of EMR (900, 1800 and 2450 MHz) on the pattern of apoptotic and necrotic cells in amygdala by using flow cytometric analysis. (A) Quadrants are labeled as Q1, Q2, Q3 and Q4 which denotes necrosis, late apoptosis, early apoptosis and live cells respectively, (B) shows the percentage of necrosis and apoptosis in amygdala in response to exposure with EMR (900, 1800 and 2450 MHz)	46-47
Figure 2.1.17	Summary	54
Figure 2.2.1	Proposed hypothesis of long term exposure to EMR causes depressive-like symptoms	58
Figure 2.2.2	Schematic representation of the experimental design. '+' denotes experiment performed	60
Figure 2.2.3	Effect of EMR-900, 1800 and 2450 MHz on (A) immobility period in forced swim test and (B) immobility period in tail	65

	suspension test in rats	
Figure 2.2.4	The effect of EMR (900, 1800 and 2450 MHz) on sucrose consumption behavior in experimental rats	66
Figure 2.2.5	The effect of sub-chronic exposure of EMR (900, 1800 and 2450 MHz) on cortical blood flow in rats	67
Figure 2.2.6	The effect of EMR on the level of VEGF in cortical region of brain in rats	68
Figure 2.2.7	Effect of EMR-900, 1800 and 2450 MHz on expression of BDNF mRNA in PFC of experimental rats	70
Figure 2.2.8	Effect of EMR-900, 1800 and 2450 MHz on flow cytometric analysis as four quadrants labeled as Q1, Q2, Q3 and Q4 which denotes necrosis, late apoptosis, early apoptosis and live cells respectively	71
Figure 2.2.8	Summary	76
Figure 2.3.1	Proposed hypothesis of long term exposure to EMR causes learning and memory deficits	80
Figure 2.3.2	Schematic representation of the experimental design. '+' denotes experiment performed	82
Figure 2.3.3	Effect of EMR (900, 1800, and 2450 MHz) exposure ensuing alterations in the total arm entries in trials 1 and 2 (curiosity; A), total novel arm entries (spatial recognition memory) (B), and coping behavior to novel arm (anxiety-like behavior; C) in Y-maze paradigm	88
Figure 2.3.4	EMR (900, 1800 and 2450 MHz)-induced alterations in the spatial recognition memory on D-1 (A), D-7 (B), D-14 (C), D-21 (D) and D-28 (E)	90
Figure 2.3.5	Effect of EMR (900, 1800 and 2450 MHz)-induced changes in concentration of Ach and AchE activity in hippocampus	91
Figure 2.3.6	The effect of EMR (900, 1800 and 2450 MHz)-induced changes in the level of $\beta$ Amyloid in brain tissues of rodents	92
Figure 2.3.7	The effect of discrete range of EMR induced alterations in MMP in EMR subjected rats	93
Figure 2.3.8	Effect of EMR (900, 1800 and 2450 MHz)-induced alterations in the levels of (A) cytochrome-C, (B) caspase-9 and (C) caspase-3 in hippocampus	94
Figure 2.3.9	Summary	101
Figure 3.1	Proposed hypothesis of long term exposure to EMR alters	105

	pharmacokinetics of fluoxetine and liver dysfunction in rats.	
Figure 3.2	Schematic representation of the experimental design. '+' denotes experiment performed	107
Figure 3.3	Effect of fluoxetine (5, 10 and 20 mg/kg) exhibited antidepressant activity in EMR-2450 MHz exposed rats during FST and TST paradigm	111
Figure 3.4	The effect of fluoxetine (5, 10 and 20mg/kg) exhibited antidepressant activity in EMR-2450 MHz exposed rats during SPT paradigms.	113
Figure 3.5	EMR-2450 MHz altered level of serotonin and dopamine of PFC in Fluoxetine (5, 10 and 20 mg/kg) treated rats	114
Figure 3.6	The effect of EMR on the level of CYP2D6 in liver and PFC in fluoxetine treated rats	115
Figure 3.7	The effect of fluoxetine on the level of ALT and AST in long term exposure of EMR in liver of rats	118
Figure 3.8	Summary	123
Figure 4.1	Proposed hypothesis of long term exposure to EMR causes gastric integrity and pharmacology in rats	128
Figure 4.2	Schematic representation of the experimental design. '+' denotes experiment performed	130
Figure 4.3	Effect of repeated exposure of EMR-2450 MHz on ulcer index in rat stomach (A) Control (B) EMR-2450MHz (E) (C) EMR- 2450 MHz (E1) (D) EMR-2450MHz (E2) (E) EMR-2450MHz (E3) (B) Shows changes in ulcer index after long term exposure of EMR in rat stomach. (A) Control (B) EMR- 2450MHz (E) (C) E1+OMZ-10 mg (D) E2+OMZ-20 mg (E) E3+OMZ-30 mg	139
Figure 4.4	The effect of EMR (2450 MHz) on mucus content in OMZ treated gastric ulcer in experimental rodents	140
Figure 4.5	Effect of repeated exposure to EMR-2450 MHz on $H^+/K^+$ ATPase activity in experimental rats	141
Figure 4.6	Effect of repeated exposure to EMR-2450 MHz on VEGF level in animals	143
Figure 4.7	Effect of repeated exposure to EMR-2450 MHz on (A) TNF-α, (B) IL-6 and (C) IL-10 in experimental rats	144

Figure 4.8	Repeated exposure of EMR showed complete thrashing of outer layer of gastric mucosa leads to necrosis and hemorrhage in experimental rats	145
Figure 4.9	Summary	150
Figure 5.1	Proposed hypothesis of Effect of EMR on cardiovascular system in rats.	153
Figure 5.2	Schematic representation of the experimental design. '+' denotes experiment performed	155
Figure 5.3	Effects of EMR-900, 1800 and 2450 MHz on heart rate (A) and mean arterial pressure (B) in experimental rats.	159
Figure 5.4	Effects of EMR-900, 1800 and 2450 MHz on electrocardiography analyses of heart in experimental rats	160
Figure 5.5	Effects of EMR-900, 1800 and 2450 MHz on cardiac blood flow in experimental rats	161
Figure 5.6	Effects of EMR-900, 1800 and 2450 MHz on morphological changes of cardiac tissues by using SEM in experimental rats	162
Figure 5.7	Summary	165
Figure 6.1	Conclusion	169