

# *Bibliography*

**BIBLIOGRAPHY**

Al-Soud, Y. A., Al-Sadoni, H. H., Saeed, B., Jaber, I. H., Beni-Khalid, M. O., Al-Masoudi, N. A., Abdul-Kadir, T., Colla, P. L., Busonera, B. and Sanna, T. (2008). Synthesis and in vitro antiproliferative activity of new benzothiazole derivatives. *Arkivoc*, 15, 225-238.

Ali, A., Taylor, G. E. and Graham, D. W. Int PCT, Appl WO 20011028561, 2001.

Amir, M., Asif, S., Ali, I. and Hassan, M. Z. (2012). Synthesis of benzothiazole derivatives having acetamido and carbothioamido pharmacophore as anticonvulsant agents. *Med. Chem. Res.*, 21, 2661-2670.

Anzini, M., Chelini, A., Mancini, A., Cappelli, A., Frosini, M., Ricci, L., Valoti, M., Magistretti, J., Castelli, L., Giordani, A. (2010). Synthesis and biological evaluation of amidine, guanidine, and thiourea derivatives of 2-amino(6-trifluoromethoxy)benzothiazole as neuroprotective agents potentially useful in brain diseases. *J. Med. Chem.*, 53, 734-744.

Arora, P., Das, S., Ranawat, M.S., Arora, N., Gupta, M.M., (2010). Synthesis and biological evaluation of some novel chromene- 2-one derivatives for antipsychotic activity. *J. Chem. Pharm. Res.*, 2, 317–323.

Atamanyuk, D., Faivre, F., Oxoby, M., Ledoussal, B., Drocourt, E., Moreau, F. and Gerusz, V. (2013). Vectorization efforts to increase Gram-negative intracellular drug concentration: a case study on HIdE-K inhibitors. *J. Med. Chem.*, 56, 1908-1921.

Aydemir, N. and Bilaloglu, R. (2003). Genotoxicity of two anticancer drugs, gemcitabine and topotecan, in mouse bone marrow in vivo. *Mutat. Res.*, 537, 43-51.

Balamurugan, R., Stalin, A. and Ignacimuthu, S. (2012). Molecular docking of gamma-sitosterol with some targets related to diabetes. *Eur. J. Med. Chem.*, 47, 38-43.

Bandyopadhyay, P., Sathe, M., Ponmariappan, S., Sharma, A., Sharma, P., Srivastava, A. K. and Kaushik, M. P. (2011). Exploration of in vitro time point quantitative evaluation of newly synthesized benzimidazole and benzothiazole derivatives as potential antibacterial agents. *Bioorg. Med. Chem. Lett.*, 21, 7306-7309.

Bayer CropScience AG. Cycloalkylamidobenzoxazoles and benzothiazoles as fungicides and their preparation and use for fighting harmful microorganisms in plants. EP2277869, 2011.

Bayer CropScience AG. Substituted benzoxazoles and benzothiazoles as fungicides and their preparation and use for fighting harmful microorganisms. EP2277870, 2011.

- Beneteau, V., Besson, T., Guillard, J., Leonce, S. and Pfeiffer, B. (1999). Synthesis and in vitro antitumour evaluation of benzothiazole-2-carbonitrile derivatives. *Eur. J. Med. Chem.*, 34, 1053-1060.
- Ben-Alloum, A., Bakkas, S. and Soufiaoui, M. (1997). Nouvelle voie de synthase des 2-arylbenzothiazoles transfert d'electrons active par micro-ondes. *Tetrahedron Lett.*, 38, 6395-6396.
- Bergman, J. M., Coleman, P. J., Cox, C. D., Hartman, G. D., Lindsley, C., Mercer, S. P., Roecker, A. J. and Whitman, D. B. Proline bis-amide orexin receptor antagonists. Int PCT Appl WO 2006127550, 2006.
- Bharti, S. K., Nath, G., Tilak, R. and Singh, S. K. (2010a). Synthesis, anti-bacterial and anti-fungal activities of some novel Schiff bases containing 2, 4-disubstituted thiazole ring. *Eur. J. Med. Chem.*, 45, 651-660.
- Bharti, S. K., Patel, S. K., Nath, G., Tilak, R. and Singh, S. K. (2010b). Synthesis, characterization, DNA cleavage and in vitro antimicrobial activities of copper (II) complexes of Schiff bases containing a 2, 4-disubstituted thiazole. *Transit. Metal Chem.*, 35, 917-925.
- Bhusari, K. P., Amnerkar, N. D., Khedekar, P. B., Kale, M. K. and Bhole, R. P. (2008). Synthesis and in vitro antimicrobial activity of some new 4-amino-N-(1, 3-benzothiazol-2-yl) benzenesulphonamide derivatives. *Asian J. Res. Chem.*, 1, 53-57.
- Bhuva, H. A. and Kini, S. G. (2010). Synthesis, anticancer activity and docking of some substituted benzothiazoles as tyrosine kinase inhibitors. *J. Mol. Graph. Modell.*, 29, 32-37.
- Bjornsti, M.-A. and Houghton, P. J. (2004). The TOR pathway: a target for cancer therapy. *Nat. Rev. Cancer*, 4, 335-348.
- Bolelli, K., Yalcin, I., Ertan-Bolelli, T., Ozgen, S., Kaynak-Onurdag, F., Yildiz, I. and Aki, E. (2012). Synthesis of novel 2-[4-(4-substitutedbenzamido/phenylacetamido) phenyl] benzothiazoles as antimicrobial agents. *Med. Chem. Res.*, 21, 3818-3825.
- Bondock, S., Fadaly, W. and Metwally, M. A. (2009). Enaminonitrile in heterocyclic synthesis: Synthesis and antimicrobial evaluation of some new pyrazole, isoxazole and pyrimidine derivatives incorporating a benzothiazole moiety. *Eur. J. Med. Chem.*, 44, 4813-4818.
- Bonnefous, C., Payne, J. E., Roppe, J., Zhuang, H., Chen, X., Symons, K. T., Nguyen, P. M., Sablad, M., Rozenkrants, N. and Zhang, Y. (2009). Discovery of inducible nitric oxide synthase (iNOS) inhibitor development candidate KD7332, part 1: Identification of a novel, potent, and

selective series of quinolinone iNOS dimerization inhibitors that are orally active in rodent pain models. *J. Med. Chem.*, 52, 3047-3062.

Bradshaw, T. D., Matthews, C. S., Cookson, J., Chew, E. H., Shah, M., Bailey, K., Monks, A., Harris, E., Westwell, A. D., Wells, G. et al. (2005). Elucidation of thioredoxin as a molecular target for antitumor quinols. *Cancer Res.*, 65, 3911-3919.

Bradshaw, T. D., Trapani, V., Vasselin, D. A. and Westwell, A. D. (2002). The aryl hydrocarbon receptor in anticancer drug discovery: friend or foe? *Curr. Pharm. Des.*, 8, 2475-2490.

Bradshaw, T. D. and Westwell, A. D. (2004). The development of the antitumour benzothiazole prodrug, Phortress, as a clinical candidate. *Curr. Med. Chem.*, 11, 1009-1021.

Bradshaw, T. D., Wrigley, S., Shi, D. F., Schultz, R. J., Paull, K. D. and Stevens, M. F. (1998). 2-(4-Aminophenyl)benzothiazoles: novel agents with selective profiles of in vitro anti-tumour activity. *Br. J. Cancer*, 77, 745-752.

Bujdakova, H. and Muckova, M. (1994). Antifungal activity of a new benzothiazole derivative against *Candida* in vitro and in vivo. *Int. J. Antimicrob. Agents*, 4, 303-308.

Caputo, R., Calabro, M. L., Micale, N., Schimmer, A. D., Ali, M., Zappala, M. and Grasso, S. (2012). Synthesis of benzothiazole derivatives and their biological evaluation as anticancer agents. *Med. Chem. Res.*, 21, 2644-2651.

Casewit, C. J., Colwell, K. S. and Rappe, A. K. (1992). Application of a universal force field to organic molecules. *J. Am. Chem. Soc.*, 114, 10035-10046.

Catalano, A., Carocci, A., Defrenza, I., Muraglia, M., Carrieri, A., Van Bambeke, F., Rosato, A., Corbo, F. and Franchini, C. (2013). 2-Aminobenzothiazole derivatives: Search for new antifungal agents. *Eur. J. Med. Chem.*, 64, 357-364.

Chen, J., Qian, L., Shen, Y., Chen, L. and Zheng, K. (2008). A QSAR study and molecular design of benzothiazole derivatives as potent anticancer agents. *Sci. China, Ser. B, Chem.*, 51, 111-119.

Chen, X., Sassano, M. F., Zheng, L., Setola, V., Chen, M., Bai, X., Frye, S. V., Wetsel, W. C., Roth, B. L. and Jin, J. (2012). Structure functional selectivity relationship studies of  $\beta$ -arrestin-biased dopamine D<sub>2</sub> receptor agonists. *J. Med. Chem.*, 55, 7141-7153.

Choi, S. J., Park, H. J., Lee, S. K., Kim, S. W., Han, G. and Choo, H. Y. (2006). Solid phase combinatorial synthesis of benzothiazoles and evaluation of topoisomerase II inhibitory activity. *Bioorg. Med. Chem.*, 14, 1229-1235.

Christoffersen, R.E. and Marr, J.J., The management of drug discovery: principles and practice, In: Chang, Y.L., Klug, Y.L., Wolff, M.E. Eds., Burger's Medicinal Chemistry and Drug Discovery, 5<sup>th</sup> edition, John Wiley and Sons, Inc., New York, USA, 1, 10-20, 1995.

Chua, M. S., Shi, D. F., Wrigley, S., Bradshaw, T. D., Hutchinson, I., Shaw, P. N., Barrett, D. A., Stanley, L. A. and Stevens, M. F. (1999). Antitumor benzothiazoles. 7. Synthesis of 2-(4-acylamino-phenyl)benzothiazoles and investigations into the role of acetylation in the antitumor activities of the parent amines. *J. Med. Chem.*, 42, 381-392.

Clark, D. E. (2003). In silico prediction of blood brain barrier permeation. *Drug Discov. Today*, 8, 927-933.

Cocco, M. T., Congiu, C., Omnis, V., Pellerano, M. L. and De Logu, A. (2002). Synthesis and antimycobacterial activity of new S-alkylisothiosemicarbazone derivatives. *Bioorg. Med. Chem.*, 10, 501-506.

Cumings, J., Ward, T. H., Ranson, M. and Dive, C. (2004). Apoptosis pathway-targeted drugs from the bench to the clinic. *Biochim. Biophys. Acta*, 1705, 53-66.

Daidone, G., Maggio, B. and Schillaci, D. (1990). Salicylanilide and its heterocyclic analogues. A comparative study of their antimicrobial activity. *Pharmazie*, 45, 441-442.

Damour, D. Preparation of 6-[(6-alkylor6-cycloalkyl-triazolopyridazin-3-yl)sulfanyl]-1,3-benzothiazole derivatives as c-Met inhibitors and their pharmaceutical compositions. WO121223, 2011.

Didziapetris, R., Japertas, P., Avdeef, A. and Petrauskas, A. (2003). Classification analysis of P-glycoprotein substrate specificity. *J. Drug Target.*, 11, 391-406.

Ding Z, Zhou W, Jiang J. Preparation of pyridothiazole and benzothiazole derivatives as antibacterial agents. CN103724360, 2014.

Donde, K. J., Patil, V. R., Utekar, S. S. and Malve, S. P. (2002). Synthesis, structural characterization and antimicrobial studies of hydrazone derivatives of 3-hydroxyimino-5-methyl-2-hexanone. *Acta Pol. Pharm.*, 59, 291-294.

El-Aziz, D. M. A., Etaiw, S. E. H. and Ali, E. A. (2013). Synthesis, spectroscopic, cytotoxic aspects and computational study of N-(pyridine-2-ylmethylene) benzo [d] thiazol-2-amine Schiff base and some of its transition metal complexes. *J. Mol. Str.*, 1048, 487-499.

Fabbro, D., Ruetz, S., Buchdunger, E., Cowan-Jacob, S. W., Fendrich, G., Liebetanz, J., Mestan, J., O'Reilly, T., Traxler, P. and Chaudhuri, B. (2002). Protein kinases as targets for anticancer agents: from inhibitors to useful drugs. *Pharmacol. Ther.*, 93, 79-98.

Fan, X., He, Y., Wang, Y., Xue, Z., Zhang, X. and Wang, J. (2011). A novel and practical synthesis of 2-benzoylbenzothiazoles and 2-benzylbenzothiazoles. *Tetrahedron Lett.*, 52, 899-902.

Friedrich, C. L., Moyles, D., Beveridge, T. J. and Hancock, R. E. W. (2000). Antibacterial action of structurally diverse cationic peptides on gram-positive bacteria. *Antimicrob. Agents Chemother.*, 44, 2086-2092.

Fukuoka, T., Ohya, S., Narita, T., Katsuta, M., Iijima, M., Masuda, N., Yasuda, H., Trias, J. and Nikaido, H. (1993). Activity of the carbapenem panipenem and role of the OprD (D2) protein in its diffusion through the *Pseudomonas aeruginosa* outer membrane. *Antimicrob. Agents Chemother.*, 37, 322-327.

Ghate, M. D., Baksariya, A., Gupta, N. and Patel, B. D. (2013). Pharmacophore Mapping, Virtual Screening and Molecular Docking Studies of DPP-IV Inhibitors. *Int. J. Drug Des. Discov.*, 4, 1231-1248.

Ghittoni, R., Accardi, R., Chiocca, S. and Tommasino, M. (2015). Role of human papillomaviruses in carcinogenesis. *E cancer medical science*, 9, 526.

Gilani, S. J., Nagarajan, K., Dixit, S. P., Taleuzzaman, M. and Khan, S. A. (2012). Benzothiazole incorporated thiazolidin-4-ones and azetidin-2-ones derivatives: Synthesis and in vitro antimicrobial evaluation. *Arabian J. Chem.*, doi:10.1016/j.arabjc.2012.04.004.

Gill, R. K., Singh, G., Sharma, A., Bedi, P. M. S. and Saxena, A. K. (2013). Synthesis, cytotoxic evaluation, and in silico studies of substituted N-alkylbromo-benzothiazoles. *Med. Chem. Res.*, 22, 4211-4222.

Gong, B., Hong, F., Kohm, C., Bonham, L. and Klein, P. (2004). Synthesis and SAR of 2-arylbenzoxazoles, benzothiazoles and benzimidazoles as inhibitors of lysophosphatidic acid acyltransferase-beta. *Bioorg. Med. Chem. Lett.*, 14, 1455-1459.

Gullick, W. J. (1991). Prevalence of aberrant expression of the epidermal growth factor receptor in human cancers. *British Med. Bull.*, 47, 87-98.

Gupta A and Rawat S. (2010). Synthesis and anti-inflammatory study of novel fluorobenzothiazole derivatives. *J. Chem. Pharm. Res.*, 2, 244-258.

- Hania, M. M. (2009). Synthesis of some imines and investigation of their biological activity. *J. Chem.*, 6, 629-632.
- Haugwitz, R. D., Angel, R. G., Jacobs, G. A., Maurer, B. V., Narayanan, V. L., Cruthers, L. R. and Szanto, J. (1982). Antiparasitic agents. 5. Synthesis and anthelmintic activities of novel 2-heteroaromatic-substituted isothiocyanatobenzoxazoles and-benzothiazoles. *J. Med. Chem.*, 25, 969-974.
- Havrylyuk, D., Mosula, L., Zimenkovsky, B., Vasylenko, O., Gzella, A. and Lesyk, R. (2010). Synthesis and anticancer activity evaluation of 4-thiazolidinones containing benzothiazole moiety. *Eur. J. Med.Chem.*, 45, 5012-5021.
- Haydon, D.J., Czaplewski, L.G., Palmer, N.J., et al., Preparation of benzothiazole derivatives as antibacterial agents. US0004221, 2012.
- Henriksen, G., Hauser, A. I., Westwell, A. D., Yousefi, B. H., Schwaiger, M., Drzezga, A. and Wester, H. J. (2007). Metabolically stabilized benzothiazoles for imaging of amyloid plaques. *J. Med. Chem.*, 50, 1087-1089.
- Hisano, T., Ichikawa, M., Tsumoto, K. and Tasaki, M. (1982). Synthesis of Benzoxazoles, Benzothiazoles and Benzimidazoles and Evaluation of Their Antifungal, Insecticidal and Herbicidal Activities. *Chem. Pharm. Bull.*, 30, 2996-3004.
- Hong, S.H., Hong, I.S., Han, E.J., Lee, G.H. Preparation of benzothiazole derivatives for the treatment of cancer. KR128693, 2013.
- Hong, S., Kim, J., Yun, S.-M., Lee, H., Park, Y., Hong, S.-S.and Hong, S. (2013). Discovery of new benzothiazole-based inhibitors of breakpoint cluster region-Abelson kinase including the T315I mutant. *J. Med. Chem.* 56, 3531-3545.
- Hutchinson, I., Bradshaw, T. D., Matthews, C. S., Stevens, M. F. and Westwell, A. D. (2003). Antitumour benzothiazoles. Part 20: 3'-cyano and 3'-alkynyl-substituted 2-(4'-aminophenyl)benzothiazoles as new potent and selective analogues. *Bioorg. Med. Chem. Lett.*, 13, 471-474.
- Jagessar, R. C. and Rampersaud, D. (2007). Amides as antimicrobial agents. *Life Sci. J.*, 4, 46-49.
- Jenkins, D. (2008). A review of cross-protection against oncogenic HPV by an HPV-16/18 AS04-adjuvanted cervical cancer vaccine: importance of virological and clinical endpoints and implications for mass vaccination in cervical cancer prevention. *Gynecol. Oncol.*, 110, S18-S25.

Jin, L., Song, B., Zhang, G., Xu, R., Zhang, S., Gao, X., Hu, D. and Yang, S. (2006). Synthesis, X-ray crystallographic analysis, and antitumor activity of N-(benzothiazole-2-yl)-1-(fluorophenyl)-O, O-dialkyl- $\alpha$ -aminophosphonates. *Bioorg. Med. Chem. Lett.*, 16, 1537-1543.

Johnson, A. C., Murphy, B. A., Matelis, C. M., Rubinstein, Y., Piebenga, E. C., Akers, L. M., Neta, G., Vinson, C. and Birrer, M. (2000). Activator protein-1 mediates induced but not basal epidermal growth factor receptor gene expression. *Mol. Medicine.*, 6, 17-27.

Joshi, S., Dewangan, R. P., Yadav, S., Rawat, D. S. and Pasha, S. (2012). Synthesis, antibacterial activity and mode of action of novel linoleic acid-dipeptide-spermidine conjugates. *Org. Biomol. Chem.*, 10, 8326-8335.

Kamal, A., Faazil, S., Ramaiah, M. J., Ashraf, M., Balakrishna, M., Pushpavalli, S., Patel, N. and Pal-Bhadra, M. (2013a). Synthesis and study of benzothiazole conjugates in the control of cell proliferation by modulating Ras/MEK/ERK-dependent pathway in MCF-7 cells. *Bioorg. Med. Chem. Lett.*, 23, 5733-5739.

Kamal, A., Reddy, K. S., Khan, M. N. A., Shetti, R. V., Ramaiah, M. J., Pushpavalli, S., Srinivas, C., Pal-Bhadra, M., Chourasia, M. and Sastry, G. N. (2010). Synthesis, DNA-binding ability and anticancer activity of benzothiazole/benzoxazolepyrrolo [2, 1-c][1, 4] benzodiazepine conjugates. *Bioorg. Med. Chem. Lett.*, 18, 4747-4761.

Kamal, A., Syed, M. A. H. and Mohammed, S. M. (2015). Therapeutic potential of benzothiazoles: a patent review (2010-2014). *Expert Opin. Ther. Pat.*, 25, 335-349.

Kamal, A., Tamboli, J. R., Nayak, V. L., Adil, S. F., Vishnuvardhan, M. and Ramakrishna, S. (2013b). Synthesis of pyrazolo [1, 5-a] pyrimidine linked aminobenzothiazole conjugates as potential anticancer agents. *Bioorg. Med. Chem. Lett.*, 23, 3208-3215.

Kamesaki, H. (1998). Mechanisms involved in chemotherapy-induced apoptosis and their implications in cancer chemotherapy. *Int. J. Hematol.*, 68, 29-43.

Karmakar, T., Kuang, Y., Neamati, N. and Baruah, J. B. (2013). Cadmium complexes and cocrystals of indium complexes of benzothiazole derivatives and anticancer activities of the cadmium complexes. *Polyhedron*, 54, 285-293.

Karthikeyan, C., Lee, C., Moore, J., Mittal, R., Suswam, E. A., Abbott, K. L., Pondugula, S. R., Manne, U., Narayanan, N. K. and Trivedi, P. (2015). IND-2, a pyrimido [1'', 2''': 1, 5] pyrazolo [3, 4-b] quinoline derivative, circumvents multi-drug resistance and causes apoptosis in colon cancer cells. *Bioorg. Med. Chem.*, 23, 602-611.



Kashiyama, E., Hutchinson, I., Chua, M. S., Stinson, S. F., Phillips, L. R., Kaur, G., Sausville, E. A., Bradshaw, T. D., Westwell, A. D. and Stevens, M. F. (1999). Antitumor benzothiazoles. 8. Synthesis, metabolic formation, and biological properties of the C- and N-oxidation products of antitumor 2-(4-aminophenyl)benzothiazoles. *J. Med. Chem.*, 42, 4172-4184.

Kasibhatla, S., Amarante-mendes, G.P, Finucane, D., Brunner, T., Bossy-Wetzel, E., Green, D.R. Analysis of DNA Fragmentation Using Agarose Gel Electrophoresis. *Cold Spring Harb Protoc.*, 2006, pdb.prot, 4429-4430.

Kok, S. H. L., Gambari, R., Chui, C. H., Yuen, M. C. W., Lin, E., Wong, R. S. M., Lau, F. Y., Cheng, G. Y. M., Lam, W. S. and Chan, S. H. (2008). Synthesis and anti-cancer activity of benzothiazole containing phthalimide on human carcinoma cell lines. *Bioorg. Med. Chem.*, 16, 3626-3631.

Komaratat, P. and Kates, M. (1975). The lipid composition of a halotolerant species of *Staphylococcus epidermidis*. *Biochim.Biophys. Acta*, 398, 464-484.

Koopman, G., Reutelingsperger, C. P., Kuijten, G. A., Keehnen, R. M., Pals, S. T. and Van Oers, M. H. (1994). Annexin V for flow cytometric detection of phosphatidylserine expression on B cells undergoing apoptosis. *Blood*, 84, 1415-1420.

Kumar, A., D'Souza, S. S., Gaonkar, S. L., Rai, K. M. L. and Salimath, B. P. (2008). Growth inhibition and induction of apoptosis in MCF-7 breast cancer cells by a new series of substituted-1, 3, 4-oxadiazole derivatives. *Invest. New Drugs*, 26, 425-435.

Kumar, A., Maurya, R. A. and Ahmad, P. (2009a). Diversity oriented synthesis of benzimidazole and benzoxa/(thia)zole libraries through polymer-supported hypervalent iodine reagent. *J. Comb. Chem.*, 11, 198-201.

Kumar, P., Narasimhan, B., Sharma, D., Judge, V. and Narang, R. (2009b). Hansch analysis of substituted benzoic acid benzylidene/furan-2-yl-methylene hydrazides as antimicrobial agents. *Eur. J. Med. Chem.*, 44, 1853-1863.

Kumbhare, R. M., Dadmal, T., Kosurkar, U., Sridhar, V. and Rao, J. V. (2012). Synthesis and cytotoxic evaluation of thiourea and N-bis-benzothiazole derivatives: a novel class of cytotoxic agents. *Bioorg. Med. Chem. Lett.*, 22, 453-455.

Kushi, L. H., Doyle, C., McCullough, M., Rock, C. L., Demark-Wahnefried, W., Bandera, E. V., Gapstur, S., Patel, A. V., Andrews, K. and Gansler, T. (2012). American Cancer Society

Guidelines on nutrition and physical activity for cancer prevention: reducing the risk of cancer with healthy food choices and physical activity. *CA Cancer J. Clin.*, 62, 30-67.

Latrofa, A., Franco, M., Lopedota, A., Rosato, A., Carone, D. and Vitali, C. (2005). Structural modifications and antimicrobial activity of N-cycloalkenyl-2-acylalkylidene-2, 3-dihydro-1, 3-benzothiazoles. *Il Farmaco*, 60, 291-297.

Lavoie, E. J., Parhi, A., Zhang, Y., et al., Preparation of benzothiazole derivatives and related compounds as antimicrobial agents. WO142712, 2013.

Leite, A. C. L., de M, M., Rodrigo, D., de O, C., Marcos, V., Hernandez, M. Z., Alves Pereira, V. r. R., Silva, R. O., Kiperstok, A. C. and da S Lima, M. (2007). Synthesis, Cruzain Docking, and in vitro Studies of Aryl Oxothiazolylylhydrazones against *Trypanosoma cruzi*. *Chem. Med. Chem.*, 2, 1339-1345.

Lerner, C. G. and Beutel, B. A. (2002). Antibacterial drug discovery in the post-genomics era. *Curr. Drug Targets Infect. Disord*, 2, 109-119.

Lewis, K. (2013). Platforms for antibiotic discovery. *Nat. Rev. Drug Discov.*, 12, 371-387.

Li, X., Liu, X., Wang, C., et al., 2-(4-Aminophenyl)benzothiazole-containing naphthalimide derivatives useful in treatment of cancers and their preparation. CN103450176, 2013.

Li, Y., Mei, L., Li, H., et al., Predation of 2-methoxy-3-substituted-sulfonylamino-5-(2-acetamido-6-benzothiazole)-benzamide derivatives as antitumor agents. CN103772317, 2014.

Li. X., Ma, L., Zhao. J., et al., Preparation of mercapto benzothiazole derivatives as antitumor agents. CN103435574, 2013.

Lokhandwala, S. R. and Desai, K. R. (2008). Novel organophosphorus compounds as potential antimicrobial agents. *Phosphorus, Sulfur, and Silicon*, 183, 1264-1271.

Luniss, C. J., Palmer, J. T., Pitt, GRW., et al., Benzothiazole derivatives as antibacterial agents and their preparation. WO138860, 2013.

MacKinnon, C. H., Lau, K., Burch, J. D., Chen, Y., Dines, J., Ding, X., Eigenbrot, C., Heifetz, A., Jaochico, A. and Johnson, A. (2013). Structure-based design and synthesis of potent benzothiazole inhibitors of interleukin-2 inducible T cell kinase (ITK). *Bioorg. Med. Chem. lett.*, 23, 6331-6335.

Maddila, S., Gorle, S., Seshadri, N., Lavanya, P. and Jonnalagadda, S. B. (2013). Synthesis, antibacterial and antifungal activity of novel benzothiazole pyrimidine derivatives. *Arabian J. Chem.*, doi:10.1016/j.arabjc.2013.04.003.

Mahran, M., William, S., Ramzy, F. and Sembel, A. (2007). Synthesis and in vitro Evaluation of New Benzothiazole Derivatives as Schistosomicidal Agents. *Molecules*, 12, 622.

Maleki, B. and Salehabadi, H. (2010). Ammonium chloride; as a mild and efficient catalyst for the synthesis of some 2-arylbenzothiazoles and bisbenzothiazole derivatives. *Eur. J. Chem.*, 1, 377-380.

Marta, M., Matthew, M., Leonard, A. and Seamus, F. (2011). Mechanisms of Antibiotic Resistance in Salmonella: Efflux Pumps, Genetics, Quorum Sensing and Biofilm Formation. *Lett. Drug Design Discov.*, 8, 114-123.

Mendelsohn, J. and Baselga, J. (2000). The EGF receptor family as targets for cancer therapy. *Oncogene*, 19, 6550-6565.

Milewski, S., Chmara, H., Andruszkiewicz, R., Borowski, E., Zaremba, M. and Borowski, J. (1988). Antifungal peptides with novel specific inhibitors of glucosamine 6-phosphate synthase. *Drugs Exp. Clin. Res.*, 14, 461-465.

Milewski, S. (2002). Glucosamine-6-phosphate synthase the multi-facets enzyme. *Biochim Biophys Acta*, 1597, 173-192.

Mishra, D., Naskar, S., Drew, M. G. B. and Chattopadhyay, S. K. (2006). Synthesis, spectroscopic and redox properties of some ruthenium (II) thiosemicarbazone complexes: structural description of four of these complexes. *Inorganica Chimica Acta*, 359, 585-592.

Mortimer, C. G., Wells, G., Crochard, J.-P., Stone, E. L., Bradshaw, T. D., Stevens, M. F. G. and Westwell, A. D. (2006). Antitumor benzothiazoles. 26. 1 2-(3, 4-dimethoxyphenyl)-5-fluorobenzothiazole (GW 610, NSC 721648), a simple fluorinated 2-arylbenzothiazole, shows potent and selective inhibitory activity against lung, colon, and breast cancer cell lines. *J. Med. Chem.*, 49, 179-185.

Mukherjee, A., Westwell, A. D., Bradshaw, T. D., Stevens, M. F., Carmichael, J. and Martin, S. G. (2005). Cytotoxic and antiangiogenic activity of AW464 (NSC 706704), a novel thioredoxin inhibitor: an in vitro study. *Br. J. Cancer*, 92, 350-358.

Navarrete-Vazquez, G., Paoli, P., Leon-Rivera, I., Villalobos-Molina, R., Medina-Franco, J. L., Ortiz-Andrade, R., Estrada-Soto, S., Camici, G., Diaz-Coutino, D., Gallardo-Ortiz, I. et al. (2009). Synthesis, in vitro and computational studies of protein tyrosine phosphatase 1B inhibition of a small library of 2-arylsulfonylaminobenzothiazoles with antihyperglycemic activity. *Bioorg. Med. Chem.*, 17, 3332-3341.

Noble, M. E. M., Endicott, J. A. and Johnson, L. N. (2004). Protein kinase inhibitors: insights into drug design from structure. *Science*, 303, 1800-1805.

Normanno, N., De Luca, A., Bianco, C., Strizzi, L., Mancino, M., Maiello, M. R., Carotenuto, A., De Feo, G., Caponigro, F. and Salomon, D. S. (2006). Epidermal growth factor receptor (EGFR) signaling in cancer. *Gene*, 366, 2-16.

Novo, D. J., Perlmutter, N. G., Hunt, R. H. and Shapiro, H. M. (2000). Multiparameter Flow Cytometric Analysis of Antibiotic Effects on Membrane Potential, Membrane Permeability, and Bacterial Counts of *Staphylococcus aureus* and *Micrococcus luteus*. *Antimicrob. agents chemother.*, 44, 827-834.

Oanh, D. T. K., Van Hai, H., Park, S. H., Kim, H.-J., Han, B.-W., Kim, H.-S., Hong, J.-T., Han, S.-B. and Nam, N.-H. (2011). Benzothiazole-containing hydroxamic acids as histone deacetylase inhibitors and antitumor agents. *Bioorg. Med. Chem.lett.*, 21, 7509-7512.

Okaniwa, M., Hirose, M., Arita, T., Yabuki, M., Nakamura, A., Takagi, T., Kawamoto, T., Uchiyama, N., Sumita, A. and Tsutsumi, S. (2013). Discovery of a selective kinase inhibitor (TAK-632) targeting pan-RAF inhibition: design, synthesis, and biological evaluation of C-7-substituted 1, 3-benzothiazole derivatives. *J. Med. Chem.*, 56, 6478-6494.

Okaniwa, M., Hirose, M., Imada, T., Ohashi, T., Hayashi, Y., Miyazaki, T., Arita, T., Yabuki, M., Kakoi, K. and Kato, J. (2012). Design and synthesis of novel DFG-out RAF/vascular endothelial growth factor receptor 2 (VEGFR2) inhibitors. 1. Exploration of [5, 6]-fused bicyclic scaffolds. *J. Med. Chem.*, 55, 3452-3478.

Orlov, V. N., Antonenko, Y. N., Bulychev, A. A. and Yaguzhinsky, L. S. (1994). Combination of the electrogenic ionophores, valinomycin and CCCP, can lead to non-electrogenic K<sup>+</sup>/H<sup>+</sup> exchange on bilayer lipid membranes. *FEBS lett.*, 345, 104-106.

Osario, T. M., DelleMonache, F., Chiaradia, L. D., Mascarello, A., Stumpf, T. R., Zanetti, C. R., Silveira, D. B., Barardi, C. I. R. M., Smania E., d. F. A. and Viancelli, A. (2012) Antibacterial activity of chalcones, hydrazones and oxadiazoles against methicillin-resistant *Staphylococcus aureus*. *Bioorg. Med. Chem. lett.*, 22, 225-230.

Ouyang, L., Huang, Y., Zhao, Y., He, G., Xie, Y., Liu, J., He, J., Liu, B. and Wei, Y. (2012). Preparation, antibacterial evaluation and preliminary structureactivity relationship (SAR) study of benzothiazol-and benzoxazol-2-amine derivatives. *Bioorg. Med. Chem. lett.*, 22, 3044-3049.

Padalkar, V. S., Borse, B. N., Gupta, V. D., Phatangare, K. R., Patil, V. S., Umape, P. G. and Sekar, N. (2011). Synthesis and antimicrobial activity of novel 2-substituted benzimidazole, benzoxazole and benzothiazole derivatives. *Arabian J. Chem.*, doi:10.1016/j.arabjc.2011.12.00.

Pandeya, S. N., Srivastava, T., Gangwar, M., Bihari, B. and Nath, G. (2012). Synthesis and antimicrobial activities of 1-Naphthylamine based acetophenone semicarbazones. *Med. Chem. Drug Discov.*, 3, 94-102.

Panicker, B., Oehlen, LJWM., Tarrant, J. G., et al., Preparation of benzo[d]thiazole compounds as cytochrome P450 inhibitors and therapeutic uses thereof. WO153192, 2011.

Patel, R. V. and Park, S. W. (2014). Access to a new class of biologically active quinoline based 1, 2, 4-triazoles. *Eur. J. Med. Chem.* 71, 24-30.

Patel, R. V., Patel, P. K., Kumari, P., Rajani, D. P. and Chikhalia, K. H. (2012). Synthesis of benzimidazolyl-1, 3, 4-oxadiazol-2ylthio-N-phenyl (benzothiazolyl) acetamides as antibacterial, antifungal and antituberculosis agents. *Eur. J. Med. Chem.*, 53, 41-51.

Pathak, P., Jolly, V. S. and Sharma, K. P. (2000). Synthesis and biological activities of some new substituted arylazo Schiff bases. *Oriental J. Chem.*, 16, 161-162.

Patrick, L.G., Drug discovery, design, and development, An introduction to medicinal chemistry, 2<sup>nd</sup> edition, Oxford University Press Inc., New York, 153-167, 2001.

Patterson, L. H. and Murray, G. I. (2002). Tumour cytochrome P450 and drug activation. *Curr. Pharm. Des.*, 8, 1335-1347.

Perin, D.D., Armargo, W.L.; Perin, D.R. Purification of Laboratory Chemicals, Pergamon, London, 1966.

Pieczonka, A. M., Strzelczyk, A., Sadowska, B., Mloston, G. and Staczek, P. (2013) Synthesis and evaluation of antimicrobial activity of hydrazones derived from 3-oxido-1H-imidazole-4-carbohydrazides. *Eur. J. Med. Chem.*, 64, 389-395.

Piscitelli, F., Ballatore, C. and Smith, A. B., 3rd. (2010). Solid phase synthesis of 2-aminobenzothiazoles. *Bioorg. Med. Chem. Lett.*, 20, 644-648.

Prabhu, P. P., Panneerselvam, T., Shastry, C. S., Sivakumar, A. and Pande, S. S. (2015). Synthesis and anticancer evaluation of 2-phenyl thiaolidinone substituted 2-phenyl benzothiazole-6-carboxylic acid derivatives. *J. Saudi Chem. Soc.*, 19, 181-185.

Prasad, S. B., Yadav, S. S., Das, M., Govardhan, H. B., Pandey, L. K., Singh, S., Pradhan, S. and Narayan, G. (2014). Down regulation of FOXO1 promotes cell proliferation in cervical cancer. *J. Cancer*, 5, 655.

Prasad, S. B., Yadav, S. S., Das, M., Modi, A., Kumari, S., Pandey, L. K., Singh, S., Pradhan, S. and Narayan, G. (2015). PI3K/AKT pathway-mediated regulation of p27Kip1 is associated with cell cycle arrest and apoptosis in cervical cancer. *Cellular Oncology*, 38, 215-225.

Racane, L., Pavelic, S.K., Nhili, R., Depauw, S., Paul-Constant, C., Ratkaj, I., David-Cordonnier, M.H., Pavelic, K., Tralic-Kulenovic, V. and Karminski-Zamola, G. (2013). New anticancer active and selective phenylene-bisbenzothiazoles: synthesis, antiproliferative evaluation and DNA binding. *Eur. J. Med. Chem.*, 63, 882-891.

Rai, G., Vyjayanti, V. N., Dorjsuren, D., Simeonov, A., Jadhav, A., Wilson I., D. M. and Maloney, D. J. (2012). Synthesis, biological evaluation, and structureactivity relationships of a novel class of apurinic/aprimidinic endonuclease 1 inhibitors. *J. Med. Chem.*, 55, 3101-3112.

Rao, A. J., Rao, P. V., Rao, V. K., Mohan, C., Raju, C. N. and Reddy, C. S. (2010). Microwave assisted one-pot synthesis of novel  $\alpha$ -aminophosphonates and their biological activity. *Bull. Korean Chem. Soc.*, 31, 1863-1868.

Reddy, P.V.G.; Lin, Y-W.; Chang, H-T. (2007). Synthesis of novel benzothiazole compounds with an extended conjugated system. *Arkivoc*, 15, 113-122.

Riadi, Y., Mamouni, R., Azzalou, R., El Haddad, M., Routier, S., Guillaumet, G. and Lazar, S. (2011). An efficient and reusable heterogeneous catalyst animal bone meal for facile synthesis of benzimidazoles, benzoxazoles, and benzothiazoles. *Tetrahedron Lett.*, 52, 3492-3495.

Sadjadi, S. and Sepehrian, H. (2011). Cu(OAc)<sub>2</sub>/MCM-41: An efficient and solid acid catalyst for synthesis of 2-arylbenzothiazoles under ultrasound irradiation. *Ultrasonics sonochemistry*, 18, 480-483.

Saeed, S., Rashid, N., Jones, P. G., Ali, M. and Hussain, R. (2010). Synthesis, characterization and biological evaluation of some thiourea derivatives bearing benzothiazole moiety as potential antimicrobial and anticancer agents. *Eur. J. Med. Chem.*, 45, 1323-1331.

Sahu, P. K., Gupta, S. K., Thavaselvam, D. and Agarwal, D. D. (2012). Synthesis and evaluation of antimicrobial activity of 4H-pyrimido[2,1-b]benzothiazole, pyrazole and benzyldiene derivatives of curcumin. *Eur. J. Med. Chem.*, 54, 366-378.

Scaltriti, M. and Baselga, J. (2006). The epidermal growth factor receptor pathway: a model for targeted therapy. *Clinical Cancer Res.*, 12, 5268-5272.

Schlessinger, J. (2002). Ligand-induced, receptor-mediated dimerization and activation of EGF receptor. *Cell*, 110, 669-672.

Schlessinger, J. (2004). Common and distinct elements in cellular signaling via EGF and FGF receptors. *Science*, 306, 1506-1507.

Seijas, J. A., Pilar Vazquez-Tato, M., Raquel Carballido-Reboredo, M., Crecente-Campo, J. and Romar-Lopez, L. (2007). Lawesson's reagent and microwaves: A new efficient access to benzoxazoles and benzothiazoles from carboxylic acids under solvent-free conditions. *Synlett*, 313-317.

Sharma, K. P., Jolly, V. S. and Phatak, P. (1998). Schiff bases and their derivatives as potential anticancer agents. *Ultra Scientist of Physical Sciences*, 10, 263-266.

Sharma, P. K., Kumar, M. and Mohan, V. (2010). Synthesis and antimicrobial activity of 2H-pyrimido [2, 1-b] benzothiazol-2-ones. *Res. Chem. Intermediates*, 36, 985-993.

Sheng C, Zhang W, Liu Y, et al., Substituted benzothiazoles as antifungal agents and their preparation, pharmaceutical compositions and use in the treatment of fungal infections. CN103193770, 2013.

Sheng, C., Zhu, J., Zhang, W., Zhang, M., Ji, H., Song, Y., Xu, H., Yao, J., Miao, Z. and Zhou, Y. (2007). 3D-QSAR and molecular docking studies on benzothiazole derivatives as *Candida albicans* N-myristoyltransferase inhibitors. *Eur. J. Med. Chem.*, 42, 477-486.

Shi, D.-F., Bradshaw, T. D., Wrigley, S., McCall, C. J., Lelieveld, P., Fichtner, I. and Stevens, M. F. G. (1996). Antitumor benzothiazoles. 3. 1 Synthesis of 2-(4-aminophenyl) benzothiazoles and evaluation of their activities against breast cancer cell lines in vitro and in vivo. *J. Med. Chem.*, 39, 3375-3384.

Shi, D.-Q., Rong, S.-F. and Dou, G.-L. (2010). Efficient synthesis of 2-arylbenzothiazole derivatives with the aid of a low-valent titanium reagent. *Synthetic Comm.*, 40, 2302-2310.

Siddiqui, N., Rana, A., Khan, S. A., Bhat, M. A. and Haque, S. E. (2007). Synthesis of benzothiazole semicarbazones as novel anticonvulsants "The role of hydrophobic domain". *Bioorg. Med. Chem. Lett.*, 17, 4178-4182.

Sims, P. J., Waggoner, A. S., Wang, C.-H. and Hoffman, J. F. (1974). Mechanism by which cyanine dyes measure membrane potential in red blood cells and phosphatidylcholine vesicles. *Biochemistry*, 13, 3315-3330.

Singh, M. K., Tilak, R., Nath, G., Awasthi, S. K. and Agarwal, A. (2013). Design, synthesis and antimicrobial activity of novel benzothiazole analogs. *Eur. J. Med. Chem.*, 63, 635-644.

Sleeb, B. E., Kersten, W. J. A., Kulasegaram, S., Nikolakopoulos, G., Hatzis, E., Moss, R. M., Parisot, J. P., Yang, H., Czabotar, P. E. and Fairlie, W. D. (2013). Discovery of potent and selective benzothiazole hydrazone inhibitors of Bcl-XL. *J. Med. Chem.*, 56, 5514-5540.

Solomon, V. R., Hu, C. and Lee, H. (2009). Hybrid pharmacophore design and synthesis of isatin benzothiazole analogs for their anti-breast cancer activity. *Bioorg. Med. Chem.*, 17, 7585-7592.

Song, E. Y., Kaur, N., Park, M. Y., Jin, Y., Lee, K., Kim, G., Lee, K. Y., Yang, J. S., Shin, J. H., Nam, K. Y. et al. (2008). Synthesis of amide and urea derivatives of benzothiazole as Raf-1 inhibitor. *Eur. J. Med. Chem.*, 43, 1519-1524.

Soni, B., Ranawat, M. S., Sharma, R., Bhandari, A. and Sharma, S. (2010). Synthesis and evaluation of some new benzothiazole derivatives as potential antimicrobial agents. *Eur. J. Med. Chem.*, 45, 2938-2942.

Sparks, R. B., Polam, P., Zhu, W., Crawley, M. L., Takvorian, A., McLaughlin, E., Wei, M., Ala, P. J., Gonnevill, L., Taylor, N. (2007). Benzothiazole benzimidazole (S)-isothiazolidinone derivatives as protein tyrosine phosphatase-1B inhibitors. *Bioorg. Med. Chem. Lett.*, 17, 736-740.

Spector, I., Palfrey, C. and Littauer, U. Z. (1975). Enhancement of the electrical excitability of neuroblastoma cells by valinomycin. *Nature*, 254, 121-124.

Sun, D., Hurdle, J. G., Lee, R., Lee, R., Cushman, M. and Pezzuto, J. M. (2012). Evaluation of flavonoid and resveratrol chemical libraries reveals abyssinone II as a promising antibacterial lead. *Chem Med Chem*, 7, 1541-1545.

Sun, Q., Wu, R., Cai, S., Lin, Y., Sellers, L., Sakamoto, K., He, B. and Peterson, B. R. (2011). Synthesis and biological evaluation of analogues of AKT (protein kinase B) inhibitor-IV. *J. Med. Chem.*, 54, 1126-1139.

Swinnen D, Jorand-Lebrun C, Grippi-Vallotton T, et al. Fused bicyclic derivatives as PI3 kinase inhibitors and their preparation and use for the treatment of diseases. WO100144, 2010.

Sybyl-X 1.2 (2010) <http://www.tripos.com>, St Louis.



Tabassum, S., Khan, R. A., Arjmand, F., Aziz, M., Juvekar, A. S. and Zingde, S. M. (2011). Carbohydrate-conjugate heterobimetallic complexes: synthesis, DNA binding studies, artificial nuclease activity and in vitro cytotoxicity. *Carbohydrate Res.*, 346, 2886-2895.

Thompson, C. B. (1995). Apoptosis in the pathogenesis and treatment of disease. *Science*, 267, 1456-1462.

Tomi, I. H. R., Tomma, J. H., Al-Daraji, A. H. R. and Al-Dujaili, A. H. (2015). Synthesis, characterization and comparative study the microbial activity of some heterocyclic compounds containing oxazole and benzothiazole moieties. *J. Saudi Chem. Soc.*, 19, 392-398.

Tosteson, D. C., Andreoli, T. E., Tieffenberg, M. and Cook, P. (1968). The effects of macrocyclic compounds on cation transport in sheep red cells and thin and thick lipid membranes. *The J Gen. Physiology.*, 51(5), 373-384.

Traxler, P., Bold, G., Buchdunger, E., Caravatti, G., Furet, P., Manley, P., O'Reilly, T., Wood, J. and Zimmermann, J. (2001). Tyrosine kinase inhibitors: from rational design to clinical trials. *Med. Res. Reviews*, 21, 499-512.

Verkleij, A. J., Zwaal, R. F., Roelofsen, B., Comfurius, P., Kastelijn, D. and van Deenen, L. L. (1973). The asymmetric distribution of phospholipids in the human red cell membrane. A combined study using phospholipases and freeze-etch electron microscopy. *Biochim. Biophys. Acta*, 323, 178-193.

Vicini, P., Geronikaki, A., Incerti, M., Busonera, B., Poni, G., Cabras, C. A. and La Colla, P. (2003). Synthesis and biological evaluation of benzo [d] isothiazole, benzothiazole and thiazole Schiff bases. *Bioorg. Med. Chem.*, 11, 4785-4789.

Wang, L., Doherty, G., Wang, X., et al., Preparation of substituted benzothiazoles as apoptosis-inducing agents for the treatment of cancer, immune and autoimmune diseases. WO055895, 2013.

Wang, M., Wang, L.-F., Li, Y.-Z., Li, Q.-X., Xu, Z.-D. and Qu, D.-M. (2001). Antitumour activity of transition metal complexes with the thiosemicarbazone derived from 3-acetylbulliferone. *Trans. Metal Chem.*, 26, 307-310.

Wang, X., Sarris, K., Kage, K., Zhang, D., Brown, S. P., Kolasa, T., Surowy, C., El Kouhen, O. F., Muchmore, S. W., Brioni, J. D. et al., (2009). Synthesis and evaluation of benzothiazole-based analogues as novel, potent, and selective fatty acid amide hydrolase inhibitors. *J. Med. Chem.*, 52, 170-180.

Wang, Z., Shi, X.-H., Wang, J., Zhou, T., Xu, Y.-Z., Huang, T.-T., Li, Y.-F., Zhao, Y.-L., Yang, L. and Yang, S.-Y. (2011). Synthesis, structure activity relationships and preliminary antitumor evaluation of benzothiazole-2-thiol derivatives as novel apoptosis inducers. *Bioorg. Med. Chem. Lett.*, 21, 1097-1101.

Wayne A (1997) National committee for clinical laboratory standards, NCCLS approved standard M27-A, USA.

Wojciechowski, M., Milewski, S., Mazerski, J. and Borowski, E. (2005). Glucosamine-6-phosphate synthase, a novel target for antifungal agents. Molecular modelling studies in drug design. *Acta Biochim Pol.*, 52(3), 647-653.

Wu, C., Wei, J., Gao, K. and Wang, Y. (2007). Dibenzothiazoles as novel amyloid-imaging agents. *Bioorg. Med. Chem.*, 15, 2789-2796.

Yadav, L. D. S. and Singh, S. (2001). Synthesis of antiviral acyclic C-nucleosides incorporating thiazolo-1, 3, 4-oxa (thia) diazole or thiazolo-1, 2, 4-triazole structure as a nucleobase. *Indian J. Chem.*, 40B, 440-442.

Yalcin, I., Oren, I., Sener, E., Akin, A. and Ucarturk, N. (1992). The synthesis and the structure-activity relationships of some substituted benzoxazoles, oxazolo (4,5-b) pyridines, benzothiazoles and benzimidazoles as antimicrobial agents. *Eur. J. Med. Chem.*, 27, 401-406.

Yang, Z., Sheng-Gang, Y., Yan-Ping, L., Ying, T., Ge-Fei, H., Qiong-You, W., Zhen, X. and Guang-Fu, Y. (2013). Design and synthesis of 1-(benzothiazol-5-yl)-1H-1, 2, 4-triazol-5-ones as protoporphyrinogen oxidase inhibitors. *Bioorg. Med. Chem.*, 21, 3245-3255.

Yar, M. S. and Ansari, Z. H. (2009). Synthesis and in vivo diuretic activity of biphenyl benzothiazole-2-carboxamide derivatives. *Acta Pol. Pharm.*, 66, 387-392.

Yarden, Y. and Sliwkowski, M. X. (2001). Untangling the ErbB signalling network. *Nat. Reviews Mol. Cell Bio.*, 2, 127-137.

Yarden, Y. and Ullrich, A. (1988). Growth factor receptor tyrosine kinases. *Ann. Rre. Biochem.*, 57, 443-478.

Yates, P. C., McCall, C. J. and Stevens, M. F. (1991). Structural studies on benzothiazoles. Crystal and molecular structure of 5, 6-dimethoxy-2-(4-methoxyphenyl)-benzothiazole and molecular orbital calculations on related compounds. *Tetrahedron*, 47, 6493-6502.

Yildiz-Oren, I., Yalcin, I., Aki-Sener, E. and Ucarturk, N. (2004). Synthesis and structure-activity relationships of new antimicrobial active multisubstituted benzazole derivatives. *Eur. J. Med. Chem.*, 39, 291-298.

Yin, Y., Lin, L., Ruiz, C., Cameron, M. D., Pocas, J., Grant, W., Schroter, T., Chen, W., Duckett, D., Schurer, S. et al., (2009). Benzothiazoles as Rho-associated kinase (ROCK-II) inhibitors. *Bioorg. Med. Chem. Lett.*, 19, 6686-6690.

Yoshida, T., Zhang, G. and Haura, E. B. (2010). Targeting epidermal growth factor receptor: central signaling kinase in lung cancer. *Biochem. Pharm.*, 80, 613-623.

Yuan, Y.-Q. and Guo, S.-R. (2011). TMSCl/Fe(NO<sub>3</sub>)<sub>3</sub>-Catalyzed synthesis of 2-arylbenzothiazoles and 2-arylbenzimidazoles under ultrasonic irradiation. *Syn. Comm.*, 41, 2169-2177.

Zheng, C. H., Zhou, Y. J., Zhu, J., Ji, H. T., Chen, J., Li, Y. W., Sheng, C. Q., Lu, J. G., Jiang, J. H., Tang, H. et al. (2007). Construction of a three-dimensional pharmacophore for Bcl-2 inhibitors by flexible docking and the multiple copy simultaneous search method. *Bioorg. Med. Chem.*, 15, 6407-6417.

Zou Q, Liu Y, Zhengwei Y, et al. Preparation of heterocycles as protein kinase inhibitors. WO127345, 2013.

Zwick, E., Bange, J. and Ullrich, A. (2001). Receptor tyrosine kinase signalling as a target for cancer intervention strategies. *Endocr. Relat. Cancer*, 8(3), 161-173.