

### List of Figures

**Figure 1.1:** The Lycurgus Cup, appearing green in (A) reflected light and red in (B) transmitted light – Courtesy of the British Museum.

**Figure 1.2:** Schematic illustration of surface plasmon resonance by interaction of light with the surface dipole of nanomaterials.

**Figure 1.3:** Top-down Vs Bottom-up.

**Figure 1.4:** Faraday is credited as the discoverer of metallic colloid, he prepared gold colloids that were purple red in color by the reduction of gold salts with phosphorous, courtesy – Royal Institution of Great Britain available from: <http://www.rigb.org/rimain/heritage/faradaypage.jsp>.

**Figure 1.5:** Formation of AuNPs coated with citrate and its oxidation product by reaction of sodium citrate with chloro auric acid.

**Figure 1.6:** General scheme for borohydride Brust-Schiffrin method for gold nanomaterials functionalized with thiols. In figure, where m and n represent the stoichiometric ratio.

**Figure 1.7:** Ascorbic acid mediated synthesis of gold nanoparticles.

**Figure 1.8:** Proposed reaction scheme illustrating luminol- reduced gold nanoparticles.

**Figure 1.9:** Reaction scheme for the TTF catalyzed formation of silver dendritic nanostructures.

**Figure 1.10:** Time-dependent UV-vis absorption spectra of solutions containing  $\text{HAuCl}_4$  and medium molecular weight chitosan and corresponding TEM image of the gold nanoparticles.

**Figure 1.11:** Photochemical scheme illustrating the step wise reduction of Au (III).

**Figure 1.12:** Proposed scheme illustrating the photochemical reduction of Ag (I) and the growth of polyhedral silver nanostructures.

**Figure 1.13:** Diversified applications of nanomaterials ranging from consumer applications → healthcare → electronics → commercial power devices products.

**Figure 1.14:** Thermodynamically favored bulk agglomerated particle in the absence of any stabilizer (route a). stepwise formation of metallic nanomaterials and at the same time Electrostatic/ vander-waal interaction of metal nano particles with capping agent provide the stabilization of metal nanostructures (route b).

**Figure 1.15:** Steric stabilization of nanostructured metal colloids capped with protecting agent.

**Figure 1.16:** Mechanism of electrocatalysis on chemically modified electrode.

**Figure 1.17:** Schematic outline illustrating the objectives of thesis.

**Figure 2.1:** Electronic transition in a molecule.

**Figure 2.2:** Block diagram of UV visible spectroscopy.

**Figure 2.3:** Schematic ray diagram illustrating Beer-Lambert law.

**Figure 2.4:** Block diagram for FT-IR.

**Figure 2.5:** Scheme illustrating the transfer of electron density from non-metal to metal atom, resulting into the change in the characteristics vibrational frequency.

**Figure 2.6:** Physical process involved in XPS.

**Figure 2.7:** Block diagram of modern electron spectroscopy for chemical analysis.

**Figure 2.8:** Block diagram of scanning electron microscopy for surface analysis.

**Figure 2.9:** Block diagram of transmission electron microscopy for surface imaging.

- Figure 2.10:** Ray diagram of SAED, illustrating electron diffraction.
- Figure 2.11:** Common excitation signal in voltammetry.
- Figure 2.12:** Electrochemical set up for voltammetric measurement.
- Figure 3.1:** Photochemically generated electron utilizing the reduction of noble metal ions and possible oxidation product of dithizone.
- Figure 3.2:** Reaction of Au (III) with dithizone.
- Figure 3.3:** Time dependent UV-visible spectrum of the photochemical assisted formation of gold nanoparticles.
- Figure 3.4:** Reaction scheme for the photochemical step wise reduction of Au (III) to Au (0).
- Figure 3.5:** (a) FT-IR of DTZ (b) FT-IR of Au-DTZH
- Figure 3.6:** (a) XPS survey spectrum of Au-DTZH in full range (b) X-ray photoelectron spectra of Au (4f).
- Figure 3.7:** (a) TEM studies for the formation of nano gold w.r.t. time of light exposure a=09 min, b=17 min, c=28 min, d=54 min.
- Figure 3.8:** (a) TEM image of gold nanoparticles at the completion of reaction and the respective (b) EDX and (c) SAED pattern.
- Figure 3.9:** FESEM and EDAX image of chemically modified electrode Au-DTZH platform.
- Figure 3.10:** CV Response for (a) Bare electrode (b) CV of  $\text{Fe}(\text{CN})_6^{4-/3-}$  redox couple on without modified electrode (c) CV of  $\text{Fe}(\text{CN})_6^{4-/3-}$  redox couple on Au-DTZH modified electrode.
- Figure 3.11:** Cyclic voltammogram response of successive addition of thiocyanate in 0.1 M tris buffer at pH 7.2. Asterisk marked curve: characteristics CV of gold nano crystals.

**Figure 3.12:** Schematic representation of the nano integrated platform of the Au-DTZH as a modified electrode for electro-sensing of thiocyanate.

**Figure 3.13:** Amperometric response at an applied potential of 0.547 V vs. Ag/AgCl for successive addition of thiocyanate in 0.1 M tris buffer at pH 7.2 under stirred condition. *Inset:* corresponding standard addition calibration plot of chronoamperometric current vs. thiocyanate concentration.

**Figure 4.1:** (a) Time dependent UV-vis spectrum of the formation of nanosilver, a-only dithizone, b-at 6 min, c-at 16min, d-30 min, e-55min, (b) UV-vis spectrum of the nanosilver at the completion of reaction (after 2 hrs).

**Figure 4.2:** (a) Reaction scheme for the photochemical assisted formation of nano silver by dithizone where photochemically generated electrons can easily captured by  $\text{Ag}^+$  ions to generate silver nanostructures via photo-reduction. (b) This figure schematically illustrates that how nucleation and growth phenomenon promotes the silver nanoparticles formation within DTZH matrix.

**Figure 4.3:** (a) TEM studies for the formation of nano silver w.r.t. time of light exposure a=6 min, b=16 min, c=30 min, d=60 min.

**Figure 4.4:** (a) TEM image of nanosilver at the completion (after 2 hrs.) of reaction (b) corresponding size distribution of nanoparticles.

**Figure 4.5:** SAED and EDAX spectrum of the corresponding TEM images of silver nanoparticles.

**Figure 4.6:** (a) FT-IR of DTZ (b) FT-IR of Ag-DTZH

**Figure 4.7:** (a) XPS survey spectrum of Ag-DTZH in full range (b) X-ray photoelectron spectra of Ag (3d).

**Figure 4.8:** FESEM and EDAX image of modified electrode Ag-DTZH platform.

**Figure 4.9:** (a) CV of bare unmodified electrode (b) modified electrode with silver nano particles having characteristics CV of AgNPs.

**Figure 4.10:** Cyclic voltammogram response of successive addition of cefotaxime

**Figure 4.11:** Schematic representation of the nano integrated platform of the Ag-DTZH as a modified electrode for electro-sensing of cefotaxime.

**Figure 4.12:** (a) Amperometric response at an applied potential of 0.789 V vs. Ag/AgCl for successive addition of cefotaxime in 0.1 M tris buffer at pH 7.2 under stirred condition. (b) Corresponding standard addition calibration plot of chronoamperometric current vs. cefotaxime concentration.

**Figure 5.1:** Photochemically generated electron utilizing the reduction of noble metal ions and possible oxidation product of phenothiazine.

**Figure 5.2:** UV-visible spectrum of phenothiazine.

**Figure 5.3:** UV-Vis spectrum gold nanoparticles, *Inset:* zoom view of SPR band of gold nanoparticles.

**Figure 5.4:** Time dependent UV-visible spectrum of gold nanoparticles formation, a: 0 min, b: 2 min, c: 5 min, d: 12 min, e: 22, min, f: 32 min, g: 50 min.

**Figure 5.5:** Plausible reaction scheme for the photochemical formation of gold nanoparticles and their stabilization by phenothiazine and its oxidation product.

**Figure 5.6:** (a) Cyclic voltammogram of unmodified ITO electrode (b) cyclic voltammogram of gold nanomaterials on ITO electrode surface in 0.5 M H<sub>2</sub>SO<sub>4</sub>.

**Figure 5.7:** (a) TEM studies for the formation of nano gold w.r.t. light exposure, recorded at different time interval a: 0 min, b: 5 min, c: 22 min, d: 32 min.

**Figure 5.8:** (a) TEM image (b) SAED pattern of Au nanoparticles. (c) size distribution of prepared nanomaterials.

**Figure 5.9:** FT-IR Spectrum of PTZ and Au-PTZH.

**Figure 5.10:** (a) XPS survey spectrum of Au-PTZH in full range (b) X-ray photoelectron spectra of Au (4f).

**Figure 5.11:** FESEM and EDAX image of the Au-PTZH nanostructured platform (gold nanomaterials modified electrode).

**Figure 5.12:** Cyclic voltammograms response for  $\text{Fe}(\text{CN})_6^{4-/3-}$  redox couple.

**Figure 5.13:** Cyclic voltammogram response of successive addition of phosphate in 0.1 M tris buffer at pH 7.2.

**Figure 5.14:** Schematic representation of the nano integrated platform of the Au-PTZH as a modified electrode.

**Figure 5.15:** Amperometric response at an applied potential of -0.336 volts Vs Ag/AgCl for successive addition of phosphate in 0.1 M tris buffer at pH 7.2 under stirred condition.

**Figure 5.16:** Corresponding standard addition calibration plot of chronoamperometric current Vs phosphate concentration.

**Figure 6.1:** Reaction scheme for the photochemical assisted formation of nano silver by phenothiazine.

**Figure 6.2:** UV-vis spectrum of nanosilver, Inset to the figure show zoom view of band of nano silver.

**Figure 6.3:** Bare unmodified electrode (a) and Characteristic CV of nano silver in Tris HCl buffer (b)

**Figure 6.4:** (a) FT-IR of PTZ (b) FT-IR of Ag-PTZH

**Figure 6.5:** (a) XPS survey spectrum of Ag-PTZH in full range (b) X-ray photoelectron spectra of Ag (3d).

**Figure 6.6:** (a) TEM image (b) SAED pattern (c) HRTEM images showing lattice fringes spacing of silver nanodendrites.

**Figure 6.7:** TEM images of Ag-PTZH when the reaction is carried out for longer time.

**Figure 6.8:** Scheme illustrating for formation of silver nanodendrites (a-g).

**Figure 6.9:** FESEM and EDAX image of the modified electrode surface with silver nano dendrites.

**Figure 6.10:** CV Response for (a) Bare electrode (b) CV of  $\text{Fe}(\text{CN})_6^{4-/3-}$  redox couple on without modified electrode (c) CV of  $\text{Fe}(\text{CN})_6^{4-/3-}$  redox couple on Ag-PTZH modified electrode.

**Figure 6.11:** Cyclic voltammogram response of successive addition of nitrite in 0.1 M tris buffer at pH 7.2.

**Figure 6.12:** Schematic representation of the nano integrated platform of the Ag-PTZH as a modified electrode for electro-sensing of nitrite.

**Figure 6.13:** Amperometric response at an applied potential of 0.44 volts Vs. Ag/AgCl for successive addition of nitrite in 0.1 M tris buffer at pH 7.2 under stirred condition.

**Figure 6.14:** Corresponding standard addition calibration plot of chronoamperometric current Vs nitrite concentration.