

## LIST OF FIGURES

---

---

<b>Figure No.</b>	<b>Figure Caption</b>	<b>Page No.</b>
<b>Figure 1.1</b>	Distribution of World's Water	<b>3</b>
<b>Figure 1.2</b>	The Water Cycle	<b>4</b>
<b>Figure 1.3</b>	Major sources of water pollution	<b>8</b>
<b>Figure 1.4</b>	Top down and bottom up approaches for the synthesis of nanomaterials	<b>31</b>
<b>Figure 3.1</b>	Speciation diagram of Chromium	<b>53</b>
<b>Figure 3.2</b>	Speciation diagram of Nickel	<b>56</b>
<b>Figure 4.1</b>	X-ray diffraction (XRD) pattern of nano-alumina	<b>76</b>
<b>Figure 4.2</b>	Fourier transform infra-red (FTIR) spectrum of nano-alumina	<b>77</b>
<b>Figure 4.3</b>	(A) Transmission electron micrograph; and (B) Selected area electron diffraction (SAED) pattern of nano-alumina	<b>78</b>
<b>Figure 4.4</b>	Scanning electron micrograph (SEM) of alumina nanoparticles	<b>79</b>
<b>Figure 4.5</b>	Energy dispersive x-ray (EDAX) pattern of alumina nanoparticles	<b>79</b>
<b>Figure 4.6</b>	X-ray diffraction pattern (XRD) pattern of nano-cupric oxide	<b>83</b>
<b>Figure 4.7</b>	Fourier transform infra-red (FTIR) spectrum of nano-cupric oxide	<b>83</b>
<b>Figure 4.8</b>	Transmission electron micrograph (TEM) of nano-cupric oxide	<b>84</b>
<b>Figure 4.9</b>	Scanning electron micrograph (SEM) of nano-cupric oxide	<b>85</b>
<b>Figure 4.10</b>	Energy dispersive x-ray (EDAX) pattern of nano-cupric	<b>85</b>

oxide

<b>Figure 4.11</b>	pHzpc of synthesized nano-alumina	<b>86</b>
<b>Figure 4.12</b>	pHzpc of synthesized nano-cupric oxide	<b>87</b>
<b>Figure 5.1</b>	XRD pattern of nano-alumina before and after adsorption of Cr(VI) ions	<b>91</b>
<b>Figure 5.2</b>	FT-IR spectra of nano-alumina before and after adsorption of Cr(VI) ions	<b>91</b>
<b>Figure 5.3</b>	EDS pattern of nano-alumina after adsorption of Cr(VI) ions	<b>92</b>
<b>Figure 5.4</b>	Effect of initial pH on removal (%) of chromium from aqueous solution on nano-cupric oxide (Initial concentration= 10 mg/L, adsorbent dose= 10g/L, Temperature= 303 K)	<b>93</b>
<b>Figure 5.5</b>	Effect of contact time on removal (%) of chromium from aqueous solution on nano-cupric oxide (Initial concentration=10mg/L, Initial pH=2.0, Initial dose=10g/L, Temperature= 303 K)	<b>94</b>
<b>Figure 5.6</b>	Effect of adsorbent dose on removal (%) of chromium from aqueous solution on nano-cupric oxide (Initial pH=2.0, Initial concentration=10mg/L, Temperature=303 K)	<b>94</b>
<b>Figure 5.7</b>	Effect of initial concentration on removal (%) of chromium from aqueous solution on nano-cupric oxide (Initial pH= 2.0, Initial dose =10g/L, Temperature=303 K)	<b>95</b>
<b>Figure 5.8</b>	(a) Surface plot; and (b) Contour plot of percentage removal vs. dose and concentration at hold value of pH at 10	<b>101</b>
<b>Figure 5.9</b>	(a) Surface plot; and (b) Contour plot of percentage removal vs. pH and dose at hold value of concentration at 25 mg/L	<b>102</b>
<b>Figure 5.10</b>	(a) Surface plot; and (b) Contour plot of percentage removal vs. pH and concentration at hold value of dose at 20 g/L	<b>103</b>
<b>Figure 5.11</b>	Response optimization plot for chromium removal on nano-alumina	<b>104</b>

<b>Figure 5.12</b>	Linear Langmuir isotherm plot of Cr(VI) removal on nano-alumina (symbols represent the experimental data and straight lines represent the data estimated by the model)	<b>107</b>
<b>Figure 5.13</b>	Linear Freundlich isotherm plot of Cr(VI) removal on nano- alumina (symbols represent the experimental data and straight lines represent the data estimated by the model)	<b>107</b>
<b>Figure 5.14</b>	Non-linear Langmuir isotherm plot of chromium removal on nano-alumina obtained by in-built Microcal origin function symbols (represent the experimental data and lines represent the data estimated by the model)	<b>109</b>
<b>Figure 5.15</b>	Non-linear Freundlich isotherm plot of chromium removal on nano-alumina obtained by in-built Microcal origin function (symbols represent the experimental data and lines represent the data estimated by the model)	<b>110</b>
<b>Figure 5.16</b>	Non-linear Langmuir isotherm plot of chromium removal on nano-alumina obtained by customized Microcal origin function (symbols represent the experimental data and lines data represent the estimated by the model)	<b>110</b>
<b>Figure 5.17</b>	Non-linear Freundlich isotherm plot of chromium removal on nano-alumina obtained by customized Microcal origin function (symbols represent the experimental data and lines represent the data estimated by the model)	<b>111</b>
<b>Figure 5.18</b>	Linear pseudo-first order plot of chromium removal on nano-alumina (symbols represent the experimental data and straight lines represent the data estimated by the model)	<b>114</b>
<b>Figure 5.19</b>	Linear pseudo-second order plot of chromium removal on nano-alumina (symbols represent the experimental data and straight lines represent the data estimated by the model)	<b>115</b>
<b>Figure 5.20</b>	Non-linear pseudo-first order plot of chromium removal on nano-alumina obtained by in-built Microcal origin function (symbols represent the experimental data and lines represent the data estimated by the model)	<b>117</b>
<b>Figure 5.21</b>	Non-linear pseudo-second order plot of chromium removal on nano-alumina obtained by in-built Microcal origin	<b>118</b>

	function (symbols represent the experimental data and lines represent the data estimated by the model)	
<b>Figure 5.22</b>	Non-linear pseudo-first order plot of chromium removal on nano-alumina obtained by customized Microcal origin function (symbols represent the experimental data and lines represent the data estimated by the model)	<b>118</b>
<b>Figure 5.23</b>	Non-linear pseudo-second order plot of chromium removal on nano-alumina obtained by in-built Microcal origin function (symbols represent the experimental data and straight lines represent the data estimated by the model)	<b>119</b>
<b>Figure 5.24</b>	A) Intra-particle diffusion plot for removal of chromium from aqueous solution on nano-alumina. B) Boyd model plot for removal of chromium from aqueous solution on nano-alumina	<b>122</b>
<b>Figure 5.25</b>	Effect of temperature on removal of Cr(VI) from aqueous solutions on nano-alumina	<b>124</b>
<b>Figure 5.26</b>	Arrhenius plot for removal of chromium from aqueous solution on nano-alumina	<b>127</b>
<b>Figure 6.1</b>	XRD pattern of nano-alumina before and after adsorption of Ni(II) ions	<b>132</b>
<b>Figure 6.2</b>	FT-IR spectra of nano-alumina before and after adsorption of Ni(II) ions	<b>132</b>
<b>Figure 6.3</b>	EDS pattern of nano-alumina after adsorption of Ni(II) ions	<b>133</b>
<b>Figure 6.4</b>	Effect of initial pH on removal (%) of nickel from aqueous solution on nano-alumina (Initial concentration= 20 mg/L, adsorbent dose= 10g/L, Temperature= 303 K)	<b>135</b>
<b>Figure 6.5</b>	Effect of adsorbent dose on removal (%) of nickel from aqueous solution on nano-alumina (Initial pH=7.0, Initial concentration=20mg/L, Temperature=303 K)	<b>135</b>
<b>Figure 6.6</b>	Effect of contact time on removal (%) of nickel from aqueous solution on nano-alumina (Initial concentration=10mg/L, Initial pH=7.0, Initial dose=10g/L, Temperature= 303 K)	<b>136</b>
<b>Figure 6.7</b>	Effect of initial concentration on removal (%) of nickel from aqueous solution on nano-alumina (Initial pH= 7.0,	<b>136</b>

Initial dose=10g/L, Temperature=303 K)

<b>Figure 6.8</b>	(a) Surface plot; and (b) Contour plot of percentage removal vs. dose and concentration at hold value of pH at	<b>142</b>
<b>Figure 6.9</b>	(a) Surface plot; and (b) Contour plot of percentage removal vs. pH and dose at hold value of concentration at 25 mg/L	<b>143</b>
<b>Figure 6.10</b>	(a) Surface plot; and (b) Contour plot of percentage removal vs. pH and concentration at hold value of dose at 20 g/L	<b>145</b>
<b>Figure 6.11</b>	Response optimization plot for nickel removal on nano-alumina	<b>146</b>
<b>Figure 6.12</b>	Linear Langmuir isotherm plot of nickel removal on nano-alumina (symbols represent the experimental data and straight lines represent the data estimated by the model)	<b>149</b>
<b>Figure 6.13</b>	Linear Freundlich isotherm plot of nickel removal on nano-alumina (symbols represent the experimental data and straight lines represent the data estimated by the model)	<b>149</b>
<b>Figure 6.14</b>	Non-linear Langmuir isotherm plot of nickel removal on nano-alumina obtained by in-built Microcal origin function (symbols represent the experimental data and lines represent the data estimated by the model)	<b>151</b>
<b>Figure 6.15</b>	Non-linear Freundlich isotherm plot of nickel removal on nano-alumina obtained by customized Microcal origin function (symbols represent the experimental data and lines represent the data estimated by the model)	<b>152</b>
<b>Figure 6.16</b>	Non-linear Langmuir isotherm plot of nickel removal on nano-alumina obtained by customized Microcal origin function (symbols represent the experimental data and lines represent the data estimated by the model)	<b>152</b>
<b>Figure 6.17</b>	Non-linear Freundlich isotherm plot of nickel removal on nano-alumina obtained by customized Microcal origin function (symbols represent the experimental data and lines represent the data estimated by the model)	<b>153</b>
<b>Figure 6.18</b>	Linear pseudo-first order plot of nickel removal on nano-alumina (symbols represent the experimental data and straight lines represent the data estimated by the model)	<b>156</b>

<b>Figure 6.19</b>	Linear pseudo-second order plot of nickel removal on nano-alumina (symbols represent the experimental data and straight lines represent the data estimated by the model)	<b>157</b>
<b>Figure 6.20</b>	Non-linear pseudo-first order plot of nickel removal on nano-alumina obtained by in-built Microcal origin function (symbols represent the experimental data and lines represent the data estimated by the model)	<b>159</b>
<b>Figure 6.21</b>	Non-linear pseudo-second order plot of nickel removal on nano-alumina obtained by in-built Microcal origin function (symbols represent the experimental data and lines represent the data estimated by the model)	<b>160</b>
<b>Figure 6.22</b>	Non-linear pseudo-first order plot of nickel removal on nano-alumina obtained by customized Microcal origin function (symbols represent the experimental data and lines represent the data estimated by the model)	<b>160</b>
<b>Figure 6.23</b>	Non-linear pseudo-second order plot of nickel removal on nano-alumina obtained by in-built Microcal origin function (symbols represent the experimental data and lines represent the data estimated by the model)	<b>161</b>
<b>Figure 6.24</b>	A) Intra-particle diffusion plot for removal of nickel from aqueous data solution on nano-alumina. B) Boyd model plot for removal of nickel from aqueous solution on nano-alumina	<b>162</b>
<b>Figure 6.25</b>	Effect of temperature on removal of nickel from aqueous solutions on nano-alumina	<b>165</b>
<b>Figure 6.26</b>	Arrhenius plot for removal of nickel from aqueous solution on nano-alumina	<b>168</b>
<b>Figure 7.1</b>	XRD pattern of nano-cupric oxide before and after adsorption of Cr(VI) ions	<b>173</b>
<b>Figure 7.2</b>	FT-IR spectra of nano-cupric oxide before and after adsorption of Cr(VI) ions	<b>173</b>
<b>Figure 7.3</b>	EDS pattern of nano- nano-cupric oxide after adsorption of Cr(VI) ions	<b>174</b>
<b>Figure 7.4</b>	Effect of initial pH on removal (%) of chromium from	<b>175</b>

aqueous solution on nano-cupric oxide (Initial concentration=10 mg/L, adsorbent dose= 8g/L, Temperature= 303 K)

<b>Figure 7.5</b>	Effect of adsorbent dose on removal (%) of chromium from aqueous solution on nano-cupric oxide (Initial pH=2.0, Initial concentration=10mg/L, Temperature=303 K)	<b>176</b>
<b>Figure 7.6</b>	Effect of contact time on removal (%) of chromium from aqueous solution on nano-cupric oxide (Initial concentration=10mg/L, Initial pH=2.0, Initial dose=8g/L, Temperature= 303 K)	<b>176</b>
<b>Figure 7.7</b>	Effect of initial concentration on removal (%) of chromium from aqueous solution on nano-cupric oxide (Initial pH= 2.0, Initial dose=8g/L, Temperature=303 K)	<b>177</b>
<b>Figure 7.8</b>	(a) Surface plot; and (b) Contour plot of percentage removal vs. dose and concentration at hold value of pH at 10	<b>182</b>
<b>Figure 7.9</b>	(a) Surface plot; and (b) Contour plot of percentage removal vs. pH and dose at hold value of concentration at 25 mg/L	<b>184</b>
<b>Figure 7.10</b>	(a) Surface plot; and (b) Contour plot of percentage removal vs. pH and concentration at hold value of dose at 20 g/L	<b>185</b>
<b>Figure 7.11</b>	Response optimization plot for chromium removal on nano-cupric oxide	<b>186</b>
<b>Figure 7.12</b>	Linear Langmuir isotherm plot of Cr(VI) removal on nano-cupric oxide (symbols represent the experimental data and straight lines represent the data estimated by the model)	<b>189</b>
<b>Figure 7.13</b>	Linear Freundlich isotherm plot of Cr(VI) removal on nano-cupric oxide (symbols represent the experimental data and straight lines represent the data estimated by the model)	<b>190</b>
<b>Figure 7.14</b>	Non-linear Langmuir isotherm plot of chromium removal on nano-cupric oxide obtained by in-built Microcal origin function (symbols represent the experimental data and lines represent the data estimated by the model)	<b>192</b>

<b>Figure 7.15</b>	Non-linear Freundlich isotherm plot of chromium removal on nano-cupric oxide obtained by in-built Microcal origin function (symbols represent the experimental data and lines represent the data estimated by the model)	<b>193</b>
<b>Figure 7.16</b>	Non-linear Langmuir isotherm plot of chromium removal on nano-cupric oxide obtained by customized Microcal origin function (symbols represent the experimental data and lines represent the data estimated by the model)	<b>193</b>
<b>Figure 7.17</b>	Non-linear Freundlich isotherm plot of chromium removal on nano-cupric oxide obtained by customized Microcal origin function (symbols represent the experimental data and lines represent the data estimated by the model)	<b>194</b>
<b>Figure 7.18</b>	Linear pseudo-first order plot of chromium removal on nano-cupric oxide (symbols represent the experimental data and straight lines represent the data estimated by the model)	<b>197</b>
<b>Figure 7.19</b>	Linear pseudo-second order plot of chromium removal on nano-cupric oxide (symbols represent the experimental data and straight lines represent the data estimated by the model)	<b>198</b>
<b>Figure 7.20</b>	Non-linear pseudo-first order plot of chromium removal on nano-cupric oxide obtained by in-built Microcal origin function (symbols represent the experimental data and lines represent the data estimated by the model)	<b>200</b>
<b>Figure 7.21</b>	Non-linear pseudo-second order plot of chromium removal on nano-cupric oxide obtained by in-built Microcal origin function (symbols represent the experimental data and lines represent the data estimated by the model)	<b>200</b>
<b>Figure 7.22</b>	Non-linear pseudo-first order plot of chromium removal on nano-cupric oxide obtained by customized Microcal origin function represent the experimental data and lines represent the data estimated by the model)	<b>201</b>
<b>Figure 7.23</b>	Non-linear pseudo-second order plot of chromium removal on nano-cupric oxide obtained by in-built Microcal origin function (symbols represent the experimental data and lines represent the data estimated by the model)	<b>201</b>
<b>Figure 7.24</b>	A) Intra-particle diffusion plot for removal of chromium from aqueous solution on nano- cupric oxide. B) Boyd	<b>203</b>



	model plot for removal of chromium from aqueous solution on nano-cupric oxide	
<b>Figure 7.25</b>	Effect of temperature on removal of chromium from aqueous solutions on nano-cupric oxide	<b>206</b>
<b>Figure 7.26</b>	Arrhenius plot for removal of chromium from aqueous solution on nano-cupric oxide	<b>209</b>
<b>Figure 8.1</b>	XRD pattern of nano-cupric oxide before and after adsorption of Ni(II) ions	<b>215</b>
<b>Figure 8.2</b>	FT-IR spectra of nano-cupric oxide before and after adsorption of Ni(II) ions	<b>215</b>
<b>Figure 8.3</b>	EDS pattern of nano-cupric oxide after adsorption of Ni(II) ions	<b>216</b>
<b>Figure 8.4</b>	Effect of initial pH on removal (%) of nickel from aqueous solution on nano-cupric oxide (Initial concentration= 20 mg/L, adsorbent dose= 6g/L, Temperature= 303K)	<b>217</b>
<b>Figure 8.5</b>	Effect of adsorbent dose on removal (%) of nickel from aqueous solution on nano-cupric oxide (Initial pH=7.0, Initial concentration=20 mg/L, Temperature=303K)	<b>218</b>
<b>Figure 8.6</b>	Effect of contact time on removal (%) of nickel from aqueous solution on nano-cupric oxide (Initial concentration=20mg/L, Initial pH=7.0, Initial dose=8g/L, Temperature= 303K)	<b>219</b>
<b>Figure 8.7</b>	Effect of initial concentration on removal (%) of nickel from aqueous solution on nano-cupric oxide (Initial pH= 7.0, Initial dose=6g/L, Temperature=303 K)	<b>219</b>
<b>Figure 8.8</b>	(a) Surface plot; and (b) Contour plot of percentage removal vs. dose and concentration at hold value of pH at 7.0	<b>225</b>
<b>Figure 8.9</b>	(a) Surface plot; and (b) Contour plot of percentage removal vs. pH and dose at hold value of concentration at 25 mg/L	<b>226</b>
<b>Figure 8.10</b>	(a) Surface plot; and (b) Contour plot of percentage removal vs. pH and concentration at hold value of dose at 20 g/L	<b>227</b>

<b>Figure 8.11</b>	Response optimization plot for nickel removal on nano-cupric oxide	<b>228</b>
<b>Figure 8.12</b>	Linear Langmuir isotherm plot of nickel removal on nano-cupric oxide (symbols represent the experimental data and straight lines represent the data estimated by the model)	<b>231</b>
<b>Figure 8.13</b>	Linear Freundlich isotherm plot of nickel removal on nano-cupric oxide (symbols represent the experimental data and straight lines represent the data estimated by the model)	<b>232</b>
<b>Figure 8.14</b>	Non-linear Langmuir isotherm plot of nickel removal on nano-cupric oxide obtained by in-built Microcal origin function (symbols represent the experimental data and lines represent the data estimated by the model)	<b>233</b>
<b>Figure 8.15</b>	Non-linear Freundlich isotherm plot of nickel removal on nano-cupric oxide obtained by in-built Microcal origin function (symbols represent the experimental data and lines represent the data estimated by the model)	<b>234</b>
<b>Figure 8.16</b>	Non-linear Langmuir isotherm plot of nickel removal on nano-cupric oxide obtained by customized Microcal origin function (symbols represent the experimental data and lines represent the data estimated by the model)	<b>234</b>
<b>Figure 8.17</b>	Non-linear Freundlich isotherm plot of nickel removal on nano-cupric oxide obtained by customized Microcal origin function (symbols represent the experimental data and lines represent the data estimated by the model)	<b>235</b>
<b>Figure 8.18</b>	Linear pseudo-first order plot of nickel removal on nano-cupric oxide (symbols represent the experimental data and straight lines represent the data estimated by the model)	<b>238</b>
<b>Figure 8.19</b>	Linear pseudo-second order plot of nickel removal on nano-cupric oxide (symbols represent the experimental data and lines represent the data estimated by the model)	<b>239</b>
<b>Figure 8.20</b>	Non-linear pseudo-first order plot of nickel removal on nano-cupric oxide obtained by in-built Microcal origin function (symbols represent the experimental data and lines represent the data estimated by the model)	<b>241</b>
<b>Figure 8.21</b>	Non-linear pseudo-second order plot of nickel removal on nano-cupric oxide obtained by in-built Microcal origin	<b>242</b>

function (symbols represent the experimental data and lines represent the data estimated by the model)

- Figure 8.22** Non-linear pseudo-first order plot of nickel removal on nano-cupric oxide obtained by customized Microcal origin function (symbols represent the experimental data and lines represent the data estimated by the model) **242**
- Figure 8.23** Non-linear pseudo-second order plot of nickel removal on nano-cupric oxide obtained by in-built Microcal origin function (symbol represent the experimental data and lines represent the data estimated by the model) **243**
- Figure 8.24** A) Intra-particle diffusion plot for removal of nickel from aqueous solution on nano-cupric oxide. B) Boyd model plot for removal of nickel from aqueous solution on nano-cupric oxide **245**
- Figure 8.25** Effect of temperature on removal of nickel from aqueous solutions on nano-cupric oxide **247**
- Figure 8.26** Arrhenius plot for removal of nickel from aqueous solution on nano-cupric oxide **251**