List of Figures

Figure No.	Title	Page No.
Figure 2.1	Percentage contributions of different mining activities to total dust emissions from a typical Indian opencast coal mine	7
Figure 2.2	Residence times for homogeneously distributed dust particles of different aerodynamic diameters	11
Figure 2.3	Three regions of respiratory system and their parts	12
Figure 2.4	A comparison of the actual dust dispersion within a mine with Gaussian plume and CFD models	31
Figure 3.1	Satellite image of Mine 'A' Opencast Coal Project	50
Figure 3.2	Receptors location in the study area of Mine 'A'	55
Figure 3.3	Wind - rose diagram of Jharsuguda District for the month of January 2013	58
Figure 3.4	Wind - rose diagram of Jharsuguda District for the month of May 2013	59
Figure 3.5	Digital Elevation Model of Mine 'A' generated by AERMAP	61
Figure 3.6	1st Highest predicted concentration level of PM_{10} for daily averaging time period in January 2013	63
Figure 3.7	2nd Highest predicted concentration level of PM_{10} for daily averaging time period in January 2013	63
Figure 3.8	3rd Highest predicted concentration level of PM_{10} for daily averaging time period in January 2013	64
Figure 3.9	4th Highest predicted concentration level of PM_{10} for daily averaging time period in January 2013	64
Figure 3.10	5th Highest predicted concentration level of PM_{10} for daily averaging time period in January 2013	65
Figure 3.11	6th Highest predicted concentration level of PM_{10} for daily averaging time period in January 2013	65
Figure 3.12	7th Highest predicted concentration level of PM_{10} for daily averaging time period in January 2013	66
Figure 3.13	8th Highest predicted concentration level of PM_{10} for daily averaging time period in January 2013	66
Figure 3.14	9th Highest predicted concentration level of PM_{10} for daily averaging time period in January 2013	67
Figure 3.15	10th Highest predicted concentration level of PM_{10} for daily averaging time period in January 2013	67
Figure 3.16	11th Highest predicted concentration level of PM_{10} for daily averaging time period in January 2013	68

Figure 3.17	12th Highest predicted concentration level of PM_{10} for daily averaging time period in January 2013	68
Figure 3.18	13th Highest predicted concentration level of PM_{10} for daily averaging time period in January 2013	69
Figure 3.19	14th Highest predicted concentration level of PM_{10} for daily averaging time period in January 2013	69
Figure 3.20	15th Highest predicted concentration level of PM_{10} for daily averaging time period in January 2013	70
Figure 3.21	16th Highest predicted concentration level of PM_{10} for daily averaging time period in January 2013	70
Figure 3.22	17th Highest predicted concentration level of PM_{10} for daily averaging time period in January 2013	71
Figure 3.23	18th Highest predicted concentration level of PM_{10} for daily averaging time period in January 2013	71
Figure 3.24	19th Highest predicted concentration level of PM_{10} for daily averaging time period in January 2013	72
Figure 3.25	20th Highest predicted concentration level of PM_{10} for daily averaging time period in January 2013	72
Figure 3.26	21st Highest predicted concentration level of PM_{10} for daily averaging time period in January 2013	73
Figure 3.27	22nd Highest predicted concentration level of PM_{10} for daily averaging time period in January 2013	73
Figure 3.28	23rd Highest predicted concentration level of PM_{10} for daily averaging time period in January 2013	74
Figure 3.29	24th Highest predicted concentration level of PM_{10} for daily averaging time period in January 2013	74
Figure 3.30	25th Highest predicted concentration level of PM_{10} for daily averaging time period in January 2013	75
Figure 3.31	26th Highest predicted concentration level of PM_{10} for daily averaging time period in January 2013	75
Figure 3.32	27th Highest predicted concentration level of PM_{10} for daily averaging time period in January 2013	76
Figure 3.33	1st Highest predicted concentration level of PM_{10} for daily averaging time period in May 2013	76
Figure 3.34	2nd Highest predicted concentration level of PM_{10} for daily averaging time period in May 2013	77
Figure 3.35	3rd Highest predicted concentration level of PM_{10} for daily averaging time period in May 2013	77
Figure 3.36	4th Highest predicted concentration level of PM_{10} for daily averaging time period in May 2013	78
Figure 3.37	5th Highest predicted concentration level of PM_{10} for daily averaging time period in May 2013	78
Figure 3.38	6th Highest predicted concentration level of PM_{10} for daily averaging time period in May 2013	79
Figure 3.39	7th Highest predicted concentration level of PM_{10} for daily averaging time period in May 2013	79

Figure 3.40	8th Highest predicted concentration level of PM_{10} for daily averaging time period in May 2013	80
Figure 3.41	9th Highest predicted concentration level of PM_{10} for daily averaging time period in May 2013	80
Figure 3.42	10th Highest predicted concentration level of PM_{10} for daily averaging time period in May 2013	81
Figure 3.43	11th Highest predicted concentration level of PM_{10} for daily averaging time period in May 2013	81
Figure 3.44	12th Highest predicted concentration level of PM_{10} for daily averaging time period in May 2013	82
Figure 3.45	13th Highest predicted concentration level of PM_{10} for daily averaging time period in May 2013	82
Figure 3.46	14th Highest predicted concentration level of PM_{10} for daily averaging time period in May 2013	83
Figure 3.47	15th Highest predicted concentration level of PM_{10} for daily averaging time period in May 2013	83
Figure 3.48	16th Highest predicted concentration level of PM_{10} for daily averaging time period in May 2013	84
Figure 3.49	17th Highest predicted concentration level of PM_{10} for daily averaging time period in May 2013	84
Figure 3.50	18th Highest predicted concentration level of PM_{10} for daily averaging time period in May 2013	85
Figure 3.51	19th Highest predicted concentration level of PM_{10} for daily averaging time period in May 2013	85
Figure 3.52	20th Highest predicted concentration level of PM_{10} for daily averaging time period in May 2013	86
Figure 3.53	21st Highest predicted concentration level of PM_{10} for daily averaging time period in May 2013	86
Figure 3.54	22nd Highest predicted concentration level of PM_{10} for daily averaging time period in May 2013	87
Figure 3.55	23rd Highest predicted concentration level of PM_{10} for daily averaging time period in May 2013	87
Figure 3.56	24th Highest predicted concentration level of PM_{10} for daily averaging time period in May 2013	88
Figure 3.57	25th Highest predicted concentration level of PM_{10} for daily averaging time period in May 2013	88
Figure 3.58	26th Highest predicted concentration level of PM_{10} for daily averaging time period in May 2013	89
Figure 3.59	27th Highest predicted concentration level of PM_{10} for daily averaging time period in May 2013	89
Figure 3.60	28th Highest predicted concentration level of PM_{10} for daily averaging time period in May 2013	90
Figure 3.61	29th Highest predicted concentration level of PM_{10} for daily averaging time period in May 2013	90
Figure 3.62	30th Highest predicted concentration level of PM_{10} for daily averaging time period in May 2013	91
Figure 3.63	31st Highest predicted concentration level of PM_{10} for daily averaging time period in May 2013	91

Figure 3.64	Predicted concentration level of PM_{10} for monthly averaging time period in the month of January 2013	92
Figure 3.65	Predicted concentration level of PM_{10} for monthly averaging time period in the month of May 2013	93
Figure 3.66	Q-Q plots of predicted and observed concentration for daily averaging time for all four receptors in the month of January 2013	109
Figure 3.67	Q-Q plots of predicted and observed concentration for monthly averaging time for all four receptors in the month of January 2013	110
Figure 3.68	Q-Q plots of predicted and observed concentration for daily averaging time for all four receptors in the month of May 2013	111
Figure 3.69	Q-Q plots of predicted and observed concentration for monthly averaging time for all four receptors in the month of May 2013	112
Figure 4.1	Terrain Contours of Mine 'B' at 50 m depth	115
Figure 4.2	Terrain Contours of Mine 'B' at 100 m depth	116
Figure 4.3	Terrain Contours of Mine 'B' at 150 m depth	117
Figure 4.4	Terrain Contours of Mine 'B' at 200 m depth	118
Figure 4.5	Terrain Contours of Mine 'B' at 250 m depth	119
Figure 4.6	50 m , 100 m and 150 m depth section of Mine 'B'	120
Figure 4.7	200 m and 250 m depth section of Mine 'B'	121
Figure 4.8	Wind - rose diagram of Easterly wind variation	126
Figure 4.9	Wind - rose diagram of Westerly wind variation	127
Figure 4.10	Wind - rose diagram of Southerly wind variation	128
Figure 5.1	Contours of PM_{10} concentrations generated from the dust sources at 50 m depth of mine 'B' with the easterly wind	129
Figure 5.2	Contours of PM_{10} concentrations generated from the dust sources at 100 m depth of mine 'B' with the easterly wind	130
Figure 5.3	Contours of PM_{10} concentrations generated from the dust sources at 150 m depth of mine 'B' with the easterly wind	131
Figure 5.4	Contours of PM_{10} concentrations generated from the dust sources at 200 m depth of mine 'B' with the easterly wind	132
Figure 5.5	Contours of PM_{10} concentrations generated from the dust sources at 250 m depth of mine 'B' with the easterly wind	134
Figure 5.6	Contours of PM_{10} concentrations generated from the dust sources at 50 m depth of mine 'B' with the westerly wind	135
Figure 5.7	Contours of PM_{10} concentrations generated from the dust sources at 100 m depth of mine 'B' with the westerly wind	136
Figure 5.8	Contours of PM_{10} concentrations generated from the dust sources at 150 m depth of mine 'B' with the westerly wind	137
Figure 5.9	Contours of PM_{10} concentrations generated from the dust sources at 200 m depth of mine 'B' with the westerly wind	138

Figure 5.10	Contours of PM_{10} concentrations generated from the dust sources at 250 m depth of mine 'B' with the westerly wind	139
Figure 5.11	Contours of PM_{10} concentrations generated from the dust sources at 50 m depth of mine 'B' with the southerly wind	141
Figure 5.12	Contours of PM_{10} concentrations generated from the dust sources at 100 m depth of mine 'B' with the southerly wind	142
Figure 5.13	Contours of PM_{10} concentrations generated from the dust sources at 150 m depth of mine 'B' with the southerly wind	143
Figure 5.14	Contours of PM_{10} concentrations generated from the dust sources at 200 m depth of mine 'B' with the southerly wind	144
Figure 5.15	Contours of PM_{10} concentrations generated from the dust sources at 250 m depth of mine 'B' with the southerly wind	145
Figure 5.16	Comparative variation of the maximum dust concentration generated from the dust sources for the different depths of the mine 'B' with the different wind directions	146
Figure 5.17	Contours of PM_{10} concentrations generated from the internal overburden dump at 50 m depth of the mine 'B' with the easterly wind	148
Figure 5.18	Contours of PM_{10} concentrations generated from the internal overburden dump at 100 m depth of the mine 'B' with the easterly wind	149
Figure 5.19	Contours of PM_{10} concentrations generated from the internal overburden dump at 150 m depth of the mine 'B' with the easterly wind	150
Figure 5.20	Contours of PM_{10} concentrations generated from the internal overburden dump at 200 m depth of the mine 'B' with the easterly wind	151
Figure 5.21	Contours of PM_{10} concentrations generated from the internal overburden dump at 250 m depth of the mine 'B' with the easterly wind	152
Figure 5.22	Contours of PM_{10} concentrations generated from the internal overburden dump at 50 m depth of the mine 'B' with the westerly wind	154
Figure 5.23	Contours of PM_{10} concentrations generated from the internal overburden dump at 100 m depth of the mine 'B' with the westerly wind	155
Figure 5.24	Contours of PM_{10} concentrations generated from the internal overburden dump at 150 m depth of the mine 'B' with the westerly wind	156
Figure 5.25	Contours of PM_{10} concentrations generated from the internal overburden dump at 200 m depth of the mine 'B' with the westerly wind	157
Figure 5.26	Contours of PM_{10} concentrations generated from the internal overburden dump at 250 m depth of the mine 'B' with the westerly wind	158
Figure 5.27	Contours of PM_{10} concentrations generated from the internal overburden dump at 50 m depth of the mine 'B' with the southerly wind	160

Figure 5.28	Contours of PM_{10} concentrations generated from the internal overburden dump at 100 m depth of the mine 'B' with the southerly wind	161
Figure 5.29	Contours of PM_{10} concentrations generated from the internal overburden dump at 150 m depth of the mine 'B' with the southerly wind	162
Figure 5.30	Contours of PM_{10} concentrations generated from the internal overburden dump at 200 m depth of the mine 'B' with the southerly wind	163
Figure 5.31	Contours of PM_{10} concentrations generated from the internal overburden dump at 250 m depth of the mine 'B' with the southerly wind	165
Figure 5.32	Comparative variation of the maximum dust concentration generated from the internal overburden dumps at the different depths of the mine 'B' with the different wind directions	166
Figure 5.33	Contours of PM_{10} concentrations generated from the haul road at 50 m depth of the mine 'B' with the easterly wind	168
Figure 5.34	Contours of PM_{10} concentrations generated from the haul road at 100 m depth of the mine 'B' with the easterly wind	169
Figure 5.35	Contours of PM_{10} concentrations generated from the haul road at 150 m depth of the mine 'B' with the easterly wind	170
Figure 5.36	Contours of PM_{10} concentrations generated from the haul road at 200 m depth of the mine 'B' with the easterly wind	171
Figure 5.37	Contours of PM_{10} concentrations generated from the haul road at 250 m depth of the mine 'B' with the easterly wind	172
Figure 5.38	Contours of PM_{10} concentrations generated from the haul road at 50 m depth of the mine 'B' with the westerly wind	174
Figure 5.39	Contours of PM_{10} concentrations generated from the haul road at 100 m depth of the mine 'B' with the westerly wind	175
Figure 5.40	Contours of PM_{10} concentrations generated from the haul road at 150 m depth of the mine 'B' with the westerly wind	176
Figure 5.41	Contours of PM_{10} concentrations generated from the haul road at 200 m depth of the mine 'B' with the westerly wind	177
Figure 5.42	Contours of PM_{10} concentrations generated from the haul road at 250 m depth of the mine 'B' with the westerly wind	179
Figure 5.43	Contours of PM_{10} concentrations generated from the haul road at 50 m depth of the mine 'B' with the southerly wind	180
Figure 5.44	Contours of PM_{10} concentrations generated from the haul road at 100 m depth of the mine 'B' with the southerly wind	181
Figure 5.45	Contours of PM_{10} concentrations generated from the haul road at 150 m depth of the mine 'B' with the southerly wind	182
Figure 5.46	Contours of PM_{10} concentrations generated from the haul road at 200 m depth of the mine 'B' with the southerly wind	183
Figure 5.47	Contours of PM_{10} concentrations generated from the haul road at 250 m depth of the mine 'B' with the southerly wind	184

Figure 5.48 Comparative variations of the maximum dust concentration generated from the haul roads at the different depth of the mine ‘B’ with the different wind directions 186