Líst of Tables

Table No.	Description	Page No.
2.1	Reported development carried out by various research groups to improve bioavailability of ATR	23
4.1	Chemicals used in study	37
4.2	The following major instruments were used in the various stages of the experimental study	38
4.3	List of surfactants used and their critical micelle concentration (cmc) with aggregation number (N)	46
4.4	List of selected independent and dependent variables of AERSNs with coded and exact value used in CCD	49
4.5	Experimental variables and their levels with coded and exact value in APLNs formulation preparation	50
4.6	List of selected ALPNs formulation/process and response variables with coded and exact value	50
4.7	Batches of AERSNs designed by central composite experimental design using 4 factors and five levels with coded value	51
4.8	APLNs batches designed by central composite design using four factors and five levels	52
4.9	ALPNs batches designed by central composite design using four factors and five levels	53
4.10	Treatment regimen during 4th-6th week to different group	60
5.1	Validation parameters of calibration plot developed by HPLC method	66
5.2	Extraction ratio of ATR from rat plasma	66
5.3	The intra-day and inter-day precision of the RP-HPLC method	67
5.4	Release kinetic parameters of optimized AERSNs in PBS (pH 7.4)	85
5.5	Pharmacokinetic parameters of AERSNs and drug suspension after single oral dose administration	89
5.6	Plasma efficacy and safety related biochemical parameters in different groups of rats at the end of 6 th week	93
5.7	Drug release kinetic models of Optimized batch	102
5.8	Pharmacokinetic parameters of APLNs and drug suspension after single dose administration	106
5.9	Plasma efficacy and safety related biochemical parameters in different groups of rats at the end of 6 th week	110
5.10	Drug release kinetic models of Optimized batch of ALPNs	118
5.11	Pharmacokinetic parameters of ATR after single dose oral administration of ALPNs and ATR suspension	121
5.12	Plasma efficacy and safety related biochemical parameters in different groups of rats at the end of 6^{th} week	125