

List of Figures

| Figure No. | Description | Page No. |
|-------------------|---|-----------------|
| 4.1 | FTIR spectrum of docetaxel (sample drug) | 55 |
| 4.2 | UV spectrum of Docetaxel in acetonitrile mixed with pH 7.4 PBS (30:70) | 57 |
| 4.3 | Preparation of PHBV PVA polymeric nanoparticles | 64 |
| 4.4 | Ishikawa fishbone diagram is showing risk assessment studies | 65 |
| 4.5 | Preparation of PHBV PF127 polymeric nanoparticles | 70 |
| 4.6 | Risk management flow chart containing material attributes and process parameters | 71 |
| 4.7 | Preparation of PHBV TPGS polymeric nanoparticles | 76 |
| 4.8 | Preparation of PHBV PLH PE-PEG polymeric nanoparticles | 80 |
| 5.1 | Calibration curve of docetaxel in acetonitrile and pH 7.4 saline phosphate buffer (70:30) at 231 nm | 91 |
| 5.2 | Graphical illustrations are representing the effect of independent variables (column temp., flow rate and % organic phase) on the number of theoretical plates. (A1 & A2) represents interaction effect of column temp. and flow rate on theoretical plate; (B1 & B2) describes interaction effect of column temp and % organic phase on theoretical plate; whereas (C1 & C2) represents interaction effect of flow rate and % organic phase on the theoretical plate | 95 |
| 5.3 | Optimized predicted values for factors as per Design Expert 10 | 96 |
| 5.4 | Chromatogram of docetaxel at 5, 10, 25, 50 and 100 ppm | 97 |
| 5.5 | Results of linearity data and the calibration curve of docetaxel in acetonitrile and PBS pH 7.4 (30:70) at 235 nm | 99 |
| 5.6 | Degradation data for docetaxel at various environmental conditions for 72 h | 101 |
| 5.7 | Pareto chart showing the influence of variables on responses; | 103 |

| | | |
|------|--|-----|
| | (A) influence of input variables on particle size, (B) influence of input variables on zeta potential, (C) influence of input variables on entrapment efficiency | |
| 5.8 | Contour and 3-D Response Surface Methodology showing effect of independent variables on responses; (PS) effect of input variables on particle size, (ZP) effect of input variables on zeta potential and (EE) effect of input variables on % entrapment efficiency | 105 |
| 5.9 | (a) HR-TEM image; (b) SEM image at 10KX of optimized formulation and (c) AFM micrographs | 110 |
| 5.10 | Time course of days represents the in vitro percentage cumulative release of the drug from NPs in PBS pH 7.4 | 111 |
| 5.11 | Pareto chart and half normal plot showing the influence of variables on responses; (A) influence of input variables on particle size, (B) influence of input variables on zeta potential, (C) influence of input variables on entrapment efficiency | 113 |
| 5.12 | Graphical illustrations of Contour and 3-D Response Surface Methodology showing the effect of independent variables on responses; (PS) effect of input variables on particle size, (ZP) effect of input variables on zeta potential and (EE) effect of input variables on percentage entrapment efficiency | 117 |
| 5.13 | (A) SEM micrographs; (B) HR-TEM image; (C) AFM image of the optimized formulation | 119 |
| 5.14 | Profile showing in vitro cumulative drug release percentage from PHBV NPs in PBS pH 7.4 | 120 |
| 5.15 | Pareto chart showing the influence of input variables on particle size and influence of input variables on entrapment efficiency | 123 |
| 5.16 | Contour, 3-D Response Surface Methodology and All factor plot showing the effect of independent variables on responses (effect of homogenizer speed and polymer concentration on particle size; effect of sonication time and polymer concentration on % entrapment efficiency) | 127 |
| 5.17 | (A) SEM image; (B) AFM micrographs; (C) HR-TEM images of optimized nanoformulation | 129 |
| 5.18 | In vitro cumulative percentage of the drug release from PHBV NPs in PBS pH 7.4 | 130 |

| | | |
|------|--|-----|
| 5.19 | Analysis of variance and lack of fit plots with error showing goodness of fit for optimized process | 134 |
| 5.20 | Optimized predicted set points of independent variables indicating the effects on particle size and entrapment efficiency and response distribution graph showing expected values of responses (particle size and entrapment efficiency) | 134 |
| 5.21 | 3-D response surface plot showing the effects on responses (effect of homogenizer speed and polymer concentration on particle size; effect of sonication time and polymer concentration on % entrapment efficiency) and contour surface plot showing the effects of independent variables on particle size and entrapment efficiency with their corresponding optimized set points | 135 |
| 5.22 | Results of particle size, PDI and zeta potential analysis for F4 optimized formulation | 138 |
| 5.23 | Multi y plot for percentage entrapment efficiency and particle size of different optimized formulations | 139 |
| 5.24 | (A) SEM, (B) HR-TEM and (C) AFM micrographs of optimized nanoformulation | 140 |
| 5.25 | ¹ H NMR spectra of F4 formulation before and after PLH-PE-PEG coating carried out in the pulsed FT mode at a 500 MHz resonance frequency for protons after dissolving samples in deuterated DMSO-d ₆ | 141 |
| 5.26 | Cumulative percentage drug release of all the formulations set against days | 142 |
| 5.27 | FTIR spectra of DTX, PHBV, F1, F2, F3 & F4 showing absorption bands | 145 |
| 5.28 | XRD analyses of DTX, PHBV, PM, F1, F2, F3 & F4 showing absorption bands | 146 |
| 5.29 | DSC thermograms of DTX, PHBV, F1, F2, F3 & F4 showing peaks | 147 |
| 5.30 | SDS-PAGE images of rat plasma with different optimized formulations along with rat plasma (RP), plasma containing drug (DRP), and free drug (D) | 148 |
| 5.31 | In vitro cytotoxicity study of negative control and optimized nanoparticles observed under the inverted microscope in | 150 |

| | | |
|------|--|-----|
| | MCF7 cell lines | |
| 5.32 | Percentage control growth versus concentration ($\mu\text{g}/\text{mL}$) followed in order of F4>F1>F2>F3 | 151 |
| 5.33 | (A) Intracellular uptake of formulations tagged with dye (FITC) demonstrated by confocal microscopy imaging; (B) photographs representing trypan blue assay on rat lymphocytes for cell viability studies | 152 |
| 5.34 | Image of red blood cells (RBC) observed after hemolysis and platelet aggregation studies by optical microscopy | 154 |
| 5.35 | Percentage hemolysis against time (h) upon i.v. administration | 154 |
| 5.36 | Plot of observed plasma concentration-time profile of the drug and prepared formulations (F1, F2, F3 & F4) after i.v. administration in rat | 156 |
| 5.37 | Experimental and simulated plasma concentration-time profile of optimized formulations | 158 |
| 5.38 | Observed in vitro- in vivo correlation of drug loaded optimized formulations | 159 |
| 5.39 | (A) Average body weight (g) of mouse mammary tumor model C3H/J altered with respect to number of days; (B) Relative tumor volume data curve of mouse mammary tumor model of control, pure drug, standard (STD) and optimized formulations (25 mg/Kg i.v. weekly x 4 weeks) from day 1 to 30; (C) Survival data (%) using Kaplan-Meier curve of mouse bearing mammary tumor model treated with all seven treatment groups by log-rank test; (D) T/C values from RTV data of mouse mammary tumor model C3H/J demonstrated significant activity of optimized formulations | 162 |
| 5.40 | Photographs representing tumor bearing mice on first and last days of treatment with the optimized formulations | 163 |
| 5.41 | Stability study concerning particle size, zeta potential and % entrapment efficiency for 6 months' period | 165 |