

Table of contents

List of figures	xv
List of tables	xxv
Nomenclature	xxv
1 Introduction	1
1.1 Classical systems and the thermodynamic averages	2
1.2 Active Matter: an out of equilibrium system	3
1.3 Details of Active matter	6
1.4 Categories of active matter systems	8
1.4.1 Polar and Apolar active matter	8
1.4.2 Dry and Wet active matter	9
1.5 Methods to study active matter	10
1.5.1 Agent based models: Vicsek Model	11
1.5.2 Agent based model : Langevin's dynamics	13

1.5.3	Hydrodynamics equations of motion	14
1.6	Systems with nematic interactions on a substrate : Active nematics	24
1.7	Domain growth and the length scale	27
1.8	Active matter with inhomogeneity	28
1.9	The Lorentz lattice gas	29
1.10	Summary of the coming chapters	35
2	Active nematics with quenched disorder	37
2.1	Introduction	37
2.2	Model and Numerical details	40
2.3	Steady state properties	42
2.4	Kinetics	46
2.4.1	Defect dynamics	46
2.4.2	Growth law and scaling properties	48
2.4.3	Morphology of ordered domains	49
2.5	Linearized hydrodynamic calculation of two-point correlation functions	49
2.6	Snapshots for $\Delta\theta$ and NOP	58
2.6.1	Snapshots for $\Delta\theta$	58
2.6.2	Fixed a_0 and varying h_0	58
2.6.3	Fixed disorder h_0 and varying activity a_0	59
2.7	Summary	59

3	Active nematic gel with quenched disorder	63
3.1	Introduction	63
3.2	Model and Numerical details	64
3.3	Results	67
3.3.1	Sapshots	67
3.3.2	Kinetics	69
3.4	Discussions	74
4	Effect of polydispersity on the dynamics of active Brownian particles	75
4.1	Introduction	75
4.2	Model and numerical details	78
4.3	Results	80
4.3.1	Enhanced Diffusivity for finite Polydispersity	80
4.3.2	Mobility order parameter	85
4.3.3	Phase diagram	86
4.4	System size independence	90
4.5	Discussion	90
5	Dynamics of particle moving in one dimensional Lorentz lattice gas	93
5.1	Introduction	93
5.2	Model and numerical details	96
5.3	Definitions: Ballistic, Anomalous diffusion and Confined motion	99

5.4	Results	99
5.4.1	Anomalous diffusion	103
5.5	Discussion	106
6	Dynamics of particle moving in a two dimensional Lorentz lattice gas	109
6.1	Introduction	109
6.2	Model and numerical details	111
6.3	Deterministic nature and closed trajectories	114
6.4	Results	116
6.4.1	Mean Square Displacement	116
6.4.2	Open and closed trajectories	118
6.4.3	Properties of Open Trajectories	119
6.5	Discussion	122
7	Summary and future scope	125
7.1	Summary	125
7.2	Future prospects	126
	References	127

List of figures

1.1	Schematic diagram of polar and apolar particles and their ordering states. Arrow shows the direction of motion, double arrow represents the head and tail symmetry.	9
1.2	Examples of the three types of configurations considered in [Mishra et al. (2016)] are shown. In (a) a completely random configuration is shown in which $r \simeq 1$. In (b) and (c), the entire lattice is divided in periodic blocks of size $l \times l$, where $l = 3$. The configuration in (b) is randomly periodic (RP) in that each block contains a single scatterer that is assigned to be a left rotator (open circle). The configuration in (c) is a completely periodic (CP) configuration, where the single left rotator is in the same relative position in each block. In both (b) and (c) the initial ratio of rotators is $r = 1/(l \times l - 1) = 1/8$.	34
2.1	(Color online) (a) NOP \mathbf{Q} vs. system size $N = K \times K$ for different h_0 (symbols) and $\sqrt{C_{\mathcal{Q}}}$, Eq. (2.3) (solid lines), (b) Scaled NOP, $\bar{Q} = Q \times N^{\mathcal{B}'/2}$ vs. $\bar{x} = N \times h_0^4$. (c) Probability distribution function $P(\Delta\theta)$ of angle fluctuations of angular orientation for different values of (h_0, a_0) i.e. $(0.0, 0.2)$, $(0.05, 0.2)$ and $(0.05, 0.3)$. Data for $h_0 = 0.05$, $a_0 = 0.3$ (data points) fitted with Gaussian for two distinct peaks (solid lines).	39

-
- 2.2 (Color online) Number fluctuation σ vs. $\langle N_s \rangle$ plot for different disorder strengths, and $K = 300$ 42
- 2.3 (Color online) (a) Snapshots of local NOP \mathcal{Q} : upper panel (P:1) is for clean-AN (i.e. $h_0 = 0.0$) and bottom panel (P:2) is for RFAN ($h_0 = 0.075$) and the number along the white arrow is the relative separation ($\Delta r(t)$) between the $+$ and $-1/2$ defects. (b) $\Delta r(t)$ vs. t plot for $h_0 = 0.0, 0.075$. u is the relative speed of defects defined as $u = |\frac{d}{dt}\Delta r(t)| \times 10^{-3}$ and plotted in the inset of (b). (c) Snapshot of density current near the defects for clean-AN, $h_0 = 0.0$ and (d) RFAN, $h_0 = 0.075$. Intensity of colors shows the magnitude of the density current. Data is generated for system size $N = 512^2$ 44
- 2.4 (Color online) Two-point correlation function $C_{\mathcal{Q},\rho}$ vs. scaled distance $r/L_{\mathcal{Q},\rho}(t)$. (a-b) Two-point correlation function for RFAN i.e. $h_0 = 0.1$ at different simulation time (t). (c-d) Two-point correlation function for different h_0 and at fixed simulation time $t = 10^5$ 46
- 2.5 (Color online) Plots of dynamic growth exponent $z_{eff}(h_0)$ vs. disorder strength h_0 (a), Δz vs. h_0 on log – log scale (b). 47
- 2.6 (Color online) Cusp's exponent α (slope of the plot) for Q and ρ field. Different symbols used for different values of disorder strength: $h_0 = 0.0$ (circle), $h_0 = 0.05$ (square), $h_0 = 0.075$ (diamond), $h_0 = 0.10$ (triangle). 50
- 2.7 (Color online) Snapshots of $\Delta\theta$ for $h_0 = 0.0, a_0 = 0.2$ (i), $h_0 = 0.05, a_0 = 0.2$ (ii) and $h_0 = 0.05, a_0 = 0.3$ (iii). 59

- 2.8 (Color online) Snapshots of local NOP, \mathcal{Q} for $a_0 = 0.2$ and for different disorder strength in the system. From top to bottom panel (P:1 to P:3) $h_0 = 0.0, 0.05$ and 0.1 respectively. Snapshots are generated at equal interval i.e. $t = 150000, 300000, 450000,$ and 600000 from left to right respectively. Notice the numbers on the color bars. 60
- 2.9 (Color online) Snapshots of local NOP, \mathcal{Q} for $h_0 = 0.05$ and for different activity in the system. From top to bottom panel (P:1 to P:3) $a_0 = 0.1, 0.2$ and 0.3 respectively. Snapshots are generated at equal interval i.e. $t = 150000, 300000, 450000,$ and 600000 from left to right respectively. Notice the numbers on the color bars. 60
- 2.10 (Color online) (a) Relative separation $\Delta(r)$ between $+1/2$ and $-1/2$ defects pair vs. time plot and (b) relative speed, $u = \left| \frac{d}{dt} \Delta r(t) \right| \times 10^{-3}$, of $\pm 1/2$ defects, for different value of a_0 and fixed $h_0 = 0.1$. t is the simulation time. 61
- 3.1 Snapshots at different time for $\alpha_2 < 0$ and $h_0 = 0.0$. Different snapshots are for (NOP) field $\mathcal{Q}(\mathbf{r}, t)$ (top panel), local density field $\rho(\mathbf{r}, t)$ (middle panel), and flow field $\mathbf{v}(\mathbf{r}, t)$ (bottom panel). A pair of $+1/2$ -defect and $-1/2$ -defect are indicated by the arrows in the first picture (top left). The range of color bars for NOP, density and flow fields are in range $(0.0, 0.5), (0.0, 0.82)$ and $(0.0, 0.016)$, respectively. 68
- 3.2 Snapshots of NOP $\mathcal{Q}(\mathbf{r}, t)$ at different time for $\alpha_2 < 0$; for different disorder strengths $h_0 = 0.0$ (top panel) and $h_0 = 0.2$ (bottom panel). Range of the color bars are same as given for fig. 3.1. 68

-
- 3.3 Two point correlation function, C_{ϱ} vs. r (top-left and top-right) and C_{ρ} vs. r (bottom-left and bottom-right) for different h_0 . All the data shown are at fixed simulation time $t = 50000$. Plot in left panel plots are for contractile case with $\alpha_2 = 0.4$, whereas right panel shows the plots for extensile case with $\alpha_2 = -0.4$ 70
- 3.4 Two point correlation function, C_{ϱ} vs. r for $h_0 = 0.05$ (top-left and top-right), and $h_0 = 0.2$ (bottom-left and bottom-right) for different t . Plot in left panel plots are for contractile case with $\alpha_2 = 0.4$, whereas right panel shows the plots for extensile case with $\alpha_2 = -0.4$ 70
- 3.5 Two point correlation function, C_{ρ} vs. r for $h_0 = 0.05$ (top-left and top-right), and $h_0 = 0.2$ (bottom-left and bottom-right) for different t . Plot in left panel plots are for contractile case with $\alpha_2 = 0.4$, whereas right panel shows the plots for extensile case with $\alpha_2 = -0.4$ 71
- 3.6 Correlation length, L_{ϱ} vs. t (top-left and top-right) and L_{ρ} vs. t (bottom-left and bottom-right) for different h_0 . Plot in left panel plots are for contractile case with $\alpha_2 = 0.4$, whereas right panel shows the plots for extensile case with $\alpha_2 = -0.4$ 71
- 3.7 Correlation length, L_{ϱ} vs. $t/\ln(t)$ (top-left and top-right) and L_{ρ} vs. t (bottom-left and bottom-right) for different h_0 (log-log scale). Plot in left panel plots are for contractile case with $\alpha_2 = 0.4$, whereas right panel shows the plots for extensile case with $\alpha_2 = -0.4$. Dashed line corresponds to correlation length in equilibrium *xy - model* 72

- 3.8 Scaled two point correlation function, C_{ϱ} vs. r (top-left and top-right) and C_{ρ} vs. r (bottom-left and bottom-right) for different h_0 . x-axis is scaled as $r \rightarrow r/L_{\varrho,\rho}(t)$. All the data shown are at fixed simulation time $t = 50000$. Plot in left panel plots are for contractile case with $\alpha_2 = 0.4$, whereas right panel shows the plots for extensile case with $\alpha_2 = -0.4$ 72
- 3.9 Scaled two point correlation function, C_{ϱ} vs. r for $h_0 = 0.05$ (top-left and top-right), and $h_0 = 0.2$ (bottom-left and bottom-right) for different t . x-axis is scaled as $r \rightarrow r/L_{\varrho,\rho}(t)$. Plot in left panel plots are for contractile case with $\alpha_2 = 0.4$, whereas right panel shows the plots for extensile case with $\alpha_2 = -0.4$ 73
- 3.10 Scaled two point correlation function, C_{ρ} vs. r for $h_0 = 0.05$ (top-left and top-right), and $h_0 = 0.2$ (bottom-left and bottom-right) for different t . x-axis is scaled as $r \rightarrow r/L_{\varrho,\rho}(t)$. Plot in left panel plots are for contractile case with $\alpha_2 = 0.4$, whereas right panel shows the plots for extensile case with $\alpha_2 = -0.4$ 73
- 4.1 (Color online) Snapshot of the system for a non-zero polydispersity i.e. $\varepsilon = 0.5$. Disks resembles the active Brownian particles with their radius represented by color bar, and the arrows shows their velocity direction. 78
- 4.2 (Color online) (a) Mean squared displacement, $\Delta(t)$ vs. t for different activity (v) and fixed polydispersity index, $\varepsilon = 8.33 \times 10^{-1}$. (b) $\Delta(t)$ vs. t for different ε and fixed activity, $v = 3.33 \times 10^{-2}$. (c) Diffusion coefficient, $D(t)$ vs. t for different activity (v) and fixed polydispersity index, $\varepsilon = 8.33 \times 10^{-1}$. (d) $D(t)$ vs. t for different ε and fixed activity, $v = 3.33 \times 10^{-2}$ 81

- 4.3 (Color online) Effective diffusion coefficient, D_{eff} vs. activity v for different polydispersity index (ε), for $\phi = 0.65$ (a), 0.75 (b), 0.85 (c). Effective diffusion coefficient D_{eff} vs. polydispersity index, ε for different activity (v). Symbols are from the simulation and the solid lines are the fit (Eq. 4.4) to the data points. Different plots are for different values of $\phi = 0.65, 0.75, 0.85$ (d-f), respectively. Error bars are smaller than the size of the symbols. 81
- 4.4 (Color online) Scaled diffusivity, D_{eff}/v_0^β vs. scaled PDI, $\varepsilon v^{-\alpha}$ for $\phi = 0.65, 0.75, 0.85$ (a-c), respectively; where $\alpha = -0.2$ and $\beta = 2.0$ 83
- 4.5 (Color online) Percentage change in the effective diffusion coefficient, ΔD_{eff} vs. PDI, ε for different activity (v), for $\phi = 0.65$ (a), 0.75 (b), 0.85 (c). 84
- 4.6 (Color online) Snapshots at equal time for $\varepsilon = 1.66 \times 10^{-1}$ (a), and $\varepsilon = 6.66 \times 10^{-1}$ (b), for fixed activity, $v = 3.33 \times 10^{-2}$, and $\phi = 0.65$. Rattlers are represented by blue disks whereas non-rattlers are in red. Arrows on the disks shows their velocity direction. 84
- 4.7 (Color online) (a) Probability distribution function, $P(\psi)$ vs ψ for different activity (v) and fixed polydispersity index, $\varepsilon = 8.33 \times 10^{-1}$. (b) $P(\psi)$ vs ψ for different ε and fixed activity, $v = 3.33 \times 10^{-2}$. (c) $P(\psi_{bs})$ vs ψ_{bs} for different activity (v) and fixed polydispersity index, $\varepsilon = 8.33 \times 10^{-1}$. (d) $P(\psi_{bs})$ vs ψ_{bs} for different ε and fixed activity, $v = 3.33 \times 10^{-2}$ 85

- 4.8 (Color online) (a) Plot for the radial distribution function $g(r)$ for different PDI, ε on semi-log scale (main), inset: plot on linear scale. We keep the activity fixed in this plot. (b) Number Fluctuation plot for $(v, \varepsilon) = (3.33 \times 10^{-3}, 1.66 \times 10^{-1})$ circles: solid-jammed phase, $(3.33 \times 10^{-3}, 6.66 \times 10^{-1})$ squares: liquid-jammed phase, $(2.00 \times 10^{-2}, 1.66 \times 10^{-1})$ diamonds: MIPS-liquid phase, $(3.33 \times 10^{-2}, 0.0)$ triangle up: MIPS-liquid phase, $(3.33 \times 10^{-2}, 8.33 \times 10^{-1})$ triangle left: pure liquid phase. The dashed line corresponds to $\Delta N/\sqrt{N} = 1$. Both the plots are for $\phi = 0.85$ and the system size $L = 40$ 87
- 4.9 (Color online) Phase diagram for $\phi = 0.65$ (a), $\phi = 0.75$ (b) and $\phi = 0.85$ (c): circles, squares, triangles and diamonds represents the solid-jammed phase, liquid-jammed phase, MIPS-liquid phase and *pure liquid* phase, respectively. Color bar shows the value of D_{eff} for a given (ε, v) 88
- 4.10 (Color online) (a) Effective diffusion coefficient D_{eff} vs. polydispersity ε for different v . (b) Scaled diffusivity, D_{eff}/v_0^β vs. scaled PDI, $\varepsilon v^{-\alpha}$, where $\alpha = -0.2$ and $\beta = 2.0$. Error bars are of the size of the symbols. (c) Phase diagram: circles, squares, triangles and diamonds represents the solid-jammed phase, liquid-jammed phase, MIPS-liquid phase and *pure liquid* phase, respectively. Color bar shows the value of D_{eff} for a given (ε, v) . All the data are generated for $\phi = 0.85$ and system size $L = 30$ 91
- 5.1 A cartoon of one dimensional Lorentz lattice Gas : filled circles:- reflector, empty circles:- transmitters. Arrow shows the direction of particle's velocity. (a) Part of a typical initial configuration. Interaction of particle with a scatterer and state of scatterer one-step after interaction when (b) $\alpha = 0$ i.e. pure flipping and (c) $\alpha = 1$ i.e. pure fixed. 96

- 5.2 (Color online) MSD ($\Delta(\tau)$) vs. time (τ) (upto 10^4 before the asymptotic behaviour observed) Plot: $r = 0.0$ (a), $r = 0.15$ (b), $r = 0.5$ (c) and $r = 1.0$ (d) for $\alpha = 0.0$ (circle), 0.1 (square), 0.2 (diamond), 0.3 (triangle up), 0.4 (triangle left), 0.5 (triangle down), 0.6 (triangle right), 0.7 (plus), 0.8 (x), 0.9 (star), 1.0 (open circle). Black and Red dotted lines have slope = 1 and 2 respectively. Data is averaged over 1000 ensembles. 98
- 5.3 Three different trajectory when particle motion is confined (or periodic): (a), (b) and (c) shows the trajectory for system parameter set $(\alpha, r)=(1.0,0.1)$, $(1.0,0.5)$ and $(1.0,0.9)$ respectively. Figure (d),(e) and (f) shows the RoG of the particle for system parameter set $(\alpha, r)=(1.0,0.1)$, $(1.0,0.5)$ and $(1.0,0.9)$ respectively. Data for $R_g = \sqrt{\Delta(t)}$ is averaged over 100 realisations. 100
- 5.4 MSD ($\Delta(\tau)$) vs. time (τ) plot when (a) $\alpha = 0.003$ and $r = 0.9$ and 0.6, (b) $\alpha = 0.999$ and $r = 0.3$, (c) different set of α and r 104
- 5.5 $F(t) = \frac{d}{dt}N(t)$ vs. $G(t) = \frac{1}{N(t)}$ plot when (a) $\alpha = 0.3$ and $r = 0.3$, (b) $\alpha = 0.003$ and $r = 0.6$, (c) $\alpha = 0.003$ and $r = 0.9$. Data (black solid line) is fitted linearly (red dashed line) which give sthe slope c 104
- 5.6 Plot of $r_{visited}$ vs time for $\alpha = 0.01$ and $r = 0.6$ 104
- 5.7 Probability distribution of particles position (bars) and fitted with Gaussian distribution (solid line) when (a) $r = 0.5$ and $\alpha = 0.5$, (b) $r = 0.5$ and $\alpha = 0.9$, (c) $r = 0.9$ and $\alpha = 0.5$, (d) $r = 0.9$ and $\alpha = 0.9$ 106
- 5.8 Phase diagram in the plane of (r, α) . Data points shows the asymptotic phases of the system, based on the values of MSD exponent (β) for different set of (r, α) , when the system reaches the steady state. 107

- 6.1 Schematic diagram of two dimensional Lorentz Lattice Gas : (a) Initial configuration of LLG with left-rotators (filled circles), right-rotators (open circles) and vacancies (empty sites). (b) State of the particle before (t) and after ($t + 1$) interaction with rotator; the arrow shows the direction of particle's velocity. 112
- 6.2 (a) Variation of MSD $\Delta(t)$ with time (t) for different vacancy densities ρ_V (Main). Inset plot shows the crossover from diffusion to subdiffusion. (b) Zoomed in image of plot (a) for $t < 1000$. (c) Effective diffusion coefficient D_{eff} vs t for different values of ρ_V . (d) $P(r,t)$ vs $r(t)$ at different times for a fully occupied lattice. 114
- 6.3 (a) Variation of probability of being in an open trajectory $P_o(\rho_V, t)$ with time (t) on a log-log plot for different values of ρ_V in the first phase, when $P_o(\rho_V, t)$ varies as a power law. The straight lines represent the fitted power law functions for each value of ρ_V . (b) Variation of $P_o(\rho_V, t)$ with time on a semi-log(y) plot during the second phase of motion, when $P_o(\rho_V, t)$ decays exponentially with time. As the particles transition from the first phase of motion to the second, $P_o(\rho_V, t)$ decays exponentially with t and deviates from the fitted function. 118
- 6.4 Exponent γ vs ρ_V plot. Value of γ is extracted by fitting power law to $P_0(\rho_V, t)$ vs t plot in diffusive regime i.e. simulation time $\mathcal{O}(10^4)$ 120
- 6.5 Variation of $\Delta_o(t)$ with time for different values of ρ_V . The dashed line represents a straight line with slope 1. (b) Fractal dimension and (c) hyperscaling condition $(2/d_f - \gamma)$ of trajectories for different values of vacancy density in the first phase of motion. 120

List of tables

- 1.1 Different cases in a two dimensional Lotentz lattice gas model. $C = C_L + C_R$,
where, C_L and C_L are the left and right rotators respectively. 31

- 5.1 List of the values of D_{eff} and D_{est} for different values of r when $\alpha = 0.5$. . . 105