## List of Tables

Page No.

Table 4.1	The absolute error $ER(x,1)$ with various choices of $M$ and $N$	120
Table 4.2	The absolute error $ER(x,1)$ with various choices of $M$ and $N$	122
Table 4.3	The absolute error $ER(x,1)$ with various choices of $M$ and $N$	124
Table 4.4	The absolute error $ER(x,1)$ with various choices of $M$ and $N$	126
Table 4.5	The absolute error $ER(x,1)$ with various choices of $M$ and $N$	128
Table 5.1	The absolute error $ER(x,1)$ with various choices of $M$ and $N$	143
Table 5.2	The absolute error $ER(x,1)$ with various choices of $M$ and $N$	145
Table 5.3	The absolute error $ER(x,1)$ with various choices of $M$ and $N$	147
Table 5.4	The absolute error $ER(x,1)$ with various choices of $M$ and $N$	149
Table 5.5	The absolute error $ER(x,1)$ with various choices of $M$ and $N$	152
Table 7.1	The absolute errors at different t with fixed values of $x = y$	173
	and $N = M_1 = M_2 = 3$ for Example 1	
Table 7.2	The absolute errors at different choices of $x = y$ with fixed	173
	values of t and $N = M_1 = M_2 = 3$ for Example 1	
Table 7.3	The absolute errors at various $t$ with fixed values of $x = y$ and	175
	$N = M_1 = M_2 = 3$ for Example 2	
Table 7.4	$N = M_1 = M_2 = 3$ for Example 2 The absolute errors at various values of $x = y$ with fixed values	175
Table 7.4		175
Table 7.4 Table 7.5	The absolute errors at various values of $x = y$ with fixed values	175 177
	The absolute errors at various values of $x = y$ with fixed values of t and $N = M_1 = M_2 = 3$ for Example 2	
	The absolute errors at various values of $x = y$ with fixed values of t and $N = M_1 = M_2 = 3$ for Example 2 The absolute errors at different t with fixed values of $x = y$	

- **Table 7.7** The absolute errors at various choices of t with fixed values of179x = y and  $N = M_1 = M_2 = 3$  for Example 4
- **Table 7.8** The absolute errors at various choices of x = y with fixed179values of t and  $N = M_1 = M_2 = 3$  for Example 4
- **Table 7.9** The absolute errors at different choices t with fixed values of181x = y and  $N = M_1 = M_2 = 3$  for Example 5
- **Table 7.10** The absolute errors at different choices of x = y with fixed182values of t and  $N = M_1 = M_2 = 3$  for Example 5