

List of Figures

Page
No.

- Fig. 2.1 Plots of $u(x, t)$ vs. t for $a_0 = 10, \beta = 0.5, k = 2, n = 1, x = 0.5$, and for different values of α 39
- Fig. 2.2 Plots of $u(x, t)$ vs. t for $a_0 = 10, \beta = 0.5, k = 2, n = 2, x = 0.5$, and for different values of α 39
- Fig. 2.3 Plots of $u(x, t)$ vs. t for $a_0 = 10, \beta = 0.5, k = 2, n = 1, x = 0.5$, and for different values of α 40
- Fig. 2.4 Plots of $u(x, t)$ vs. t for $a_0 = 10, \beta = 0.5, k = 2, n = 2, x = 0.5$, and for different values of α 40
- Fig. 2.5 Plots of $u(x, t)$ vs. t for $a_0 = -10, \beta = 0.5, k = 2, n = 1, x = 0.5$, and for different values of α 41
- Fig. 2.6 Plots of $u(x, t)$ vs. t for $a_0 = -10, \beta = 0.5, k = 2, n = 2, x = 0.5$, and for different values of α 41
- Fig. 2.7 Plots of $u(x, t)$ vs. t for $a_0 = -10, \beta = 0.5, k = 2, n = 1, x = 0.5$, and for different values of α 42
- Fig. 2.8 Plots of $u(x, t)$ vs. t for $a_0 = -10, \beta = 0.5, k = 2, n = 2, x = 0.5$, and for different values of α 42
- Fig. 2.9 Plots of $\langle X^2(t) \rangle$ vs. t for $a_0 = 5, \beta = 0.5$ and for different values of α 43

| | | |
|----------|--|----|
| Fig.2.10 | Plots of $\langle X^2(t) \rangle$ vs. t for $a_0 = 10, \beta = 0.5$ and for different values of α | 43 |
| Fig.2.11 | Plots of $\langle X^2(t) \rangle$ vs. t for $a_0 = -2, \beta = 0.5$ and for different values of α | 44 |
| Fig.2.12 | Plots of $\langle X^2(t) \rangle$ vs. t for $a_0 = -5, \beta = 0.5$ and for different values of α | 44 |
| Fig. 4.1 | Plots of exact residual error E_m vs. \hbar for $\alpha = 0.8 + 0.2 \frac{xt}{LT}, L = 10, T = 1$ | 69 |
| Fig. 4.2 | Evolution curve of $u(x=0.5, t)$ vs. t at $\hbar = -0.975296$ | 70 |
| Fig. 4.3 | Plots of exact residual error E_m vs. \hbar for $\alpha = xt$ | 70 |
| Fig. 4.4 | Evolution curve of $u(x, t)$ vs. x and t for $\alpha = xt$ | 71 |
| Fig. 5.1 | Plots of exact residual error E_m vs. \hbar for $a = 1, \varepsilon = 1$ and $\alpha = 1$ | 84 |
| Fig. 5.2 | Plots of exact residual error E_m vs. \hbar for $a = 1, \varepsilon = 1$ and $\alpha = 0.75$ | 84 |
| Fig. 5.3 | Plots of exact residual error E_m vs. \hbar for $a = 1, \varepsilon = 1$ and $\alpha = 0.5$ | 85 |

| | | |
|----------|---|-----|
| Fig. 5.4 | Phase Portrait between u_1 and u_2 (a) for $a=1, \varepsilon=1$ and $\alpha=0.5$ (b) for $a=1, \varepsilon=1$ and $\alpha=0.75$ (c) for $a=1, \varepsilon=1$ and $\alpha=1$ | 86 |
| Fig. 5.5 | Phase Portrait between u_1 and u_2 for $a=1, \alpha=0.75$ and $\varepsilon=1,2,3,4$ | 87 |
| Fig. 5.6 | Phase Portrait between u_1 and u_2 (a) for $a=1, \varepsilon=0.5$ and $\alpha=0.75$ (b) for $a=1, \varepsilon=0.5$ and $\alpha=0.85$ (c) for $a=1, \varepsilon=0.5$ and $\alpha=0.95$ | 88 |
| Fig. 5.7 | Phase Portrait between u_1 and u_2 (a) for $a=1, \varepsilon=8$ and $\alpha=0.75$ | 88 |
| Fig. 6.1 | Plot of $u(r, t)$ for the case (i) for $f(r) = r, g(r) = 1$ | 99 |
| Fig. 6.2 | Plot of $u(r, t)$ for the case (ii) for $f(r) = \sqrt{r}, g(r) = \frac{1}{\sqrt{r}}$ | 99 |
| Fig. 6.3 | Plot of $u(r, t)$ for the case (iii) for $f(r) = r^2, g(r) = r$ | 100 |
| Fig. 6.4 | Solution Plot for $u(r, t)$ in all the three cases for $r=0.5$ | 101 |
| Fig. 6.5 | Solution Plot for $u(r, t)$ in all the three cases for $r=0.9$ | 101 |
| Fig. 6.6 | Solution Plot for $u(r, t)$ in all the three cases (i), (ii) & (iii) respectively for $r=0.5$ for $\beta = 2$ and $3/2$ | 103 |