

TABLE OF CONTENTS

CHAPTER 1 INTRODUCTION	1-68
1.1.General Introduction	1
1.2.Types of corrosion	2
1.3.Forms of corrosion	6
1.4.Basic concept of corrosion	11
1.5. Factors influencing metallic corrosion	12
1.6.Corrosion thermodynamics	17
1.7.Kinetics of corrosion	23
1.8.Corrosion protection methods	28
1.9.Mechanism of corrosion inhibition in acid solution	35
1.10. Adsorptive Mechanism of corrosion inhibition	36
1.11. Inhibition of mild steel corrosion in minerals acids	42
1.12. Graphical representation of kinetic data	43
1.13. Electrochemical Impedance Spectroscopy (EIS)	49
1.14. Application of Theoretical Methods in Corrosion Inhibition Studies	52
1.15. Heterocyclic compounds as corrosion inhibitors: literature survey with particular emphasis of quantum chemical calculations and molecular dynamics simulations	58
CHAPTER 2: EXPERIEMNTAL	69-85
2.1. Materials	69
2.2. Heterocyclics inhibitors	70
2.3. Instruments and techniques	78

2.4. Determination of Thermodynamic Parameters	81
2.5. SEM, EDX and AFM measurements	82
2.6. Computational studies	83
CHAPTER 3: RESULTS AND DISCUSSIONS	86-177
3.1. 5-arylpyrimido-[4, 5-b] quinoline-diones (APQDs) as corrosion inhibitors	86
3.2. 2-amino-4-arylquinoline-3-carbonitriles (AACs) as corrosion inhibitors	111
3.3. 2, 4-diamino-5-(phenylthio)-5H-chromeno [2, 3-b] pyridine-3-carbonitriles (DHPCs) as corrosion inhibitors	132
3.4. 3-amino alkylated indoles (AAIs) as corrosion inhibitors	156
CHAPTER 4: SUMMARY	178-184
4.1. Summary	178
4.2. Conclusions	184
REFERENCES	185-213
LIST OF PUBLICATIONS	214-218

LIST OF FIGURES

Figure 1.1: liquid metal corrosion of mild steel surface by molten Copper at 1100 ⁰ C (MP: 1085 ⁰ C)	4
Figure 1.2: E-pH diagram for a generic metal M which forms hydroxides	20
Figure 1.3: Pourbaix diagram for iron/water system at 298 K	22
Figure 1.4: Schematic representation of Evans diagram for iron in acid solution	43
Figure 1.5: (a) Anodic control (b) cathodic control, (c) mixed control of a corrosion process.	44
Figure 1.6: Anodic and cathodic half-cell reactions present simultaneously on a corroding zinc surface	45
Figure 1.7: Polarization of anodic and cathodic half-cell reactions for zinc in acid solution to give a mixed potential, E_{corr} and a corrosion rate (current density)	46
Figure 1.8: “ E vs. i ” for Fe in acid solution	47
Figure 1.9: Extrapolated Tafel curves	48
Figure 1.10: Complex plane impedance spectrum	50
Figure 1.11: Nyquist plot along with constant phase element	50
Figure 3.1.1: (a) Arrhenius plots for the corrosion of mild steel in 1 M HCl (b) Langmuir adsorption isotherm plots for the adsorption of APQDs on mild steel surface in 1M HCl	92
Figure 3.1.2: Polarization curves for mild in absence and presence of different concentrations APQDs	95

Figure 3.1.3: Nyquist curves for mild in absence and presence of different concentrations of APQDs 97

Figure 3.1.4: Bode impedance modulus ($\log f$ vs $\log |Z|$) and phase angle ($\log f$ vs α^0) plots for mild steel in 1 M HCl in absence and different of different concentrations of (a) APQD-1, (b) APQD-2, (c) APQD-3 and (d) APQD-4. 99

Figure 3.1.5: SEM images of mild steel surfaces: abraded (a), in 1 M HCl in the absence of APQDs (b), and in 1 M HCl in the presence of 20 mgL⁻¹ of APQD-1 (c), APQD-2 (d), APQD-3 (e) and APQD-4 (f) 101

Figure 3.1.6: AFM images of mild steel: (a) in absence of APQDs and in the presence of 20 mgL⁻¹ of (b) APQD-1, (c) APQD-2, and (d) APQD-3 (e) APQD-4 102

Figure 3.1.7: Optimized molecular structures of (a) APQD-1, (b) APQD-2, (c) APQD-3 and (d) APQD-4 103

Figure 3.1.8: The frontier molecular orbital (left-hand side: HOMO; and right-hand side: LUMO) of the studied APQDs (a) APQD-1 (b) APQD-2, (c) APQD-3, and (d) APQD-4 105

Figure 3.1.9: Fukui indices for the electrophilic (f_x^+) and nucleophilic (f_x^-) sites for (a) APQD-1, (b) APQD-2, (c) APQD-3, and (d) APQD-4. (All the surfaces were visualized at isosurface value of 0.004, except for the f_x^+ of (c) APQD-3, and (d) APQD-4 for which isosurface value of 0.0028 and 0.0025 respectively, were used for better visualization) 108

Figure 3.1.10: Side view equilibrium adsorption of APQD-1, APQD-2, APQD-3 and APQD-4 on Fe (110) surface (a) before and (b) after molecular dynamics simulations 109

Figure 3.2.1: (a) Variation of inhibition efficiency with AACs concentration (b) Arrhenius plots of $\log C_R$ vs. $1000/T$ (c) Langmuir isotherm plot for the adsorption of the AACs on mild steel surface in 1M HCl	115
Figure 3.2.2: Polarization curves for mild in absence and presence of different concentration of AACs	117
Figure 3.2.3: Nyquist plot for mild steel in 1 M HCl without and with different concentrations of AACs	120
Figure 3.2.4: Bode ($\log f$ vs $\log Z $) and phase angle ($\log f$ vs α^0) plots for mild steel in 1 M HCl in absence and presence of different concentration of AACs	122
Figure 3.2.5: SEM images of mild steel surfaces: abraded (a), in absence of AACs (b) in the presence of 40 mgL^{-1} of AAC-1(c), AAC-2(d), and AAC-3(e)	123
Figure 3.2.6: EDX images of mild steel: (a) in absence of AACs and in the presence of 40 mgL^{-1} of (b) AAC-1, (c) AAC-2, and (d) AAC-3	125
Figure 3.2.7: AFM images of mild steel: (a) in absence of AACs and in the presence of 40 mgL^{-1} of (b) AAC-1, (c) AAC-2, and (d) AAC-3	126
Figure 3.2.8: Optimized molecular structures of the studied compounds showing the dihedral angles a-b-c-d (in degrees)	127
Figure 3.2.9: The frontier molecular orbital of studied AACs (a) AAC-1 (left, HOMO; right, LUMO), (b) AAC-2 (left, HOMO; right, LUMO) and (c) AAC-3 (left, HOMO; right, LUMO)	129
Figure 3.3.1: (a) Arrhenius plots for the corrosion of mild steel in 1 M HCl in the presence of the studied inhibitors (b) Langmuir adsorption isotherms for mild steel in 1 M	

Figure 3.3.2: Polarization curves recorded for mild steel in the absence and presence of different concentrations DHPCs 139

Figure 3.3.3: Nyquist plots recorded for mild steel in 1 M HCl in the absence and presence of different concentrations of (a) DHPC-1, (b) DHPC-2, and (c) DHPC-3 142

Figure 3.3.4: Bode impedance modulus ($\log f$ vs $\log |Z|$) and phase angle ($\log f$ vs α°) plots for mild steel in 1 M HCl in the absence and presence of different of different concentrations of (a) DHPC-1, (b) DHPC-2, and (c) DHPC-3 144

Figure 3.3.5: SEM images of mild steel surfaces in 1 M HCl in the absence of DHPCs (a), and in 1 M HCl in the presence of optimum concentration of DHPC-1 (b), DHPC-2 and (c), DHPC-3 (d) 145

Figure 3.3.6: AFM images of mild steel surfaces in 1 M HCl in the absence of DHPCs (a), and in 1 M HCl containing optimum concentration of DHPC-1 (b)HPC-2 and (c), DHPC-3(d) 146

Figure 3.3.7: Optimized molecular structures of (a) DHPC-1, (b) DHPC-2, and (c) DHPC-3 148

Figure 3.3.8: The frontier molecular orbital (left-hand side: HOMO; and right-hand side: LUMO) of the studied DHPCs (a) DHPC-1 (b) DHPC-2, and (c) DHPC-3 . 149

Figure 3.3.9: Fukui indices f^+ and f^- corresponding to the atomics sites for the nucleophilic and electrophilic attacks respectively in (a) DHPC-1, (b) DHPC-2, and (c) DHPC-3 (*isosurface value = 0.003*) 152

Figure 3.3.10: Side and top views of the most stable adsorption models of (a) DHPC-1, (b) DHPC-2, and (c) DHPC-3 on Fe (110) surface using quench molecular dynamic	154
Figure 3.4.1: (a) Arrhenius plots of $\log C_R$ vs. $1000/T$ (b) Langmuir isotherm plot for the adsorption of the AAI's on mild steel surface in 1M HCl	161
Figure 3.4.2: Polarization curves recorded for mild steel in the absence and presence of different concentrations of AAI's	163
Figure 3.4.3: Nyquist plots recorded for mild steel in 1 M HCl in the absence and presence of different concentrations of AAI's.	165
Figure 3.4.4: Bode impedance modulus ($\log f$ vs $\log Z $) and phase angle ($\log f$ vs α^0) plots for mild steel in 1 M HCl in the absence and different of different concentrations of AAI's	168
Figure 3.4.5: SEM images of mild steel surfaces in 1 M HCl in the absence of AAI's (a), and in 1 M HCl in the presence of optimum concentration of AAI-1 (b), AAI-2 and (c), AAI-3(d)	169
Figure 3.4.6: AFM images of mild steel surfaces in 1 M HCl in the absence of AAI's (a), and in 1 M HCl in the containing optimum concentration of AAI-1 (b), AAI-2 and (c), AAI-3(d)	170
Figure 3.4.7: Optimized molecular structures of studied AAI's, (a) AAI-1, (b) AAI-2 and (c) AAI-3	172
Figure 3.4.8: The frontier molecular orbital (left-hand side: HOMO; and right-hand side: LUMO) of the studied APQDs (a) AAI-1 (b) AAI-2 and (c) AAI-3	173

Figure 3.4.9: Side view equilibrium adsorption of AAI-1, AAI-2, and AAI-3 on Fe (110) surface
(left hand side: before; and right hand side: after molecular dynamics simulations)