

References

- Abazajian et al. (2004). The second data release of the Sloan Digital Sky Survey. *Astron. J.*, **128**, 502.
- Abazajian et al. (2005). The third data release of the Sloan Digital Sky Survey. *Astron. J.*, **129**, 1755.
- Abdussattar and Prajapati, S.R. (2011). Role of deceleration parameter and interacting dark energy in singularity avoidance. *Astrophys. Space Sci.*, **331**, 657.
- Adhav, K.S. (2012a). LRS Bianchi type-I cosmological model in $f(R, T)$ theory of gravity. *Astrophys. Space Sci.*, **339**, 365.
- Adhav, K. (2011). Bianchi type-VI0 cosmological models with anisotropic dark energy. *Astrophys. Space Sci.*, **323**, 497.
- Adhav, K. (2012b). Bianchi type-III cosmic string cosmological model in $f(R)$ theory of gravity. *Bulg. J. Phys.*, **39**, 197.
- Adhav, K., Ugale, M., Kale, C., and Bhende, M. (2007). Bianchi type-III anisotropic cosmological models with varying . *Bulg. J. Phys.*, **34**, 260.

References

- Agmohammadi, et al. (2009). Resembling holographic dark energy with $f(R)$ gravity as scalar field and ghost dark energy with tachyon scalar fields. *Phys. Scr.*, **80**, 065008.
- Ahmed, N. and Pradhan, A. (2013). Bianchi type-V cosmology in $f(R, T)$ gravity with Λ . *Int. J. Theor. Phys.*, **53**, 289.
- Akarsu, O. and Dereli, T. (2012). A comparison of the LVDP and Λ CDM cosmological. *Int. J. Theor. Phys.*, **51**, 2995.
- Al-Rawaf, A. (1998). *Mod. Phys. Lett. A*, **13**, 429.
- Al-Rawaf, A. and Taha, M. (1996). Cosmology of general relativity without energy-momentum conservation. *Gen. Rel. Grav.*, **28**, 935.
- Allemandi, G., Borowiec, A., Francaviglia, M., and Odintsov, S.D. (2005). Dark energy dominance and cosmic acceleration in first-order formalism. *Phys. Rev.D*, **72**, 063505.
- Alvarez, E., G.M.B. (1983). Entropy from extra dimensions. *Phys. Rev. Lett.*, **51**, 931.
- Ananda, K.N. and Carloni, S. (2008). Evolution of cosmological gravitational waves in $f(R)$ gravity. *Phys. Rev. D*, **77**, 024033.
- Appelquist, T. and Chodos, A. (1983). Quantum effects in Kaluza-Klein theories. *Phys. Rev. Lett.*, **28**, 141.
- Arbab, A.L. and Cosmo, J. (2003a). Cosmological consequences of a built-in cosmological constant model. *Antiparticle Phys.*, **05**, 8.
- Arbab, A.L. and Cosmo, J. (2003b). Cosmic acceleration with a positive cosmological constant. *Class. Quantum Gravity*, **20**, 93.
- Asseo, E. and Sol, H. (1987). Extragalactic magnetic fields. *Phys. Rep.*, **148**, 307.

References

- Astier et al. (2006). The supernova legacy survey: measurement of ω_M, ω_Λ and ω from the first year data set. *Astron. Astrophys.*, **31**, 447.
- Azizi, T. (2013). Wormhole geometries in f(R,T) gravity. *Int J Theo Phys*, **52**, 3486.
- Baade, W. and Zwicky, F. (1934). On Super-novae. *Proc. Nat. Acad. Sci.*, **20**, 254.
- Baghel, P.S. and Singh, J.P. (2012). Bianchi typeV universe with bulk viscous matter and time varying gravitational and cosmological constants. *Research in Astron. Astrophys.*, **12**, 1457.
- Bali, R. and Anjali (2004). Bianchi type-I bulk viscous fluid string dust magnetizedcosmological model in general relativity. *Pramana J. Phys.*, **63**, 481.
- Bali, R., Banerjee, R., and Banerjee, S.K. (2008a). Bianchi type VI0 magnetized bulkviscous massive string cosmological model in general relativity. *Astrophys.Space Sci.*, **317**, 21.
- Bali, R. and Chandnani, N.K. (2008). Bianchi type-III bulk vicous dust filled in lyra geometry. *Astrophys. Space. Sci.*, **318**, 225.
- Bali, R. and Chandnani, N.K. (2009). Bianchi type V barotropic perfect fluid cosmological model in Lyra geometry. *Int. J. Theor. Phys.*, **1523**, 48.
- Bali, R. and Dave, S. (2003). Bianchi type- III string cosmological model with bulkviscous fluid in general relativity. *Astrophys. Space Sci.*, **288**, 503.
- Bali, R. and Meena, B.L. (2002). Tilted cosmological models filled with disordered radiations in general relativity. *Astrophys. Space Sci.*, **281**, 565.
- Bali, R. and Pradhan, A. (2007). Bianchi type-III string cosmological models with time dependent bulk viscosity. *Chin. Phys. Lett.*, **24**, 585.

References

- Bali, R., Pradhan, A., and Hassan, A. (2008b). Bianchi type VI0 magnetized barotropic bulk viscous fluid massive string universe in general relativity. *Int. J. Theor. Phys.*, **47**, 2594.
- Bali, R. and Sharma, K. (2003). Tilted bianchi type I stiff fluid magnetized cosmological model in general relativity. *Astrophys. Space Sci.*, **283**, 11.
- Bali, R. and Swati (2012). LRS bianchi type-II inflationary universe with mass less scalar field and time varying Λ . *Chin. Phys. Lett.*, **29**, 080404.
- Bamba, K., Capozziello, S., Nojir, S., and Odintsov, S.D. (2012). Dark energy cosmology the equivalent description via different theoretical models and cosmography tests. *Astrophys Space Sci*, **342**, 155.
- Banerjee, A. and Sanyal, A.K. (1988). Irrotational Bianchi V viscous fluid cosmological with heat flux. *Gen. Rel. Gravit.*, **20**, 103.
- Barrow, J. (1984). Helium formation in cosmologies with anisotropic curvature. *Mon. Not. Astron.*, **211**, 221.
- Barrow, J. (1986). Comparison of theory and experiment in peristaltic transport. *The Journal of Fluid Mechanics*, **180**, 335.
- Barrow, J. (1997). Cosmological limits on slightly skew stresses,. *Phys. Rev.D*, **55**, 7541.
- Beesham, A. (1988). FLRW cosmological models in Lyras manifold with time dependent displacement field. *Aust. J. Phys.*, **41**, 833.
- Belinski, V.A. and Khalatnikov, I.M. (1976). Inuence of viscosity on the character of cosmological evolution. *JETP*, **42**, 205.

References

- Bennet et al. (2003). The third data release of the Sloan Digital Sky Survey. *Astrophys. J. Suppl. Ser.*, **148**, 1.
- Berman, M.S. (1983). A special law of variation for Hubble's parameter. *Nuovo Cimento B*, **74**, 182.
- Berman, M.S. and Gomide, F.M. (1988). Cosmological models with variable gravitational and cosmological constant. *Gen. Relativ. Gravit.*, **20**, 191.
- Bertolami, O., Bohmer, C.G., Harko, T., and Lobo, F.S.N. (2007). Extra force in $f(R)$ modified theories of gravity. *Phys. Rev. D*, **75**, 104016.
- Bogolyubov, O.I. (1985). Method in qualitative theory of dynamical systems in astrophysics and gas dynamics. *Sov. Math.*
- Burd, A. and Tavakol, R. (1993). Invariant Lyapunov exponents and chaos in cosmology. *Phys. Rev. D*, **47**, 5336.
- Caldwell, R.R., Dave, R., and Steinhardt, P.J. (1998). Cosmological imprint of an energy component with general equation of state. *Phys. Rev. Lett.*, **80**, 1582.
- Capozziello, S., Cardone, V.F., and Francaviglia, M. (2006c). $f(R)$ theories of gravity in the Palatini approach matched with observations. *Gen. Relativ. Gravit.*, **8**, 711.
- Capozziello, S., Cardone, V.F., and Troisi, A. (2006b). Dark energy and dark matter as curvature effects. *JCAP*, **8**, 01.
- Capozziello, S. and Faraoni, V. (2010). Beyond Einstein gravity: A Survey of gravitational theories for cosmology and astrophysics. *Springer*.
- Capozziello, S., Nojiri, S., Odintsov, S., and Troisi, A. (2006a). Cosmological viability

References

- of f(R) gravity as an ideal fluid and its compatibility with a matter dominated phase. *Phys. Lett. B*, **632**, 597.
- Capozziello, S., Stabile, A., and Troisi, A. (2007). Spherically symmetric solutions in f(R) gravity via the noether symmetry approach. *Class. Quant. Grav*, **24**, 2153.
- Capozziello, S., Stabile, A., and Troisi, A. (2008). Spherical symmetry in f(R)-gravity. *Class. Quant. Grav*, **25**, 085004.
- Carloni, S., Dunsby, P.K.S., and Troisi, A. (2008). Evolution of density perturbations in f(R) gravity. *Phys Rev. D*, **77**, 024024.
- Carroll, S.M. (2001). The Cosmological Constant. *Liv. Rev. Relat.*, **4**, 41.
- Carroll, S.M., Duvvuri, V., Trodden, M., and Turner, M.S. (2004). Is cosmic speed-up due to new gravitational physics ?. *Phys. Rev. D*, **70**, 043528.
- Carvalho, J., Lima, A.S., and Waga, I. (1992). Cosmological consequences of a time-dependent Λ term. *Phys. Rev. D*, **46**, 2404.
- Chakraborty, S. (2013). An alternative f (R, T) gravity theory and the dark energy problem. *arXiv*, 1212.3050.
- Chakraborty, S. and Ghosh, S. (2000). Generalized scalar tensor theory in four and higher dimension. *Int. J. Mod. phys. D*, **9**, 543.
- Chandel, S. and Ram, S. (2013). Anisotropic Bianchi type-III perfect uid cosmological models inf(R,T) theory of gravity. *Indian J. Phys*, **48**, 362.
- Chaterjee, S. (1992). Some remarks on luminosity distance and nucleosynthesis in higher dimensional cosmology. *Astrophys. J*, **397**, 1.

References

- Chaubey, R. and Shukla, A. (2013). A new class of Bianchi cosmological models in $f(R,T)$ gravity. *Astrophys. Space Sci.*, **343**, 415.
- Chen, W. and Wu, Y. (1990). Implications of a cosmological constant varying as r^{-2} . *Phys. Rev. D*, **41**, 695.
- Chiba, T., Smith, T.L., and Erickcek, A.L. (2007). Solar system constraints to general $f(R)$ gravity. *Phys. Rev. D*, **75**, 124014.
- Chodos, A. and Detweiler, S. (1980). Where has the fifth dimension gone ? *Phys. Rev. D*, **21**, 2167.
- Cognola, G., Elizalde, E., Nojiri, S., Odintsov, S.D., and Zerbini, S. (2006). Dark energy in modified gauss-bonnet gravity: Late-time acceleration and the hierarchy problem. *Phys. Rev. D*, **73**, 084007.
- Coley, A.A. (1990). Bianchi V imperfect fluid cosmology. *Gen. Rel. Gravit.*, **22**, 3.
- Coley, A.A. (2003). Dynamical system and cosmology. *Klu. Aca. Pub.*
- Coley, A.A.. and Wainright, J. (1991). Qualitative analysis of two fluidBianchi cosmologie. *Class. Quantum Grav.*, **9**, 651.
- Coley, A.A. and Hoogan, R.J. (1994). Qualitative analysis of diagonal Bianchi typeV imperfect fluid cosmological models. *J. Math. Phys.*, **35**, 4117.
- Coley, A.A. and Tupper, B. (1984). Viscous fluid collapse. *Phys. Rev. D*, **29**, 2701.
- Collins, C.B. (1971). More qualitative cosmology. *Commun. Math. Phys.*, **23**, 137.
- Copeland, E., Sami, M., and S.Tsujikawa (2006). Dynamics of dark energy. *Int.J. Mod. Phys.D*, **15**.

References

- Cunha, J.V., Lima, J.S., and N. Pires, N. (2002). Deflationary Λ cosmology: observational expressions. *Astronomy Astrophys*, **390**, 809.
- de sitter, W. (1917). Further remarks on the solutions of the field-equations of einsteins, theory of gravitation. *Proc. Acad. Sci., Amsterdam*, **19**, 1217.
- Dolgov, A.D. (1983). In the very early universe. eds. G.W. Gibbons, S.W.Hawking, S.T.C. Siklos, Cambridge University Press, Cambridge.
- Dolgov, A.D. (1990). Basics of modern cosmology. *Edditions Frontiers*.
- Dolgov, A.D. (1997). Higher spin fields and the problem of the cosmological constant. *Phys. Rev. D*, **55**, 5881.
- Einstein, A. (1915). *Pre. Akad. Wiss Ber*, 778.
- Einstein, A. (1917). Kosmogische betrachtungen zur allgemeinen relativistic-theorie. *Preuss . Akad. Wiss. Berlin, Sitzber*.
- Eisenstein, D.J., Zehavi, I., Hogg, D.W., Scoccimarro, R., and et al (2005). Detection of the baryon acoustic peak in the large-scale correlation function of sdss luminous red galaxies. *Astrophys. J*, **633**, 560.
- Everett, A.E. (1981). Cosmic strings in unified gauge theories. *Phys. Rev.*, **24**, 858.
- Faraoni, V., Jensen, M.N., and Theuerkauf, S.A. (2006)). Non-chaotic dynamics in general-relativistic and scalartensor cosmology. *Class. Quantum Grav*, **23**, 4215.
- Forgacs, P. and Horvath, Z. (1979). On the influence of extra dimensions on the homogeneous isotropic universe. *Gen. Relative. Gravit.*, **11**, 205.
- Freund, P.O. (1982). Kaluza-Klein cosmologies. *Nucl. Phys. B*, **209**, 146.

- Friedmann, A. (1922). *Zeitchrift fut Physik*, **10**, 377.
- Friedmann, A. (1924). *Zeitchrift fut Physik*, **21**, 326.
- Frieman, J.A. and Waga, I. (1998). Constraints from high redshift supernovae upon scalar field cosmologies. *Phys.Rev. D*, **57**, 4642.
- Goenner, F.M. and Kowalewsky, F. (1987). Exact anisotropic viscous fluid solutions of einstein's equations. *Gen. Rel. Gravit*, **21**, 467.
- Gross, D.J. and Perry, M.J. (1983). Magnetic monopoles in Kaluza-Klein theories. *Nucl. Phys. B*, **29**, 226.
- Guarnizo, A., Casta, L., and Tejeiro, J.M. (2011). Geodesic deviation equation in f(R) gravity. *Gen. Relativ. Gravit.*, **194**, 194.
- Gupta, R.C. and Pradhan, A. (2010). A theoretically-explained new-variant of modified-newtonian-dynamics (MOND). *Int. J. Theor. Phys.*, **49**, 821.
- Guth, A. (1981). Inflationary universe: A possible solution to horizon and flatness problem. *Phys. Rev. D*, **23**, 347.
- Hajj-Boutros, J. (1986a). Lecture notes in physics, gravitation. *Geometry and Relativity Physics*, **212**, 51.
- Hajj-Boutros, J. (1986b). A method for generating Bianchi type-II cosmological models. *J. Math. Phys.*, **27**, 1592.
- Halford, W.D. (1970). Cosmological theory based Lyras geometry. *Austr. J.Phys*, **23**, 863.
- Halford, W. (1972). Scalar-tensor theory of gravitation in a Lyra manifol. *J.Math. Phys.*, **13**, 1699.

References

- Harko, T. (2008). Modified gravity with arbitrary coupling between matter and geometry. *Phys. Lett. B*, **669**, 376.
- Harko, T., Koivisto, T.S., and Lobo, F.S.N. (2011a). Palatini formulation of modified gravity with a nonminimal curvature-matter coupling. *Mod.Phys.Lett.A*, **26**, 146.
- Harko, T. and Lobo, F.S.N. (2010). Extended f(R,Lm) theories of gravity. *Eur. Phys. J. C*, **70**, 373.
- Harko, T., Lobo, F.S.N., Nojiri, S., and Odintsov, S.D. (2011b). f(R,T) gravity. *Phys. Rev.D*, **84**, 2011.
- Hawkins, E. (2003). The 2df galaxy redshift survey: correlation functions, peculiar velocities and the matter density of the universe. *Man. Not. R. Astron. Soc.*, **346**, 78.
- Hinshaw, G., Larson, D., Komatsu, E., , Spergel, D.N., and et al (2009). Five -year wilkinson microwave anisotropy probe observations: Data processing, sky maps and basic results. *Astrophys. J. Suppl. Ser*, **180**, 225.
- Houndjo, M.J.S. (2012). Reconstruction of f(R, T) gravity describing matter dominated and accelerated phases. *Int. J. Mod. Phys. D*, **21**, 1250003.
- Hoyle, F. (1948). A new model for the expanding universe. *Mon. Not. R. Astron. Soc*, **108**, 327.
- Hoyle, F. and Narlikar, J.V. (1964b). Time symmetric electrodynamics and the arrow of time in cosmology. *Proc. Ray. Soc. Land. Ser A*, **277**, 1–23.
- Hoyle, F. and Narlikar, J. (1964a). On the avoidance of singularity in C-field cosmology. *Proc. Ray . Soc. Land. Ser. A*, **278**, 465.

References

- Huang, B., Li, S., and Ma, Y. (2010). Five-dimensional metric $f(R)$ gravity and the accelerated universe. *Phys. Rev.D*, **81**, 064003.
- Hubble, E. and L.Humason, M. (1934). The velocity-distance relation among extra-galactic nebulae. *Proc, Nature Acad. Sci. Philad.*, **20**, 264.
- Huterer, D. and Turner, M.S. (1999). Prospects for probing the dark energy via supernova distance measurements. *Phys. Rev. D*, **60**, 081301.
- Jafe, T.R., Banday, J., Eriksen, H.K., Gorski, K.M., and et al (2005). Evidence of vorticity and shear at large angular scales in the WaMP data:A vilation of cosmological isotropy. *Astrophys J. Lett*, **629**, 1.
- Johri, V.B. and Desikan, K. (1994). Cosmological models with constant deceleration parameter in brans-dicke theory. *Gen. Rel. Grav*, **26**, 1217.
- Kaluza, T. (1921). Zum unit atsproblem der physik, sitz. pressure, akadwiss. *Ber. Phys. Math.*, **33**, 966.
- Karami, K. and Abdolmaleki, A. (2009). Reconstructing interacting new agegraphic polytropicgas model in nonat frw. *Eur. Phys. J. C*, **64**, 85.
- Khadekar, G.S. and Tade, S. (2007). String cosmological models in five dimensional bimetric theory of gravitation. *Astrophy. Space Sci.*, **310**, 47.
- Kibble, T.W.B. (1976). Topology of cosmic domains and strings. *J. Phys.A, Math. Gen.*, **9**, 1387.
- Klein, O. (1926a.). Quantentheorie and finite dimensional relativitats theories. *Z. Phys* ., **37**, 895.

References

- Klein, O. (1926b). The atomicity of electricity as a quantum theory law. *Nature*, **118**, 516.
- Koivisto, T. (2007). Viable palatini-f(R) cosmologies with generalized dark matter. *Phys. Rev. D*, **76**, 043527.
- Krori, K.D., Chaudhury, T., and Mahanta, C. (1994). Some exact solutions in string cosmology. *Gen. Relativ. Gravit.*, **26**, 265.
- Lemaitre, G. (1917). *Ann. Soc. Sci.*, **47**, 19.
- Letelier, P.S. (1978). Unified theory of direct interaction between particles, strings, and membranes. *J.Math. Phys.*, **19**, 1908.
- Letelier, P.S. (1983). String cosmologies. *Phys. Rev.D*, **28**, 2414.
- Liddle, A.R. and Scherrer, R.J. (1999). Classification of scalar field potentials with cosmological scaling solutions. *Phys. Rev. D*, **59**, 023509.
- Linde, A. (1974). Is the cosmological constant a constant. *JETP Lett.*, **19**, 183.
- Lorentz, D. (1980). An exact Bianchi- type- II cosmological model with electromagnetic field. *Phys. Lett. A*, **79**, 19.
- Lyra, G. (1951). Sber eine modification der riemannschen geometrie. *Math.Z.*, **54**, 52.
- Maartens, R. (1995). Dissipative cosmology. *Class. Quantum Gravity*, **12**, 1455.
- Mazumdar, A. (1994). Solutions of LRS Bianchi type I space-time filled with a perfect fluid. *Gen. Relativ. Gravit.*, **26**, 307.
- Mishra, K. and Sahoo, P. (2014). Bianchi type VIh perfect fluid cosmological model f(R,T) theory. *Astrophys. Space Sci.*, **352**, 331.

References

- Misner, C.W. (1967). Transport process in the primordial fireball. *Nature*, **214**, 40.
- Misner, C.W. (1968). The isotropy of the universe. *Astrophys. J.*, **151**, 431.
- Mohanty, G. and Sahoo, P.K. (2007). *Gen. Relativ. Gravit.*, **312**, 321.
- Mohanty, G., Mahanta, K., and Sahoo, R.R. (2006). Non-existence of five dimensional perfect fluid cosmological model in Lyra manifold. *Astrophys. Space Sci.*, **306**, 269.
- Mohanty, G. and Pradhan, B.D. (1983). *Int. J. Theor. Phys.*, **14**, 303.
- Murphy, G.L. (1973). Big-bang model without singularities. *Phys. Rev. D*, **8**, 4231.
- M.Youshimura (1984). Effective action and cosmological evolution of scale factors in higher-dimensional curved spacetime. *Phys. Rev. D*, **30**, 344.
- Myrzakulov, R. (2012). FRW cosmology in F(R,T)gravity. *Eur. Phys. J.*, **72**, 2203.
- Naidu, R., Reddy, D., Ramprasad, T., and Ramana, K. (2013). Bianchi type-V bulk viscous string cosmological model in f (R,T) gravity. *Astrophys Space Sci*, **348**, 247.
- Nojiri, S. and Odintsov, S.D. (2003a). Modified gravity with negative and positive powers of the curvature: unification of the inflation and of the cosmic acceleration. *Phys. Rev.D*, **68**, 123512.
- Nojiri, S. and Odintsov, S.D. (2003b). Quantum de sitter cosmology and phantom matter. *Phys. Lett. B*, **562**, 147.
- Nojiri, S. and Odintsov, S.D. (2004.). Gravity assisted dark energy dominance and cosmic acceleration. *Phys. Lett. B*, **99**, 137.
- Nojiri, S. and Odintsov, S.D. (2006a). Modified f(R) gravity consistent with realistic cosmology: from matter dominated epoch to dark energy universe. *Phys. Rev. D*, **74**, 086005.

References

- Nojiri, S. and Odintsov, S.D. (2006b). On the way from matter-dominated era to dark energy universe. *Phys.Rev.D*, **74**, 086009.
- Nojiri, S. and Odintsov, S.D. (2007). Introduction to modified gravity and gravitational alternative for dark energy. *Int. J. Geom. Meth. Mod. Phys.*, **4**, 115.
- Nojiri, S. and Odintsov, S.D. (2008). Modified gravity as realistic candidate for dark energy, inflation and dark matter. *Phys. Lett. B*, **659**, 821.
- Nojiri, S. and Odintsov, S.D. (2011). Unified cosmic history in modified gravity: from F(R) theory to lorentz non-invariant models. *Phys. Rept.*, **505**, 59.
- Nojiri, S. and Odintsov, S. (2010). A proposal for covariant renormalizable field theory of gravity. *Phys Lett. B*, **91**, 60.
- Overdin, J.M. and Cooperstock, F.I. (1998). Evolution of the scale factor with a variable cosmological term. *Phys. Rev. D*, **58**, 043506.
- Ozer, M. and Taha, M.O. (1987). Amodel of the universe free of cosmological problems. *Nucle. Phys.B*, **287**, 776.
- Padmanabhan, T. (2003). *Phys. Repts*, **380**, 235.
- Padmanabhan, T. and Chitre, S.M. (1987). Vicious universes. *Phys. Lett. A*, **120**, 433.
- Peebles, P.J.E. and Ratra, B. (2003). The cosmological constant and dark energy. *Rev. Mod. Phys.*, **75**, 559.
- Percival, W., Cole, S., Eisenstein, D.J., Nichol, R.C., and et al (2007). Measuring the baryon acoustic oscillation scale using the sdss and 2dfgrs. *Mon. Not. Roy. Astron. Soc*, **381**, 1053.

- Perlmutter et al. (1997). Measurements of the cosmological parameters Ω and Λ from the first seven supernovae at $z = 0.35$. *Astrophys. J.*, **483**, 565.
- Perlmutter et al. (1999). Measurement of ω and Λ from 42 high-redshift supernovae. *Astrophys.J.*, **517**, 565.
- Poplawski, N. (2007). Alagrangian description of interacting dark energ. *arXiv:gr-qc*, 0608031.
- Poplawski, N.J. (2011). Matter-antimatter asymmetry and dark matter from torsion. *Phys Rev. D*, **83**, 084033.
- Pradhan, A. (2007). A new class of inhomogeneous string cosmological models in general relativity. *Astrophy. Space Sci.*, **312**, 145.
- Pradhan, A. (2009a). Thick domain walls in Lyra geometry with bulk viscosity. *Comm.Theor. Phys*, **51**, 378.
- Pradhan, A. (2009b). Cylindrically symmetric viscous fluid universe in Lyra geometry. *J. Math .Phys*, **50**, 022501.
- Pradhan, A. and Chouhan, D.S. (2011). Anisotropic Bianchi type-I models in string cosmology. *Astrophys. Space Sci.*, **331**, 697.
- Pradhan, A., Rai, V., and Otarod, S. (2006). Plane symmetric inhomogeneous bulk viscous domain wall in Lyra geometry. *Fizika B*, **15**, 57.
- Pradhan, A. and R.Bali (2008). Magnetized Bianchi type V I0 barotropic massive string universe with decaying vacuum energy density Λ . *EJTP*, **19**, 91.

References

- Pradhan, A., Srivastava, D., and Khadekar, G.S. (2008). Can Bianchi type-IIcosmological models with a decay law for Λ term be compatible with recent observations ? *Romanian Reports in Phy*, **60**, 3.
- Pradhan, A. and Vishwakarma, A.K. (2004). A new class of LRS Bianchi type-I cosmological models in Lyra geometry. *J.Geom. Phys*, **49**, 332.
- Pradhan, A., Yadav, V.K., and Chakraborty, I. (2001). Bulk viscous FRW cosmology in Lyra geometry. *Int. J. Mod. Phys. D*, **10**, 339.
- Rahaman, F., Begum, N., Bag, G., and Bhui, B.C. (2005). Cosmological models with negative constant deceleration parameter in Lyra geometry. *Astrophys. Space Sci.*, **299**, 211.
- Rahaman, F., Chakraborty, S., and Bera, G. (2002). Inhomogeneous cosmological model in Lyra geometryl. *Int. J.Mod. Phys.D*, **11**, 1501.
- Rahaman, F.S., Chakraborty, S., Das, M., and Hossain, B.J. (2003). Higherdimensional string theory in Lyra geometry. *Pramana J. Phys*, **60**, 453.
- Ram, S. (1985). Bianchi type VI₀ solutions in modified Brans-Dicke cosmology. *J. Math. Phys.*, **26**, 2916.
- Ram, S. (1986). Perfect fluid models of Bianchi type VI₀ in modified Brans-Dicke cosmology. *J. Math. Phys.*, **27**, 660.
- Ram, S. and Priyanka (2013). Some Kaluza-Klein cosmological models in $f(R, T)$ gravity theory. *Astrophys. Space Sci*, **347**, 389.
- Ram, S., Priyanka, and Singh., M.K. (2013). Anisotropic cosmological models in $f(R, T)$ theory of gravitation. *Pramana J. Phys.*, **81**, 67.

References

- Ram, S. and Singh, P. (1992). Anisotropic cosmological models of Bianchi types III and V in Lyra's geometry. *Int. J. Theor. Phys.*, **31**, 2095.
- Ram, S., Verma, M.K., and Zeyauddin, M. (2009). Homogeneous anisotropic cosmological models with viscous fluid and heat flow in lyra's geometry. *Modern Physics Lett. A*, **24**, 1847.
- Ram, S., Zeyauddin, M., and Singh, C.P. (2010). Anisotropic Bianchi type V perfect fluid cosmological models in Lyras geometry. *J.Geom.Phys*, **60**, 1671.
- Ratra, B. and Peebles, P.J.E. (1988). Cosmological consequences of a rolling homogeneous scalar field. *Rev. Mod. Phys.*, **37**, 12.
- Reddy, D.R.K. (2003). A string cosmological model in brans-dicke theory of gravitation,. *Astrophys. Space Sci.*, **286**, 365.
- Reddy, D.R.K. (2005). Plane symmetric cosmic strings in lyra manifold. *Astrophys. Space Sci.*, **300**, 381.
- Reddy, D.R.K. (2009). Kantowaski-sachs inflationary universe in general relativity. *Int.J. Theor.Phys*, **48**, 2884.
- Reddy, D.R.K. and Naidu, N.L. (2007). Five dimensional string cosmological models in a scalar-tensor theory of gravitation. *Astrophys. Space Sci.*, **307**, 395.
- Reddy, D.R.K., Naidu, R.L., and Naidu, K.D. (2013). LRSBianchi type-II universe with cosmic strings and bulk viscosity in a modified theory of gravity. *Astrophys. Space Sc*, **323**, 362.
- Reddy, D.R.K., Naidu, R.L., and Satyanarayan, B. (2012a). Kaluza-Klein cosmological model in $f(R,T)$ gravity. *Int. J. Theor. Phys.*, **51**, 3222.

References

- Reddy, D.R.K., Rao, S., Koteswara, M.V., and Rao, G. (2006). Axially symmetric radiating model in Brans - Dicke cosmology. *Astrophys Space Sci.*, **306**, 171.
- Reddy, D., Santikumar, R., and Naidu, R. (2012b). Bianchi type-III cosmological model in $f(R, T)$ theory of gravity. *Astrophys. Space Sci.*, **249**, 342.
- Riess et al. (1998). Observational evidence from supernovae for an accelerating universe and a cosmological constant. *Astrophys. J.*, **116**, 1009.
- Riess et al. (2004). Type ia supernova discoveries at $z > 1$, from the Hubble space telescope: evolution for past deceleration,. *Astrophys. J.*, **607**, 665.
- Ringstrom, H. (2001). The Bianchi IX attractor. *Anales Henri Poincaré*, **2**, 405.
- Rosquist, K. and Jantzen, R.T. (1988). Unified regularization of Bianchi cosmologies. *Phys. Rep.*, **166**, 89.
- Roy, S.R. and Singh, J.P. (1983). Some inhomogeneous viscous fluid cosmological models of plane symmetry. *Acta Physiaca Austriaca*, **55**, 57.
- Roy, S.R. and Tiwari, O.P. (1983). Some Bianchi VI0 cosmological models with free gravitational field of the magnetic type. *Ind. J. Pure Appl. Math.*, **14**, 233.
- Rubin, V. and Ford, W. (1970). Rotation from a spectro scope survey of emission regions. *Astrophys. J.*, **159**, 379.
- Saha, B. and Yadav, A.K. (2012). Dark energy model with variable q and ω in LRS Bianchi-II space-time. *Astrophys. Space Sci.*, **341**, 651.
- Sahdev, D. (1984). Towards a realistic Kaluza- Klein cosmology. *Phys. Lett. B*, **137**, 155.

References

- Sahni, V. and Starobinsky, A. (2000). The case for a positive cosmological Λ term. *Int. J. Mod. Phys. D*, **9**, 373.
- Sahoo, P., Mishra, B., and Reddy, G.C. (2014). Axially symmetric cosmological model in $f(R, T)$ gravity. *Eur. Phys. J. Plus*, **129**, 49.
- Samanta, G.C. (2013). Kantowski-sachs universe filled with perfect fluid in $f(R, T)$ theory of gravity,. *Int.J. Theor. Phys.*, **52**, 2647.
- Schutzhold, R. (2002a). Small cosmological constant from the QCD trace anomaly. *Phys. Rev. Lett.*, **89**, 081302.
- Schutzhold, R. (2002b). On the cosmological constant and the cosmic coincidence problem. *Int. J. Mod. Phys. A*, **17**, 4359.
- Schwarzchild, G. (1916). *Ber. Ber*, 186.
- Sen, D.K. (1957). Astatic cosmological model. *Z.furr. Phys.*, **149**, 311.
- Shafi, Q. and Wetterich, C. (1984). Cosmology from higher dimensional gravity. *Phys. Lett.B*, **129**, 384.
- Shamir, M.F. (2010). Some Bianchi type cosmological models in $f(R)$ gravi. *Astrophys. Space Sci.*, **330**, 183.
- Shamir, M.F. and Jhangeer, A. (2013). A note on plane symmetric solutions in $f(R)$ gravity. *Int.J.Theor. Phys.*, **52**, 1326.
- Shamir, M.F., Jhangeer, A., and Bhatt, A. (2012). Conserved quantities in $f(R)$ gravity via noether symmetry. *Chin. Phys. Lett.*, **29**, 080402.
- Shanti, K. and Rao, V.U.M. (1991). Bianchi type-II and III models in self-creation cosmology. *Astrophy. Space Sci.*, **179**, 147.

References

- Sharif, M. and Zubair, M. (2012a). Thermodynamics in $f(R, T)$ theory of gravity. *J. Cosmol. Astropart. Phys.*, **03**, 028.
- Sharif, M. and Zubair, M. (2012b). Bianchi type I and V $f(R, T)$ gravity theory. *arXiv*, 12040848.
- Sharma, N.K. and Singh, J.K. (2014). Bianchi type-II dark energy model in $f(R, T)$ gravity. *Int J Theor Phys.*, **53**, 1424.
- Sheykhi, M.S.R.A. (2010). Thermodynamics of viscous dark energy in an RSII braneworld. *Int. J. Mod. Phys. D*, **19**, 171.
- Singh, C.P. and Kumar, S. (2007). Bianchi type-II inflationary models with constant deceleration parameter in general relativity. *Pramana J. Phys.*, **68**, 707.
- Singh, C.P. and Singh, V. (2014). Reconstruction of modified $f(R, T)$ gravity with perfect fluid cosmological models. *Gen Relativ Gravit.*, **46**, 1696.
- Singh, J.K. and .Ram, S. (1997). String cosmological models of Bianchi type III. *Astrophysics and Space Science*, **246**, 65.
- Singh, J.P. and Baghel, P.S. (2009). Bianchi type V cosmological models with constant deceleration parameter in general relativity. *Int. J. Theor. Phys.*, **48**, 449.
- Singh, T. and Singh, G.P. (1993). Lyra's geometry and cosmology. *A Review, Fortschr. Phys.*, **41**, 737.
- Singh, V. and Singh, C.P. (2013). Functional form of $f(R)$ with power-law expansion in anisotropic mode. *Astrophys. Space Sci.*, **346**, 285.
- Smoot, G. (1992). Structure in the COBE differential microwave radiometer first years maps. *Astrophys. J. Lett.*, **396**, 1.

References

- Soleng, H.H. (1987). Cosmologies based on Lyra's geometry. *Gen. Relativ. Gravit.*, **19**, 1213.
- Sotiriou, T.P. and Faraoni, V. (2010). f(R) theories of gravity. *Rev. Mod. Phys.*, **82**, 45.
- Spergel et al. (2003). First-year wilkinson microwave anisotropy probe (wmap) observations: Determination of cosmological parameters. *Astrophys. J.*, **148**, 175.
- Starobinsky, A.A. (2007). Disappearing cosmological constant inf(r) gravity. *JETP*, **86**, 157.
- Steinhardt, P.J., Wang, L., and Zlatev, I. (1999). Cosmological tracking solutions. *Phys. Rev. D*, **59**, 123504.
- Stephani, H., Kramer, K., Maccallum, K., Hoensealers, C., and Herlt, E. (2003). Exact solutions to Einstein field equation, 2nd ed.,. *Cambri. Uni. Press*.
- Syzlowski, M. and Heller, M. (1983). *Acta Phys. Polonica B*, **14**, 303.
- Tegmark, et al (2004). Cosmological parameters from sdss and wmap. *Phys.Rev. D*, **69**, 103501.
- Tikekar, R. and L. K. Patel, L. (1994). Some exact solutions in Bianchi VI0 string cosmology. *Pramana . Phys.*, **42**, 483.
- Tiwari, R.K., Tiwari, D., and Shukla, P. (2012). LRS Bianchi type-II cosmological model with a decaying Λ term. *Chin. Phys. Lett.*, **29**, 010403.
- Venkateswarlu, R. and Reddy, D.R.K. (1991). Exact Bianchi type-II, VIII, and ixcosmological models with matter and electromagnetic fields in Lyra's manifold. *Astrophy. Space Sci.*, **182**, 97.

References

- Verma, M.K. and Ram, S. (2010). Bulk viscous Bianchi type-III cosmological model with time-dependent g and Λ . *Int. J. Theor. Phys.*, **49**, 693.
- Verma, M.K., Zeyauddin, M., and Ram, S. (2011). Anisotropic cosmological model with negative constant deceleration parameter and time-decaying term Λ . *Romanian J. Phys.*, **56**, 616.
- Vishwakarma, R.G. (2003). Is the present expansion of the universe really accelerating. *Mon. Not. R. Astron. Soc.*, 345, 545.
- Wang, Y. and Tegmark, M. (2004). New dark energy constraints from supernovae, microwave background, and galaxy clustering. *Phys. Rev. Lett.*, **92**, 241301.
- Weinberg, S. (1967). A model of leptons. *Phys. Rev. Lett.*, **19**, 1264.
- Weinberg, S. (1972). Gravitation and cosmology principle and application of general theory of relativity , gravitation and cosmology. *Wiley*, **15**.
- Weinberg, S. (1989). The cosmological constant problem. *Rev. Mod. Phys.*, **1**, 61.
- Wely, H. (1918). Gravitation und elektrizitt, sitzungsberichte der kniglic preussiche akademie der wissenschaften. *Z. Berlin*, **465**, 1918.
- Wu, X. (2010). A new interpretation of zero Lyapunov exponents in BKL time for Mixmaster cosmology. *Research in Astron. Astrophys*, **10**, 211.
- Yadav, A. and Saha, B. (2012.). LRS Bianchi-I anisotropic cosmological model with dominance of dark energy. *Astrophys. Space Sci.*, **337**, 759.
- Yadav, V.K. and Yadav, L. (2011). Bianchi type-III bulk viscous and barotropic perfect fluid cosmological models in Lyras geometry. *Int. J. Theor. Phy*, **50**, 1382.