

Summary

The thesis entitled “**Functional trialkoxysilane mediated rapid synthesis of noble metal nanoparticles and their multimetallic analogues for catalytic applications**” has been divided into five chapters.

- General introduction
- Trialkoxysilane-functionalized monometallic, bimetallic, and trimetallic noble metal nanoparticles mediated non-enzymatic sensing of glucose by resonance Rayleigh scattering
- Functional trialkoxysilane mediated controlled synthesis of fluorescent gold nanoparticles and fluoremetric sensing of dopamine
- Trialkoxysilane-functionalized synthesis of mesoporous supported palladium-nickel nanocatalyst for selective hydrazine decomposition and sensing
- Synthetic incorporation of palladium-nickel bimetallic nanoparticles within mesoporous silica/silica nanoparticles as efficient and cheaper catalyst for both cationic and anionic dyes degradation

Chapter 1

The first chapter gives a brief overview of noble metal nanoparticles and a discussion of the conventional methods for noble metal nanoparticle synthesis. The various sizes and shapes of noble metal nanoparticles, as well as their properties and applications, have been investigated in this chapter. Furthermore, the analytical behaviour of these noble metal nanoparticles in terms of their physical and chemical properties has been extensively studied. The significance of multimetallic analogues over monometallic noble metal nanoparticles has been discussed briefly. Throughout the chapter, the justification for utilizing trialkoxysilane as a nanomaterial stabiliser is explored. The introductory chapter also discusses the properties of trialkoxysilane, specifically 3-APTMS, which make it a useful material for functionalization.

Chapter 2

Chapter 2 addresses the microwave-assisted controlled synthesis of noble metal nanoparticles using 3-APTMS and 3-GPTMS as reducing, stabilising, and structure-directing agents that influence the geometry and crystallinity of noble metal nanoparticles. By changing the amount of functional trialkoxysilane, the particle size of noble metal nanoparticles was regulated. Under controlled concentrations of trialkoxysilane, large nanostructures are transformed into tiny spherical particles. This chapter discusses the synthesis of monometallic nanoparticles such as AuNPs, AgNPs, and PdNPs in a homogeneous solution, as well as bimetallic nanoparticles such as Au-Ag, Au-Pd, and Ag-Pd, and trimetallic nanoparticles such as Au-Ag-Pd. The presence of different sizes of noble metal nanoparticles was revealed by TEM microscopy, which supported the size of synthesised noble metal nanoparticles. XRD measurements revealed the crystalline nature of noble metal nanoparticles. Noble metal nanoparticles have the potential to be used in fluorescence-based applications. This chapter used several noble metal nanoparticles to present a synchronous fluorescence-based glucose sensor. When compared to other noble metal nanoparticles, synchronous fluorescence intensity data showed that gold and palladium (metal ratio of 80:20) have enhanced catalytic non-enzymatic sensing of glucose. The specific changes in the intensity of synchronous fluorescence emission for various noble metal nanoparticles were also investigated.

Chapter 3

Chapter 3 describes the successful synthesis of fluorescent gold nanoparticles were successfully synthesized using a controlled concentration of 3-APTMS and 3-GPTMS. The size of the synthesized fluorescent gold nanoparticles was confirmed by TEM microscopy, which revealed the average particle size was 6 nm. EDX analysis confirmed the prepared

fluorescent gold nanoparticles were composed of Au and Si. In the XRD spectrum, plane (111) has shown high intensity and some other planes were observed with low intensity, confirmed the crystalline nature of fluorescent gold nanoparticles. The chemical bonding present in NMNPs, their elemental state, and composition were further determined by the XPS analysis. Furthermore, the optical properties of synthesized fluorescent gold nanoparticles were analyzed which exhibited blue fluorescence at the excitation wavelength of 365 nm under UV light along with a CIE co-ordinate index (0.16, 0.28). Moreover, the as-prepared fluorescent gold nanoparticles demonstrated QY up to 13% using quinine sulfate as a reference. Application of synthesized fluorescent gold nanoparticles for DA detection with excellent sensitivity and selectivity for clinical applications.

Chapter 4

Functional trialkoxysilane containing an amino-glycidoxy moiety was used to synthesize the pair of monometallic (AuNPs-1 and AuNPs-2) and bimetallic (Pd-Ni NPs-1 and Pd-Ni NPs-2) nanomaterials. Monometallic and bimetallic nanomaterials have been demonstrated to be effective for sensing and decompose hydrous hydrazine, respectively. A concentration-dependent enhancement in resonance Rayleigh scattering is enabled by these materials, which might lead to the detection of hydrazine hydrous in water. The size of the synthesized gold nanoparticles (AuNPs-1 and AuNPs-2) was measured by TEM microscopy, which revealed the average particle size was 9 nm and 17 nm, respectively. Further, it was tested for hydrous hydrazine recognition using synchronous fluorescence spectroscopy, resulting in AuNPs-1 having greater catalytic efficiency for detection of hydrous hydrazine than AuNPs-2. Furthermore, Pd-Ni NPs-1 and Pd-Ni NPs-2 have been used to decompose hydrous hydrazine. A study conducted by EDX has shown that the Pd-Ni NPs 1 and 2 had a 1:1 and 1:5 ratio of

Pd to Ni metal, respectively. Therefore, the Pd-Ni NPs-1 exhibit a higher decomposition rate for hydrous hydrazine, which could be explained by their surface area.

Chapter 5

Chapter 5 is the study of palladium and palladium-nickel bimetallic nanocrystallite-supported mesoporous silica/mesoporous silica nanoparticles (MSPs/MSNPs) synthesized at a fixed ratio of Pd/Ni by functional trialkoxysilane-mediated reduction of palladium cations. Bimetallic Pd-Ni NPs-1 and Pd-Ni NPs-2 nanocrystallites with metal ratios of 1:1 and 1:5 are stabilised and assembled using the functional trialkoxysilane as a template. A TEM microscope was used to measure the particle size of the synthesized Pd and Pd-Ni NPs-1. The average particle size was 6 nm and 9 nm, respectively. The elemental mapping of the synthesised Pd and Pd-Ni nanocrystallite confirmed the presence of respective elements in the nanocrystallite, and EDX analysis confirmed the composition of Pd, Si, and Pd, Ni, Si, respectively. An XRD spectrum confirmed the crystalline nature of Pd and Pd-Ni nanocrystallite. According to an XRD spectrum, Pd and Pd-Ni nanocrystallite are crystalline. Both cationic and anionic dyes, such as Congo red and Rh B, have been degraded using these nanocrystallite supported MSPs/MSNPs. The degradation of 15 ppm Rh B within 50 s as compared to that of mesoporous silica within 110 s under similar conditions is justified by the nanocrystallite inserted MSNPs of diameter 200 nm, with a rate constant to the order of $6.9 \times 10^{-2} \text{ s}^{-1}$, as opposed to that for MSPs of 50 m to the order of $3.2 \times 10^{-2} \text{ s}^{-1}$. An analogous observation has been made for the degradation of (Rh B). The heterogeneously supported nanocrystallite continues to exhibit 100% catalytic activity even after five further applications.

Future projection

The potential role of trialkoxysilane-based noble metal nanoparticles has been investigated in this Ph.D. thesis. In order to analyze the flexibility of the fabricated materials, we carried out detailed investigations involving many different aspects in order to analyse the capabilities of the system. Organically functionalized alkoxy silanes (APTM_s, GPTMS) were used as stabilisers, reducing agents, and secondary capping agents for the homogenous and heterogeneous phase synthesis of noble metal nanoparticles, along with variable concentrations of metal precursor. Functional trialkoxysilane together with a stablizer effectively controlled the morphological properties, promoting the directional growth of nanoparticles. This allowed the use of NMNPs as template-based multifunctional nanoparticles. Further, amino groups of functionalized noble metal nanoparticles reacted actively with foreign molecules to produce a class of nanohybrids and microscale composites. These nanoparticle model systems have shown promising catalytic properties for a number of reactions, as well as improved selectivity and efficiency.

However, there are considerable challenges to adopting these techniques as reliable tools in order to assure that the real-time demands of the catalyst can be met. NMNPs functionalised with trialkoxysilane have been shown to be a good alternative to those prepared via conventional routes. However, an even better system requires more technical analysis to be implemented for significant practical purposes. To achieve high sensitivity, the next step will be to develop more catalytically efficient models of nanoparticles. In addition, studies on heterogeneous matrix based on the stabilization and enhancement of nanoparticle catalytic activity are currently underway and will be concluded in the near future. Several studies of zeolites in the context of catalytic synthesis concluded that they are highly crystalline and porous in nature, and exhibit metal ions oriented along their axes, which facilitate the

alignment of nanoscale particles. By confining functionalized nanoparticles in a mesoporous silica matrix, more nanodimensional metal particles can be accommodated while further controlling agglomeration possibilities. A heterogeneous silica matrix with cavities allows for the exposure of both mono-and multi-metallic nanoparticles or those in bulk to catalysed reactions. As a mesoporous silica-based solid matrix, nanoparticles bound within it will interact highly with the solid matrix. This interaction may lead to more application-oriented nanoparticles that are also commercially viable. The present work is being further extended by considering the advantages of immobilizing metal nanoparticles in aluminosilicate networks and mesoporous materials for their possible applications.

References

- ❖ Agusta, M. K., David, M., Nakanishi, H., & Kasai, H. (2010). Hydrazine (N₂H₄) adsorption on Ni (1 0 0)–density functional theory investigation. *Surf. Sci.* 604(3-4), 245-251.
- ❖ Ahmad, R., Griffete, N., Lamouri, A., Felidj, N., Chehimi, M. M., & Mangeney, C. (2015). Nanocomposites of gold nanoparticles@ molecularly imprinted polymers: chemistry, processing, and applications in sensors. *Chem. Mater.* 27(16), 5464-5478.
- ❖ Ahmed, K. B. R., Nagy, A. M., Brown, R. P., Zhang, Q., Malghan, S. G., & Goering, P. L. (2017). Silver nanoparticles: Significance of physicochemical properties and assay interference on the interpretation of in vitro cytotoxicity studies. *Toxicol. In Vitro*. 38, 179-192.
- ❖ Ajitha, B., Reddy, Y. A. K., Reddy, P. S., Jeon, H. J., & Ahn, C. W. (2016). Role of capping agents in controlling silver nanoparticles size, antibacterial activity and potential application as optical hydrogen peroxide sensor. *RSC adv.* 6(42), 36171-36179.
- ❖ Alberas, D. J., Kiss, J., Liu, Z. M., & White, J. M. (1992). Surface chemistry of hydrazine on Pt (111). *Surf. Sci.* 278(1-2), 51-61.
- ❖ Alex, S., & Tiwari, A. (2015). Functionalized gold nanoparticles: synthesis, properties and applications. *J. Nanosci. Nanotechnol.* 15(3), 1869-1894.
- ❖ Ali, A. M., Said, D. A., Khayyat, M., Boustimi, M., & Seoudi, R. (2020). Improving the efficiency of the organic solar cell (CuPc/C₆₀) via PEDOT: PSS as a photoconductor layer doped by silver nanoparticles. *Results. Phys.* 16, 102819-102827.
- ❖ Ali, S., Sharma, A. S., Ahmad, W., Zareef, M., Hassan, M. M., Viswadevarayalu, A., & Chen, Q. (2021). Noble metals based bimetallic and trimetallic nanoparticles: Controlled

- synthesis, antimicrobial and anticancer applications. *Crit Rev Anal Chem.* 51(5), 454-481.
- ❖ Ali, S., Sharma, A. S., Ahmad, W., Zareef, M., Hassan, M. M., Viswadevarayalu, A., ... & Chen, Q. (2021). Noble metals based bimetallic and trimetallic nanoparticles: Controlled synthesis, antimicrobial and anticancer applications. *Critical Reviews in Analytical Chemistry*, 51(5), 454-481.
- ❖ Al-Syadi, A. M., Faisal, M., Harraz, F. A., Jalalah, M., & Alsaiari, M. (2021). Immersion-plated palladium nanoparticles onto meso-porous silicon layer as novel SERS substrate for sensitive detection of imidacloprid pesticide. *Sci. Rep.* 11(1), 1-14.
- ❖ Al-Thabaiti, S. A., Aazam, E. S., Khan, Z., & Bashir, O. (2016). Aggregation of Congo red with surfactants and Ag-nanoparticles in an aqueous solution. *Spectrochim. Acta A Mol. Biomol. Spectrosc.* 156, 28-35.
- ❖ Ameen, S.; Park, D.R.; Shin, H.S. (2016). Silicon nanowires arrays for visible light driven photocatalytic degradation of rose bengal dye. *J Mater. Sci-Mater El.* 27(10), 10460–10467.
- ❖ Amendola, V., Pilot, R., Frasconi, M., Maragò, O. M., & Iatì, M. A. (2017). Surface plasmon resonance in gold nanoparticles: a review. *J. Phys. Condens. Matter.* 29(20), 203002-203040.
- ❖ Aminabad, N. S., Farshbaf, M., & Akbarzadeh, A. (2019). Recent advances of gold nanoparticles in biomedical applications: state of the art. *Cell Biochem. Biophys.* 77(2), 123-137.
- ❖ Ammam, M. (2013). Polyoxometalates: formation, structures, principal properties, main deposition methods and application in sensing. *J. Mater. Chem. A.* 1(21), 6291-6312.

- ❖ An, J., Chen, M., Hu, N., Hu, Y., Chen, R., Lyu, Y., ... & Liu, Y. (2020). Carbon dots-based dual-emission ratiometric fluorescence sensor for dopamine detection. *Spectrochim. Acta A Mol. Biomol. Spectrosc.* 243, 118804-118810.
- ❖ An, K., & Somorjai, G. A. (2012). Size and shape control of metal nanoparticles for reaction selectivity in catalysis. *ChemCatChem.* 4(10), 1512-1524.
- ❖ Anitha, K., Mohan, S. V., & Reddy, S. J. (2004). Development of acetylcholinesterase silica sol-gel immobilized biosensor—an application towards oxydemeton methyl detection. *Biosens. Bioelectron.* 20(4), 848-856.
- ❖ Antonello, S., Holm, A. H., Instuli, E., & Maran, F. (2007). Molecular electron-transfer properties of Au₃₈ clusters. *J. Am. Chem. Soc.* 129(32), 9836-9837.
- ❖ Aragay, G., & Merkoçi, A. (2012). Nanomaterials application in electrochemical detection of heavy metals. *Electrochim. Acta.* 84, 49-61.
- ❖ Ariga, K., Ahn, E., Park, M., & Kim, B. S. (2019). Layer-by-layer assembly: recent progress from layered assemblies to layered nanoarchitectonics. *Chem. Asian J.* 14(15), 2553-2566.
- ❖ Armaroli, N., & Balzani, V. (2011). The hydrogen issue. *ChemSusChem*, 4(1), 21-36.
- ❖ Aromal, S. A., & Philip, D. (2012). Green synthesis of gold nanoparticles using *Trigonella foenum-graecum* and its size-dependent catalytic activity. *Spectrochim. Acta A Mol. Biomol. Spectrosc.* 97, 1-5.
- ❖ Asefa, T. (2016). Metal-free and noble metal-free heteroatom-doped nanostructured carbons as prospective sustainable electrocatalysts. *Acc. Chem. Res.* 49(9), 1873-1883.
- ❖ Aslan, K., Lakowicz, J. R., & Geddes, C. D. (2004). Nanogold-plasmon-resonance-based glucose sensing. *Anal. Bioch.* 330(1), 145-155.

- ❖ Astruc, D., Lu, F., & Aranzaes, J. R. (2005). Nanoparticles as recyclable catalysts: the frontier between homogeneous and heterogeneous catalysis. *Angew. Chem. Int. Ed.* 44(48), 7852-7872.
- ❖ Atwan, Q. S., & Hayder, N. H. (2020). Eco-friendly synthesis of Silver nanoparticles by using green method: Improved interaction and application in vitro and in vivo. *Iraqi J. Agric. Sci.* 51, 201-216.
- ❖ Baig, N., Kammakakam, I., & Falath, W. (2021). Nanomaterials: A review of synthesis methods, properties, recent progress, and challenges. *Adv. Mater.* 2(6), 1821-1871.
- ❖ Baig, N., Kammakakam, I., & Falath, W. (2021). Nanomaterials: A review of synthesis methods, properties, recent progress, and challenges. *Adv. Mater.* 2(6), 1821-1871.
- ❖ Bailie, J., & Hutchings, G. (1999). Promotion by sulfur of gold catalysts for crotyl alcohol formation from crotonaldehyde hydrogenation. *Chem. Commun.* (21), 2151-2152.
- ❖ Balanta, A., Godard, C., & Claver, C. (2011). Pd nanoparticles for C–C coupling reactions. *Chem. Soc. Rev.* 40(10), 4973-4985.
- ❖ Banazadeh, A.; Salimi, H.; Khaleghi, M.; Shafiei-Haghghi, S. (2016). Highly efficient degradation of hazardous dyes in aqueous phase by supported palladium nanocatalyst - A green approach. *J. Environ. Chem. Eng.* 4(2), 2178–2186.
- ❖ Bano, D., Kumar, V., Singh, V. K., & Hasan, S. H. (2018). Green synthesis of fluorescent carbon quantum dots for the detection of mercury (II) and glutathione. *New J. Chem.* 42(8), 5814-5821.

- ❖ Bassareo, V., & Di Chiara, G. (1999). Differential responsiveness of dopamine transmission to food-stimuli in nucleus accumbens shell/core compartments. *Neuroscience*. 89(3), 637-641.
- ❖ Bastús, N. G., Merkoçi, F., Piella, J., & Puntes, V. (2014). Synthesis of highly monodisperse citrate-stabilized silver nanoparticles of up to 200 nm: kinetic control and catalytic properties. *Chem. Mater.* 26(9), 2836-2846.
- ❖ Bavykin, D. V. (2008). A. Zuttel, A. Borgschulte, L. Schlapbach (eds.): *Hydrogen as a Future Energy Carrier*. 1483-1483.
- ❖ Beer, P. D., & Gale, P. A. (2001). Anion recognition and sensing: the state of the art and future perspectives. *Angew. Chem. Int. Ed.* 40(3), 486-516.
- ❖ Beik, J., Khateri, M., Khosravi, Z., Kamrava, S. K., Kooranifar, S., Ghaznavi, H., & Shakeri-Zadeh, A. (2019). Gold nanoparticles in combinatorial cancer therapy strategies. *Coord. Chem. Rev.* 387, 299-324.
- ❖ Beshkar, F.; Amiri, O.; Salehi, Z. (2017). Synthesis of ZnSnO₃ nanostructures by using novel gelling agents and their application in degradation of textile dye. *Sep. Purif. Technol.* 184, 66–71.
- ❖ Bhattacharjee, S., & Bruening, M. L. (2008). Selective hydrogenation of monosubstituted alkenes by Pd nanoparticles embedded in polyelectrolyte films. *Langmuir*, 24(6), 2916-2920.
- ❖ Bijanzadeh, A. R., Vakili, M. R., & Khordad, R. (2012). A study of the surface plasmon absorption band for nanoparticles. *Phys. Sci. Int. J.* 7(12), 1943-1948.

- ❖ Bilal, M., Zhao, Y., Rasheed, T., & Iqbal, H. M. (2018). Magnetic nanoparticles as versatile carriers for enzymes immobilization: A review. *Int. J. Biol. Macromol.* 120, 2530-2544.
- ❖ Biskos, G., Vons, V., Yurteri, C. U., & Schmidt-Ott, A. (2008). Generation and sizing of particles for aerosol-based nanotechnology. *KONA Power Part. J.* 26, 13-35.
- ❖ Boettcher, S. W., Strandwitz, N. C., Schierhorn, M., Lock, N., Lonergan, M. C., & Stucky, G. D. (2007). Tunable electronic interfaces between bulk semiconductors and ligand-stabilized nanoparticle assemblies. *Nat. Mater.* 6(8), 592-596.
- ❖ Bommel, M.R. van; Berghe, I. Vanden; Wallert, A.M.; Boitelle, R.; Wouters, J. 2007. Highperformance liquid chromatography and non-destructive three-dimensional fluorescence analysis of early synthetic dyes. *J. Chromatogr. A* , 1157(1–2), 260–272.
- ❖ Brust, M., Walker, M., Bethell, D., Schiffrian, D. J., Whyman R. (1994). Synthesis of thiol- derivatised gold nanoparticles in a two-phase Liquid -Liquid system. *J. Chem. Soc. Chem. Commun.* 801-802.
- ❖ Bunker, C. E., & Smith, M. J. (2011). Nanoparticles for hydrogen generation. *J. Mater. Chem.* 21(33), 12173-12180.
- ❖ Cai, Z., & Park, S. (2020). Synthesis of Pd nanoparticle-decorated SnO₂ nanowires and determination of the optimum quantity of Pd nanoparticles for highly sensitive and selective hydrogen gas sensor. *Sens. Actuators B Chem.* 322, 28651-128665.
- ❖ Çakar, S., Çakıroğlu, B., Şen, Ş., & Özcar, M. (2022). A photo-sensitive BiVO₄@Bi₂O₃@ g-C₃N₄ sensor for the detection of dopamine. *Microchem. J.* 178, 107360-107371

- ❖ Carnovale, C., Bryant, G., Shukla, R., & Bansal, V. (2016). Size, shape and surface chemistry of nano-gold dictate its cellular interactions, uptake and toxicity. *Prog. Mater. Sci.* 83, 152-190.
- ❖ Charrier, J., Najar, A., & Pirasteh, P. (2013). Study of optical absorbance in porous silicon nanowires for photovoltaic applications. *Appl. Surf. Sci.* 283, 828-832.
- ❖ Chellasamy, G., Ankireddy, S. R., Lee, K. N., Govindaraju, S., & Yun, K. (2021). Smartphone-integrated colorimetric sensor array-based reader system and fluorometric detection of dopamine in male and female geriatric plasma by bluish-green fluorescent carbon quantum dots. *Mater. Today Bio.* 12, 100168-100181.
- ❖ Chen, S., & Murray, R. W. (1999). Electrochemical quantized capacitance charging of surface ensembles of gold nanoparticles. *J. Phys. Chem. B.* 103(45), 9996-10000.
- ❖ Chen, Y., Wang, L., Zhai, Y., Chen, H., Dou, Y., Li, J., ... & Cao, R. (2017). Pd–Ni nanoparticles supported on reduced graphene oxides as catalysts for hydrogen generation from hydrazine. *RSC adv.* 7(51), 32310-32315.
- ❖ Cheng, M. L., Tsai, B. C., & Yang, J. (2011). Silver nanoparticle-treated filter paper as a highly sensitive surface-enhanced Raman scattering (SERS) substrate for detection of tyrosine in aqueous solution. *Anal. Chim. Acta.* 708(1-2), 89-96.
- ❖ Cheng, Y., Wu, J., Guo, C., Li, X. G., Ding, B., & Li, Y. (2017). A facile water-stable MOF-based “off-on” fluorescent switch for label-free detection of dopamine in biological fluid. *J. Mater. Chem. B.* 5(13), 2524-2535.
- ❖ Chiu, T. C., & Huang, C. C. (2009). Aptamer-functionalized nano-biosensors. *Sensors*, 9(12), 10356-10388.

- ❖ Choi, J. K., Hwang, I. S., Kim, S. J., Park, J. S., Park, S. S., Jeong, U., ... & Lee, J. H. (2010). Design of selective gas sensors using electrospun Pd-doped SnO₂ hollow nanofibers. *Sensors and actuators B: chemical*, 150(1), 191-199.
- ❖ Contour, J. P., & Pannetier, G. (1972). Hydrazine decomposition over a supported iridium catalyst. *J. Catal.* 24(3), 434-445.
- ❖ Cookson, J. (2012). The preparation of palladium nanoparticles. *Platin. Met. Rev.* 56(2), 83-98.
- ❖ Corma, A., Garcia, H. I., & Llabrés i Xamena, F. X. (2010). Engineering metal organic frameworks for heterogeneous catalysis. *Chem. Rev.* 110(8), 4606-4655.
- ❖ Dai, J., & Bruening, M. L. (2002). Catalytic nanoparticles formed by reduction of metal ions in multilayered polyelectrolyte films. *Nano Lett.* 2(5), 497-501.
- ❖ Dalirirad, S., & Steckl, A. J. (2020). Lateral flow assay using aptamer-based sensing for on-site detection of dopamine in urine. *Anal. Biochem.* 596, 113637-113648.
- ❖ Darabdhara, G., Boruah, P. K., Borthakur, P., Hussain, N., Das, M. R., Ahamad, T., ... & Yamauchi, Y. (2016). Reduced graphene oxide nanosheets decorated with Au-Pd bimetallic alloy nanoparticles towards efficient photocatalytic degradation of phenolic compounds in water. *Nanoscale*, 8(15), 8276-8287.
- ❖ Daripa, S., Verma, R., Guin, D., Chakraborty, C., Awasthi, K., & Kuila, B. K. (2021). Metal-immobilized micellar aggregates of a block copolymer from a mixed solvent for a SERS-active sensing substrate and versatile dip catalysis. *Langmuir*, 37(7), 2445-2456.

- ❖ Deng, W., & Goldys, E. M. (2014). Chemical sensing with nanoparticles as optical reporters: from noble metal nanoparticles to quantum dots and upconverting nanoparticles. *Analyst*. 139(21), 5321-5334.
- ❖ Deng, W., & Goldys, E. M. (2014). Chemical sensing with nanoparticles as optical reporters: from noble metal nanoparticles to quantum dots and upconverting nanoparticles. *Analyst*, 139(21), 5321-5334.
- ❖ Deraedt, C., & Astruc, D. (2014). “Homeopathic” palladium nanoparticle catalysis of cross carbon–carbon coupling reactions. *Acc. Chem. Res.* 47(2), 494-503.
- ❖ Deshmukh, K. M., Qureshi, Z. S., Bhatte, K. D., Venkatesan, K. A., Srinivasan, T. G., Rao, P. V., & Bhanage, B. M. (2011). One-pot electrochemical synthesis of palladium nanoparticles and their application in the Suzuki reaction. *New J. Chem.* 35(12), 2747-2751.
- ❖ Dhand, C., Dwivedi, N., Loh, X. J., Ying, A. N. J., Verma, N. K., Beuerman, R. W., ... & Ramakrishna, S. (2015). Methods and strategies for the synthesis of diverse nanoparticles and their applications: a comprehensive overview. *RSC Adv.* 5(127), 105003-105037.
- ❖ Ding, J., Liu, Z., Liu, X., Liu, J., Deng, Y., Han, X., ... & Hu, W. (2019). Mesoporous decoration of freestanding palladium nanotube arrays boosts the electrocatalysis capabilities toward formic acid and formate oxidation. *Adv. Energy Mater.* 9(25), 1900955.
- ❖ Domènech, B., Muñoz, M., Muraviev, D. N., & Macanás, J. (2011). Polymer-stabilized palladium nanoparticles for catalytic membranes: ad hoc polymer fabrication. *Nanoscale Res. Lett.* 6(1), 1-5.

- ❖ Dong, J., Zhang, Z., Zheng, H., & Sun, M. (2015). Recent progress on plasmon-enhanced fluorescence. *Nanophotonics*. 4(4), 472-490.
- ❖ Doria, G., Conde, J., Veigas, B., Giestas, L., Almeida, C., Assunção, M., ... & Baptista, P. V. (2012). Noble metal nanoparticles for biosensing applications. *Sensors*. 12(2), 1657-1687.
- ❖ Dos Santos, C. A., Seckler, M. M., Ingle, A. P., Gupta, I., Galdiero, S., Galdiero, M., ... & Rai, M. (2014). Silver nanoparticles: therapeutical uses, toxicity, and safety issues. *J. Pharm. Sci.* 103(7), 1931-1944.
- ❖ Du, H., Kong, R. M., Guo, X., Qu, F., & Li, J. (2018). Recent progress in transition metal phosphides with enhanced electrocatalysis for hydrogen evolution. *Nanoscale*. 10(46), 21617-21624.
- ❖ Du, Y., Su, J., Luo, W., & Cheng, G. (2015). Graphene-supported nickel–platinum nanoparticles as efficient catalyst for hydrogen generation from hydrous hydrazine at room temperature. *ACS Appl. Mater. Inte.* 7(2), 1031-1034.
- ❖ Du, Y., Su, J., Luo, W., & Cheng, G. (2015). Graphene-supported nickel–platinum nanoparticles as efficient catalyst for hydrogen generation from hydrous hydrazine at room temperature. *ACS Appl. Mater. Interfaces* 7(2), 1031-1034.
- ❖ Eberle, U., Felderhoff, M., & Schueth, F. (2009). Chemical and physical solutions for hydrogen storage. *Angew. Chem., Int. Ed.* 48(36), 6608-6630.
- ❖ El Kurdi, R., & Patra, D. (2018). Tuning the surface of Au nanoparticles using poly (ethylene glycol)-block-poly (propylene glycol)-block-poly (ethylene glycol): enzyme free and label free sugar sensing in serum samples using resonance Rayleigh scattering spectroscopy. *Phy. Chem. Chem. Phy.* 20(14), 9616-9629.

- ❖ El Kurdi, R., & Patra, D. (2018). Tuning the surface of Au nanoparticles using poly (ethylene glycol)-block-poly (propylene glycol)-block-poly (ethylene glycol): enzyme free and label free sugar sensing in serum samples using resonance Rayleigh scattering spectroscopy. *Phys. Chem.Chem. Phys.* 20(14), 9616-9629.
- ❖ El-Brolossy, T. A., Abdallah, T., Mohamed, M. B., Abdallah, S., Easawi, K., Negm, S., & Talaat, H. (2008). Shape and size dependence of the surface plasmon resonance of gold nanoparticles studied by Photoacoustic technique. *Eur. Phys. J. Spec. Top.* 153(1), 361-364.
- ❖ Eustis, S., & El-Sayed, M. A. (2006). Why gold nanoparticles are more precious than pretty gold: noble metal surface plasmon resonance and its enhancement of the radiative and nonradiative properties of nanocrystals of different shapes. *Chem. Soc. Rev.* 35(3), 209-217.
- ❖ Fahmy, H. M., Mosleh, A. M., Abd Elghany, A., Shams-Eldin, E., Serea, E. S. A., Ali, S. A., & Shalan, A. E. (2019). Coated silver nanoparticles: Synthesis, cytotoxicity, and optical properties. *RSC advances.* 9(35), 20118-20136.
- ❖ Faraday, M. (1857). The Bakerian lecture: Experimental relations of gold (and other metals) to Light *Philos. Trans. R. Soc. London.* 147,145-181.
- ❖ Farjami, E., Campos, R., Nielsen, J. S., Gothelf, K. V., Kjems, J., & Ferapontova, E. E. (2013). RNA aptamer-based electrochemical biosensor for selective and label-free analysis of dopamine. *Anal. Chemi.* 85(1), 121-128.
- ❖ Farzin, L., Shamsipur, M., Samandari, L., Sadjadi, S., & Sheibani, S. (2020). Biosensing strategies based on organic-scaffolded metal nanoclusters for ultrasensitive detection of tumor markers. *Talanta.* 214, 120886.

- ❖ Fei, Z., Geldbach, T. J., Zhao, D., & Dyson, P. J. (2006). From dysfunction to bis-function: on the design and applications of functionalised ionic liquids. *Chemi. Eur. J.* 12(8), 2122-2130.
- ❖ Feng, H., Elam, J. W., Libera, J. A., Setthapun, W., & Stair, P. C. (2010). Palladium catalysts synthesized by atomic layer deposition for methanol decomposition. *Chem. Mater.* 22(10), 3133-3142.
- ❖ Feng, X., Ouyang, M., Liu, X., Lu, L., Xia, Y., & He, X. (2018). Thermal runaway mechanism of lithium ion battery for electric vehicles. A review. *Energy. Stor. Mater.* 10, 246-267.
- ❖ Fihri, A., Len, C., Varma, R. S., & Solhy, A. (2017). Hydroxyapatite: A review of syntheses, structure and applications in heterogeneous catalysis. *Coord. Chem. Rev.* 347, 48-76.
- ❖ François, A., Laroche, A., Pinaud, N., Salmon, L., Ruiz, J., Robert, J., & Astruc, D. (2011). Encapsulation of docetaxel into PEGylated gold nanoparticles for vectorization to cancer cells. *Chem. Med. Chem.* 6(11), 2003-2008.
- ❖ Frey, G. D., Schütz, J., Herdtweck, E., & Herrmann, W. A. (2005). Synthesis and characterization of N-heterocyclic carbene phospha-palladacycles and their properties in heck catalysis. *Organometallics.* 24(18), 4416-4426.
- ❖ Fu, Y., Zeng, G., Lai, C., Huang, D., Qin, L., Yi, H., ... & Pi, Z. (2020). Hybrid architectures based on noble metals and carbon-based dots nanomaterials: A review of recent progress in synthesis and applications. *Chem. Eng. Trans.* 399, 125743.
- ❖ Gawande, M. B., Monga, Y., Zboril, R., & Sharma, R. K. (2015). Silica-decorated magnetic nanocomposites for catalytic applications. *Coord. Chem. Rev.* 288, 118-143.

- ❖ Ge, S., Zhang, Y., Zhang, L., Liang, L., Liu, H., Yan, M., ... & Yu, J. (2015). Ultrasensitive electrochemical cancer cells sensor based on trimetallic dendritic Au@PtPd nanoparticles for signal amplification on lab-on-paper device. *Sens. Actuators B Chem.* 220, 665-672.
- ❖ Gebre, S. H., & Sendeku, M. G. (2022). Trimetallic nanostructures and their applications in electrocatalytic energy conversions. *J. Energy Chem.* 65, 329-351.
- ❖ Ghasemi, F., Hormozi-Nezhad, M. R., & Mahmudi, M. (2015). A colorimetric sensor array for detection and discrimination of biothiols based on aggregation of gold nanoparticles. *Anal. Chim. Acta.* 882, 58-67.
- ❖ Ghosh, S. K., & Pal, T. (2007). Interparticle coupling effect on the surface plasmon resonance of gold nanoparticles: from theory to applications. *Chem. Rev* 107(11), 4797-4862.
- ❖ Gomez, S., Philippot, K., Colliere, V., Chaudret, B., Senocq, F., Lecante, Gold Nanoparticles from self-assembled Gold (1) Amine Precursors. *Chem. Commun.* 2000, 1945-1946.
- ❖ Govindaraju, S., Ankireddy, S. R., Viswanath, B., Kim, J., & Yun, K. (2017). Fluorescent gold nanoclusters for selective detection of dopamine in cerebrospinal fluid. *Sci. Rep.* 7(1), 1-12.
- ❖ Greeley, J., Jaramillo, T. F., Bonde, J., Chorkendorff, I. B., & Nørskov, J. K. (2006). Computational high-throughput screening of electrocatalytic materials for hydrogen evolution. *Nat. mater.* 5(11), 909-913.

- ❖ Guo, H., Xing, B., & He, L. (2016). Development of a filter-based method for detecting silver nanoparticles and their heteroaggregation in aqueous environments by surface-enhanced Raman spectroscopy. *Environ. Pollut.* 211, 198-205.
- ❖ Handy, R.D., Owen, R. and Valsami-Jones, E. (2008) The ecotoxicology of nanoparticles and nanomaterials: current status, knowledge gaps, challenges, and future needs. *Ecotoxicology* . 17(5), 315-325.
- ❖ Harraz, F. A., El-Hout, S. E., Killa, H. M., & Ibrahim, I. A. (2012). Palladium nanoparticles stabilized by polyethylene glycol: Efficient, recyclable catalyst for hydrogenation of styrene and nitrobenzene. *J. Catal.* 286, 184-192.
- ❖ Haruta, M., Yamada, N., Kobayashi, T., & Iijima, S. (1989). Gold catalysts prepared by coprecipitation for low-temperature oxidation of hydrogen and of carbon monoxide. *J. Catal.* 115(2), 301-309.
- ❖ He, W., Gui, R., Jin, H., Wang, B., Bu, X., & Fu, Y. (2018). Ratiometric fluorescence and visual imaging detection of dopamine based on carbon dots/copper nanoclusters dual-emitting nanohybrids. *Talanta*, 178, 109-115.
- ❖ Hebbalalu, D., Lalley, J., Nadagouda, M. N., & Varma, R. S. (2013). Greener techniques for the synthesis of silver nanoparticles using plant extracts, enzymes, bacteria, biodegradable polymers, and microwaves. *ACS Sustain. Chem. Eng.* 1(7), 703-712.
- ❖ Henglein, A., & Giersig, M. (1999). Formation of colloidal silver nanoparticles: capping action of citrate. *J. Phys. Chem. B.* 03(44), 9533-9539.
- ❖ Hoseini Chopani, S. M., Asadi, S., & Heravi, M. M. (2020). Application of bimetallic and trimetallic nanoparticles supported on graphene as novel heterogeneous catalysts in

- the reduction of nitroarenes, homo-coupling, Suzuki-Miyaura and Sonogashira reactions. *Curr. Org. Chem.* 24(19), 2216-2234.
- ❖ Hosseini, M., Aghazadeh, M., & Ganjali, M. R. (2017). A facile one-pot synthesis of cobalt-doped magnetite/graphene nanocomposite as peroxidase mimetics in dopamine detection. *New J. Chem.* 41(21), 12678-12684.
- ❖ Huang, T., & Murray, R. W. (2002). Quenching of [Ru (bpy) 3] 2+ fluorescence by binding to Au nanoparticles. *Langmuir*, 18(18), 7077-7081.
- ❖ Huang, W., Chen, S., Liu, Y., Fu, H., & Wu, G. (2010). The controlled synthesis of stable gold nanoparticles in quaternary ammonium ionic liquids by simple heating. *Nanotechnology*. 22(2), 025602.
- ❖ Huang, Y., Zhang, X., Ringe, E., Ma, L., Zhai, X., Wang, L., & Zhang, Z. (2018). Detailed correlations between SERS enhancement and plasmon resonances in subwavelength closely spaced Au nanorod arrays. *Nanoscale*, 10(9), 4267-4275.
- ❖ Hussain, Z., Thu, H. E., Sohail, M., & Khan, S. (2019). Hybridization and functionalization with biological macromolecules synergistically improve biomedical efficacy of silver nanoparticles: Reconceptualization of in-vitro, in-vivo and clinical studies. *J. Drug. Deliv. Sci. Technol.* 54, 101169-101196.
- ❖ Indrasekara, A. D. S., Meyers, S., Shubeita, S., Feldman, L. C., Gustafsson, T., & Fabris, L. (2014). Gold nanostar substrates for SERS-based chemical sensing in the femtomolar regime. *Nanoscale*. 6(15), 8891-8899.
- ❖ Islam, M. S., Chen, L., Sisler, J., & Tam, K. C. (2018). Cellulose nanocrystal (CNC)-inorganic hybrid systems: synthesis, properties and applications. *J. Mater. Chem. B*. 6(6), 864-883.

- ❖ Itoh, H., Naka, K., & Chujo, Y. (2004). Synthesis of gold nanoparticles modified with ionic liquid based on the imidazolium cation. *J. Am. Chem. Soc.* 126(10), 3026-3027.
- ❖ Jahn, M., Patze, S., Hidi, I. J., Knipper, R., Radu, A. I., Mühlig, A., ... & Popp, J. (2016). Plasmonic nanostructures for surface enhanced spectroscopic methods. *Analyst*, 141(3), 756-793.
- ❖ Jain, I. P. (2009). Hydrogen the fuel for 21st century. *Int. j. hydrog. Energy.* 34(17), 7368-7378.
- ❖ Jain, P. K., Huang, X., El-Sayed, I. H., & El-Sayed, M. A. (2008). Noble metals on the nanoscale: optical and photothermal properties and some applications in imaging, sensing, biology, and medicine. *Acc. Chem. Res.* 41(12), 1578-1586.
- ❖ Jang, Y., Chung, J., Kim, S., Jun, S. W., Kim, B. H., Lee, D. W., ... & Hyeon, T. (2011). Simple synthesis of Pd–Fe₃O₄ heterodimer nanocrystals and their application as a magnetically recyclable catalyst for Suzuki cross-coupling reactions. *Phys. Chem. Chem. Phys.* 13(7), 2512-2516.
- ❖ Jebaranjitham, J. N., Mageshwari, C., Saravanan, R., & Mu, N. (2019). Fabrication of amine functionalized graphene oxide–AgNPs nanocomposite with improved dispersibility for reduction of 4-nitrophenol. *Compos. B. Eng.* 171, 302-309.
- ❖ Jeyaraj, M., Gurunathan, S., Qasim, M., Kang, M. H., & Kim, J. H. (2019). A comprehensive review on the synthesis, characterization, and biomedical application of platinum nanoparticles. *Nanomaterials.* 9(12), 1719.
- ❖ Jia, C. J., & Schüth, F. (2011). Colloidal metal nanoparticles as a component of designed catalyst. *Phys. Chem. Chem. Phys.* 13(7), 2457-2487.

- ❖ Jiang, H. L., Singh, S. K., Yan, J. M., Zhang, X. B., & Xu, Q. (2010). Liquid-phase chemical hydrogen storage: catalytic hydrogen generation under ambient conditions. *ChemSusChem: Chemistry & Sustainability Energy & Materials*, 3(5), 541-549.
- ❖ Jiang, Z. J., Liu, C. Y., & Sun, L. W. (2005). Catalytic properties of silver nanoparticles supported on silica spheres. *J. Phys. Chem. B*. 109(5), 1730-1735.
- ❖ Justino, C. I., Rocha-Santos, T. A., & Duarte, A. C. (2010). Review of analytical figures of merit of sensors and biosensors in clinical applications. *TrAC - Trends Anal. Chem.* 9(10), 1172-1183.
- ❖ Kamel, S., & Khattab, T. A. (2021). Recent advances in cellulose supported metal nanoparticles as green and sustainable catalysis for organic synthesis. *Cellulose*. 28(8), 4545-4574.
- ❖ Kannan, P., Maiyalagan, T., & Opallo, M. (2013). One-pot synthesis of chain-like palladium nanocubes and their enhanced electrocatalytic activity for fuel-cell applications. *Nano Energy*. 2(5), 677-687.
- ❖ Kidambi, S., & Bruening, M. L. (2005). Multilayered polyelectrolyte films containing palladium nanoparticles: synthesis, characterization, and application in selective hydrogenation. *Chem. Mater.* 17(2), 301-307.
- ❖ Kim, Y., Johnson, R. C., & Hupp, J. T. (2001). Gold nanoparticle-based sensing of “spectroscopically silent” heavy metal ions. *Nano lett.* 1(4), 165-167.
- ❖ Kondrat, S. A., Miedziak, P. J., Douthwaite, M., Brett, G. L., Davies, T. E., Morgan, D. J., ... & Hutchings, G. J. (2014). Base-Free Oxidation of Glycerol Using Titania-Supported Trimetallic Au–Pd–Pt Nanoparticles. *ChemSusChem*. 7(5), 1326-1334.

- ❖ Kulkarni, M. B., & Goel, S. (2020). Microfluidic devices for synthesizing nanomaterials—A review. *Nano express.* 1(3), 032004.
- ❖ Kumar, A., Mandal, S., Selvakannan, P. R., Pasricha, R., Mandale, A. B., & Sastry, M. (2003). Investigation into the interaction between surface-bound alkylamines and gold nanoparticles. *Langmuir.* 19(15), 6277-6282.
- ❖ Kumar, N., Chamoli, P., Misra, M., Manoj, M. K., & Sharma, A. (2022). Advanced metal and carbon nanostructures for medical, drug delivery and bio-imaging applications. *Nanoscale.* **14(11)**, 3987-4017.
- ❖ Kummer, J. T. (1986). Use of noble metals in automobile exhaust catalysts. *J. Phys. Chem.* 90(20), 4747-4752.
- ❖ Kwon, M. S., Kim, N., Park, C. M., Lee, J. S., Kang, K. Y., & Park, J. (2005). Palladium nanoparticles entrapped in aluminum hydroxide: Dual catalyst for alkene hydrogenation and aerobic alcohol oxidation. *Org. Lett.* 7(6), 1077-1079.
- ❖ Lai, Y. C., Lin, C. Y., Chung, M. R., Hung, P. Y., Horng, J. C., Chen, I. C., & Chu, L. K. (2017). Distance-dependent excited-state electron transfer from tryptophan to gold nanoparticles through polyproline helices. *J. Phys. Chem. C.* 121(9), 4882-4890.
- ❖ Lan, M., Zhao, S., Wei, X., Zhang, K., Zhang, Z., Wu, S., ... & Zhang, W. (2019). Pyrene-derivatized highly fluorescent carbon dots for the sensitive and selective determination of ferric ions and dopamine. *Dyes Pigm.* 170, 107574-107580.
- ❖ Landes, C. F., Braun, M., & El-Sayed, M. A. (2001). On the nanoparticle to molecular size transition: fluorescence quenching studies. *J. Phys. Chem. B.* 105(43), 10554-10558.

- ❖ Langer, J., Novikov, S. M., & Liz-Marzán, L. M. (2015). Sensing using plasmonic nanostructures and nanoparticles. *Nanotechnology*. 26(32), 322001-322029
- ❖ Lee, J., Shim, W., Lee, E., Noh, J. S., & Lee, W. (2011). Highly mobile palladium thin films on an elastomeric substrate: Nanogap-based hydrogen gas sensors. *Angew. Chem. Int. Ed.* 50(23), 5301-5305.
- ❖ Lee, S. H., & Jun, B. H. (2019). Silver nanoparticles: synthesis and application for nanomedicine. *Int. J. Mol. Sci.* 20(4), 865-889.
- ❖ Lewandowski, W., Wójcik, M., & Górecka, E. (2014). Metal nanoparticles with liquid-crystalline ligands: controlling nanoparticle superlattice structure and properties. *ChemPhysChem*. 15(7), 1283-1295.
- ❖ Li, B. R., Hsieh, Y. J., Chen, Y. X., Chung, Y. T., Pan, C. Y., & Chen, Y. T. (2013). An ultrasensitive nanowire-transistor biosensor for detecting dopamine release from living PC12 cells under hypoxic stimulation. *J. Am. Chem. Soc.* 135(43), 16034-16037.
- ❖ Li, D., Dong, M., Besenbacher, F., Huang, Y., & Chen, M. (2016). The preparation of a recyclable catalyst of silver nanoparticles dispersed in a mesoporous silica nanofiber matrix. *RSC advances*, 6(70), 65613-65618.
- ❖ Combes Li, J., Zhao, T., Chen, T., Liu, Y., Ong, C. N., & Xie, J. (2015). Engineering noble metal nanomaterials for environmental applications. *Nanoscale*, 7(17), 7502-7519.
- ❖ Li, L., Zhao, H., Wang, J., & Wang, R. (2014). Facile fabrication of ultrafine palladium nanoparticles with size-and location-control in click-based porous organic polymers. *ACS nano*, 8(5), 5352-5364.

- ❖ Li, X.; Wan, T.; Qiu, J.; Wei, H.; Qin, F.; Wang, Y.; Liao, Y.; Huang, Z.; Tan, X. 2017. In-situ photocalorimetry-fluorescence spectroscopy studies of RH B photocatalysis over Z-scheme gC₃N₄@Ag@Ag₃PO₄ nanocomposites: A pseudo-zero-order rather than a first-order process. *Appl. Catal. B-Environ.* 217, 591–602.
- ❖ Li, Z., Li, X., Jian, M., Geleta, G. S., & Wang, Z. (2019). Two-dimensional layered nanomaterial-based electrochemical biosensors for detecting microbial toxins. *Toxin.* 12(1), 20-43.
- ❖ Liao, G., Fang, J., Li, Q., Li, S., Xu, Z., & Fang, B. (2019). Ag-based nanocomposites: synthesis and applications in catalysis. *Nanoscale.* 11(15), 7062-7096.
- ❖ Liguori, F., & Barbaro, P. (2014). Green semi-hydrogenation of alkynes by Pd@ borate monolith catalysts under continuous flow. *J. Catal.* 311, 212-220.
- ❖ Link, S., & El-Sayed, M. A. (2003). Optical properties and ultrafast dynamics of metallic nanocrystals. *Annu. Rev. Phys. Chem.* 54(1), 331-366.
- ❖ Liu, C. J., Burghaus, U., Besenbacher, F., & Wang, Z. L. (2010). Preparation and characterization of nanomaterials for sustainable energy production. *ACS Nano.* 4(10), 5517-5526.
- ❖ Liu, J., & Lu, Y. (2006). Preparation of aptamer-linked gold nanoparticle purple aggregates for colorimetric sensing of analytes. *Natu.protoc.* 1(1), 246-252.
- ❖ Liu, K., Qu, S., Zhang, X., Tan, F., & Wang, Z. (2013). Improved photovoltaic performance of silicon nanowire/organic hybrid solar cells by incorporating silver nanoparticles. *Nanoscale Res. Lett.* 8(1), 1-6.

- ❖ Liu, N., Tang, M. L., Hentschel, M., Giessen, H., & Alivisatos, A. P. (2011). Nanoantenna-enhanced gas sensing in a single tailored nanofocus. *Nat. Mater.* 10(8), 631-636.
- ❖ Liu, W., Herrmann, A. K., Bigall, N. C., Rodriguez, P., Wen, D., Oezaslan, M., ... & Eychmüller, A. (2015). Noble Metal Aerogels Synthesis, Characterization, and Application as Electrocatalysts. *Acc. Chem. Res.* 48(2), 154-162.
- ❖ Loza, K., Heggen, M., & Epple, M. (2020). Synthesis, structure, properties, and applications of bimetallic nanoparticles of noble metals. *Adv. Funct. Mater.* 30(21), 1909260.
- ❖ Lu, L. M., Li, H. B., Qu, F., Zhang, X. B., Shen, G. L., & Yu, R. Q. (2011). In situ synthesis of palladium nanoparticle-graphene nanohybrids and their application in nonenzymatic glucose biosensors. *Biosens. Bioelectron.* 26(8), 3500-3504.
- ❖ Luo, J., Li, S., Pang, C., Wang, M., Ma, X., & Zhang, C. (2022). Highly selective fluorescence probe for imidacloprid measurement based on fluorescence resonance energy transfer. *Microchem. J.* 107172-107178.
- ❖ Luo, S., Zeng, Z., Zeng, G., Liu, Z., Xiao, R., Chen, M., ... & Jiang, D. (2019). Metal organic frameworks as robust host of palladium nanoparticles in heterogeneous catalysis: Synthesis, application, and prospect. *ACS Appl. Mater. Interfaces.* 11(36), 32579-32598.
- ❖ Ma, W., Ma, H., Yang, Z. Y., & Long, Y. T. (2018). Single Ag nanoparticle electro-oxidation: potential-dependent current traces and potential-independent electron transfer kinetic. *J. Phys. Chem. Lett.* 9(6), 1429-1433.

- ❖ Mak, K. F., He, K., Shan, J., & Heinz, T. F. (2012). Control of valley polarization in monolayer MoS₂ by optical helicity. *Nat. Nanotechnol.* 7(8), 494-498.
- ❖ Malekzad, H., Zangabad, P. S., Mirshekari, H., Karimi, M., & Hamblin, M. R. (2017). Noble metal nanoparticles in biosensors: recent studies and applications. *Nanotechnol. Rev.* 6(3), 301-329.
- ❖ Malik, A., & Nath, M. (2020). Synthesis of Ag/ZIF-7 by immobilization of Ag nanoparticles onto ZIF-7 microcrystals: A heterogeneous catalyst for the reduction of nitroaromatic compounds and organic dyes. *J. Environ. Chem. Eng.* 8(6), 104547-104562.
- ❖ Mallik, K., Mandal, M., Pradhan, N., & Pal, T. (2001). Seed mediated formation of bimetallic nanoparticles by UV irradiation: a photochemical approach for the preparation of “core– shell” type structures. *Nano Lett.* 1(6), 319-322.
- ❖ Mallin, M. P., & Murphy, C. J. (2002). Solution-phase synthesis of sub-10 nm Au–Ag alloy nanoparticles. *Nano Lett.* 2(11), 1235-1237.
- ❖ Manukyan, K. V., Cross, A., Rouvimov, S., Miller, J., Mukasyan, A. S., & Wolf, E. E. (2014). Low temperature decomposition of hydrous hydrazine over FeNi/Cu nanoparticles. *Appl. Catal. A.* 476, 47-53.
- ❖ Martin, C. R. (1994). Nanomaterials: a membrane-based synthetic approach. *Science.* 266(5193), 1961-1966.
- ❖ Martínez-Castañon, G. A., Nino-Martinez, N., Martinez-Gutierrez, F., Martinez-Mendoza, J. R., & Ruiz, F. (2008). Synthesis and antibacterial activity of silver nanoparticles with different sizes. *J. Nanoparticle Res.* 10(8), 1343-1348.

- ❖ Matouq, M.; Al-Anber, Z.; Susumu, N.; Tagawa, T.; Karapanagioti, H. (2014). The kinetic of dyesdegradation resulted from food industry in wastewater using high frequency of ultrasound. *Sep. Purif. Technol.* 135, 42–47.
- ❖ Mauter, M. S., &Elimelech, M. (2008). Environmental applications of carbon-based nanomaterials. *Environ. Sci. Technol.* 42(16), 5843-5859.
- ❖ Meng, Y., Lai, Y., Jiang, X., Zhao, Q., & Zhan, J. (2013). Silver nanoparticles decorated filter paper via self-sacrificing reduction for membrane extraction surface-enhanced Raman spectroscopy detection. *Analyst.* 138(7), 2090-2095.
- ❖ Mijatovic, D., Eijkel, J. C., & Van den Berg, A. (2005). Technologies for nanofluidic systems: top-down vs. bottom-up—a review. *Lab Chip.* 5(5), 492-500.
- ❖ Minami, T., Kaneko, K., Nagasaki, T., & Kubo, Y. (2008). Isothiouronium-based amphiphilic gold nanoparticles with a colorimetric response to hydrophobic anions in water: a new strategy for fluoride ion detection in the presence of a phenylboronic acid. *Tetrahedron Lett.* 49(3), 432-436.
- ❖ Minh, P. N., Hoang, V. T., Dinh, N. X., Van Hoang, O., Van Cuong, N., Hop, D. T. B., ... & Le, A. T. (2020). Reduced graphene oxide-wrapped silver nanoparticles for applications in ultrasensitive colorimetric detection of Cr (VI) ions and the carbaryl pesticide. *New J. Chem.* 44(18), 7611-7620.
- ❖ Mohaghegh, S., Osouli-Bostanabad, K., Nazemiyeh, H., Javadzadeh, Y., Parvizpur, A., Barzegar-Jalali, M., & Adibkia, K. (2020). A comparative study of eco-friendly silver nanoparticles synthesis using *Prunus domestica* plum extract and sodium citrate as reducing agents. *Adv. Powder Technol.* 31(3), 1169-1180.

- ❖ Mohamed, A.; Ghobara, M.M.; Abdelmaksoud, M.K.; Mohamed, G.G. (2019). A novel and highly efficient photocatalytic degradation of malachite green dye via surface modified polyacrylonitrile nanofibers/biogenic silica composite nanofibers. *Sep. Purif. Technol.* 210, 935–942.
- ❖ Mohamed, M. B., Volkov, V., Link, S., & El-Sayed, M. A. (2000). The lightning'gold nanorods: fluorescence enhancement of over a million compared to the gold metal. *Chem. Phys. Lett.* 317(6), 517-523.
- ❖ Moumen, A., Halim, W., Jaffal, S., Abderrafi, K., Eddahbi, A., Sebti, S., & Ouaskit, S. (2017). Microwave assisted synthesis of palladium nanoparticles in an aqueous emulsion of copolymer: application to catalysis. *J. Clust. Sci.* 28(5), 2817-2832.
- ❖ Mubeen, S., Zhang, T., Yoo, B., Deshusses, M. A., & Myung, N. V. (2007). Palladium nanoparticles decorated single-walled carbon nanotube hydrogen sensor. *J. Phys. Chem. C.* 111(17), 6321-6327.
- ❖ Navalon, S., Dhakshinamoorthy, A., Alvaro, M., & Garcia, H. (2016). Metal nanoparticles supported on two-dimensional graphenes as heterogeneous catalysts. *Coord. Chem. Rev.* 312, 99-148.
- ❖ Ndolomingo, M. J., Bingwa, N., & Meijboom, R. (2020). Review of supported metal nanoparticles: synthesis methodologies, advantages and application as catalysts. *J. Mater. Sci.* 55(15), 6195-6241.
- ❖ Nguyen, D.C.T.; Oh, W.C. (2018). Ternary self-assembly method of mesoporous silica and Cu₂O combined graphene composite by nonionic surfactant and photocatalytic degradation of cationic anionic dye pollutants. *Sep. Purif. Technol.* (190), 77–89.

- ❖ Nishan, U., Sabba, U., Rahim, A., Asad, M., Shah, M., Iqbal, A., & Muhammad, N. (2021). Ionic liquid tuned titanium dioxide nanostructures as an efficient colorimetric sensing platform for dopamine detection. *Mater. Chemi. Phys.* 262, 124289-124397.
- ❖ Nuchprayoon, K., & Phuaksaman, C. (2018). Evaluation of Key Success Factors in Project Management of Information Systems and Selection of Operators using Analytical Hierarchy Process. *Int. J. Appl. Eng. Res.* 13(7), 5515-5521.
- ❖ Ogilby, P. R. (2010). Singlet oxygen: there is indeed something new under the sun. *Chem. Soc. Rev.* 39(8), 3181-3209.
- ❖ Oliaee, S. N., Zhang, C., Hwang, S. Y., Cheung, H. M., & Peng, Z. (2016). Hydrogen production via hydrazine decomposition on model platinum–nickel nanocatalyst with a single (111) facet. *J. Phys. Chem. C.* 120(18), 9764-9772.
- ❖ Orimo, S. I., Nakamori, Y., Eliseo, J. R., Züttel, A., & Jensen, C. M. (2007). Complex hydrides for hydrogen storage. *Chem. Rev.* 107(10), 4111-4132.
- ❖ P C Pandey and Govind Pandey, (2016) Synthesis of gold nanoparticles resistant to pH and salt for biomedical applications; functional activity of organic amine, *J. Mater. Res.* **31**, 3313-3323.
- ❖ P C Pandey and Govind Pandey, (2018a) 3-Aminopropyltrimethoxysilane mediated controlled synthesis of functional noble metal nanoparticles and its multi-metallic analogues in the presence of small organic reducing agents for selective application, *MRS. Adv.* 3, 789-801.
- ❖ P C Pandey and Shubhangi Shukla, (2017). Mesoporous silica beads encapsulated with functionalized palladium nanocrystallites: Novel catalyst for selective hydrogen evolution, *J. Mater. Res.* **32**, 3574-3584.

- ❖ Pacardo, D. B., Ardman, E., & Knecht, M. R. (2014). Effects of substrate molecular structure on the catalytic activity of peptide-templated Pd nanomaterials. *J. Phys. Chem. C.* 118(5), 2518-2527.
- ❖ Pandey, P. C., & Chauhan, D. S. (2012). 3-Glycidoxypropyltrimethoxysilane mediated in situ synthesis of noble metal nanoparticles: Application to hydrogen peroxide sensing. *Analyst*, 137(2), 376-385.
- ❖ Pandey, P. C., & Panday, D. (2016a). Tetrahydrofuran and hydrogen peroxide mediated conversion of potassium hexacyanoferrate into Prussian blue nanoparticles: Application to hydrogen peroxide sensing. *Electrochim. Acta*, 190, 758-765.
- ❖ Pandey, P. C., & Pandey, G. (2014c). Tunable functionality and nanogeometry in tetrahydrofuran hydroperoxide and 3-aminopropyl-trimethoxysilane mediated synthesis of gold nanoparticles; functional application in glutathione sensing. *J. Mater. Chem. B.* 2(21), 3383-3390.
- ❖ Pandey, P. C., & Pandey, G. (2014c). Tunable functionality and nanogeometry in tetrahydrofuran hydroperoxide and 3-aminopropyl-trimethoxysilane mediated synthesis of gold nanoparticles; functional application in glutathione sensing. *J. Mater. Chem B.* 2(21), 3383-3390.
- ❖ Pandey, P. C., & Pandey, G. (2016b). Synthesis of gold nanoparticles resistant to pH and salt for biomedical applications; functional activity of organic amine. *J. Mater. Res.* 31(21), 3313-3323.
- ❖ Pandey, P. C., & Pandey, G. (2016c). One-pot two-step rapid synthesis of 3-aminopropyltrimethoxysilane-mediated highly catalytic Ag@(PdAu) trimetallic nanoparticles. *Catal. Sci. Technol.* 6(11), 3911-3917.

- ❖ Pandey, P. C., & Prakash, R. (1998). Electrochemical synthesis of polyindole and its evaluation for rechargeable battery applications. *Journal of the Electrochemical Society*, 145(3), 999.
- ❖ Pandey, P. C., & Shukla, S. (2018b). Solvent dependent fabrication of bifunctional nanoparticles and nanostructured thin films by self-assembly of organosilanes. *J. Sol-Gel Sci. Technol.* 86(3), 650-663.
- ❖ Pandey, P. C., & Singh, R. (2015a). Controlled synthesis of functional silver nanoparticles dispersible in aqueous and non-aqueous medium. *J. Nanosci. Nanotechnol.* 15(8), 5749-5759.
- ❖ Pandey, P. C., & Singh, R. (2015b). Controlled synthesis of Pd and Pd–Au nanoparticles: effect of organic amine and silanol groups on morphology and polycrystallinity of nanomaterials. *RSC. Adv.* 5(15), 10964-10973.
- ❖ Pandey, P. C., & Upadhyay, B. C. (2005). Studies on differential sensing of dopamine at the surface of chemically sensitized ormosil-modified electrodes. *Talanta*. 67(5), 997-1006.
- ❖ Pandey, P. C., Chauhan, D. S., & Singh, V. (2009). Poly (indole-6-carboxylic acid) and tetracyanoquinodimethane-modified electrode for selective oxidation of dopamine. *Electrochim. Acta*. 54(8), 2266-2270.
- ❖ Pandey, P. C., Chauhan, D. S., & Singh, V. (2012b). Effect of processable polyindole and nanostructured domain on the selective sensing of dopamine. *Mater. Sci. Eng. C*. 32(1), 1-11.

- ❖ Pandey, P. C., Katyal, N., Pandey, G., & Narayan, R. J. (2020). Synthesis of self-assembled siloxane–polyindole–gold nanoparticle polymeric nanofluid for biomedical membranes. *MRS. Commun.* 10(3), 482-486.
- ❖ Pandey, P. C., Panday, D., & Pandey, G. (2014b). 3-Aminopropyltrimethoxysilane and organic electron donors mediated synthesis of functional amphiphilic gold nanoparticles and their bioanalytical applications. *RSC adv.* 4(105), 60563-60572.
- ❖ Pandey, P. C., Pandey, A. K., & Pandey, G. (2014a). Functionalized alkoxy silane mediated controlled synthesis of noble metal nanoparticles dispersible in aqueous and non-aqueous medium. *J. Nanosci. Nanotechnol.* 14(9), 6606-6613.
- ❖ Pandey, P. C., Pandey, G., Haider, J., & Pandey, G. (2016). Role of organic carbonyl moiety and 3-aminopropyltrimethoxysilane on the synthesis of gold nanoparticles specific to pH-and salt-tolerance. *J. Nanosci. Nanotechnol.*, 16(6), 6155-6163.
- ❖ Pandey, P. C., Shukla, S., & Pandey, Y. (2016d). 3-Aminopropyltrimethoxysilane and graphene oxide/reduced graphene oxide-induced generation of gold nanoparticles and their nanocomposites: electrocatalytic and kinetic activity. *RSC advances*, 6(84), 80549-80556.
- ❖ Pandey, P. C., Shukla, S., & Pandey, Y. (2017). Mesoporous silica beads encapsulated with functionalized palladium nanocrystallites: Novel catalyst for selective hydrogen evolution. *J. Mater. Res.* 32(18), 3574-3584.
- ❖ Pandey, P. C., Shukla, S., Pandey, G., & Narayan, R. J. (2020). Organotrialkoxysilane-mediated controlled synthesis of noble metal nanoparticles and their impact on selective fluorescence enhancement and quenching. *J. Vac. Sci. Technol. B, Nanotechnol. Microelectron.* 38(5), 052801.

- ❖ Pandey, P. C., Singh, R., & Pandey, A. K. (2014d). Tetrahydrofuran hydroperoxide and 3-Aminopropyltrimethoxysilane mediated controlled synthesis of Pd, Pd-Au, Au-Pd nanoparticles: Role of Palladium nanoparticles on the redox electrochemistry of ferrocene monocarboxylic acid. *Electrochim. Acta.* 138, 163-173.
- ❖ Pandey, P. C., Singh, R., & Pandey, A. K. (2014e). Tetrahydrofuran hydroperoxide and 3-Aminopropyltrimethoxysilane mediated controlled synthesis of Pd, Pd-Au, Au-Pd nanoparticles: Role of Palladium nanoparticles on the redox electrochemistry of ferrocene monocarboxylic acid. *Electrochim. Acta.* 38, 163-173.
- ❖ Pandey, P. C., Upadhyay, S., & Pathak, H. C. (1999). A new glucose biosensor based on sandwich configuration of organically modified sol-gel glass. *Electroanalysis*, 11(1), 59-64.
- ❖ Pandey, P. C., Upadhyay, S., & Pathak, H. C. (1999a). A new glucose biosensor based on sandwich configuration of organically modified sol-gel glass. *Electroanalysis*, 11(1), 59-64.
- ❖ Pandey, P. C., Upadhyay, S., & Sharma, S. (2003). Functionalized ormosils-based biosensor probing a horseradish peroxidase-catalyzed reaction. *J. Electrochem. Soc.* 150(4), H85-H92.
- ❖ Pandey, P. C., Upadhyay, S., & Sharma, S. (2003b). Functionalized ormosils-based biosensor: Probing a horseradish peroxidase-catalyzed reaction. *J. Electrochem. Soc.* 150(4), H85-H95
- ❖ Pandey, P. C., Upadhyay, S., Pathak, H. C., & Pandey, C. M. D. (1999b). Studies on ferrocene immobilized sol-gel glasses and its application in the construction of a novel solid-state ion sensor. *Electroanalysis*. 11(13), 950-956.

- ❖ Pandey, P. C., Upadhyay, S., Shukla, N. K., & Sharma, S. (2003a). Studies on the electrochemical performance of glucose biosensor based on ferrocene encapsulated ORMOSIL and glucose oxidase modified graphite paste electrode. *Biosens. Bioelectron.* 18(10), 1257-1268.
- ❖ Pandey, P. C., Upadhyay, S., Tiwari, I., & Sharma, S. (2001b). A Novel Ferrocene-Encapsulated Palladium-Linked Ormosil-Based Electrocatalytic Biosensor. The Role of the Reactive Functional Group. *Electroanal.* 13(18), 1519-1527.
- ❖ Pandey, P. C., Upadhyay, S., Tiwari, I., & Tripathi, V. S. (2001a). An ormosil-based peroxide biosensor—a comparative study on direct electron transport from horseradish peroxidase. *Sens. Actuators B: Chem.* 72(3), 224-232.
- ❖ Pandey, P., Pandey, G., & Narayan, R. (2021). Polyethylenimine-mediated controlled synthesis of Prussian blue-gold nanohybrids for biomedical applications. *J. Biomater. Appl.* 36(1), 26-35.
- ❖ Pandey, P.C.; Pandey, G.; Narayan, R.J. 2017. Polyethylenimine-mediated synthetic insertion of gold nanoparticles into mesoporous silica nanoparticles for drug loading and biocatalysis. *Biointerphases.* 12(1), 011005-011016.
- ❖ Pandey, P.C.; Shukla, S.; Pandey, Y. (2017). Mesoporous silica beads encapsulated with functionalized palladium nanocrystallites: Novel catalyst for selective hydrogen evolution. *J. Mater. Res.* 32(18), 3574–3584.
- ❖ Paramasivam, G., Kayambu, N., Rabel, A. M., Sundramoorthy, A. K., & Sundaramurthy, A. (2017). Anisotropic noble metal nanoparticles: Synthesis, surface functionalization and applications in biosensing, bioimaging, drug delivery and theranostics. *Acta Biomater.* 49, 45-65.

- ❖ Patra, S., Viswanath, B., Barai, K., Ravishankar, N., & Munichandraiah, N. (2010). High-surface step density on dendritic Pd leads to exceptional catalytic activity for formic acid oxidation. *ACS Appl. Mater. Interfaces.* 2(11), 2965-2969.
- ❖ Pei, Y., Maligal-Ganesh, R. V., Xiao, C., Goh, T. W., Brashler, K., Gustafson, J. A., & Huang, W. (2015). An inorganic capping strategy for the seeded growth of versatile bimetallic nanostructures. *Nanoscale*, 7(40), 16721-16728.
- ❖ Perry, M., Li, Q., & Kennedy, R. T. (2009). Review of recent advances in analytical techniques for the determination of neurotransmitters. *Anal. Chim. Acta.* 653(1), 1-22.
- ❖ Piktel, E., Niemirowicz, K., Wątek, M., Wollny, T., Deptuła, P., & Bucki, R. (2016). Recent insights in nanotechnology-based drugs and formulations designed for effective anti-cancer therapy. *J. nanobiotechnology.* 14(1), 1-23.
- ❖ Pirzada, M., & Altintas, Z. (2019). Nanomaterials for healthcare biosensing applications. *Sens.* 19(23), 5311.
- ❖ Prospisito, P., Burratti, L., & Venditti, I. (2020). Silver nanoparticles as colorimetric sensors for water pollutants. *Chemosensors.* 8(2), 26, 1-29.
- ❖ Qiao, Z., Wang, Z., Zhang, C., Yuan, S., Zhu, Y., Wang, J., & Wang, S. (2013). PVAm–PIP/PS composite membrane with high performance for CO₂/N₂ separation. *AIChE. J.* 59(1), 215-228.
- ❖ Rad, A. S., Mirabi, A., Binaian, E., & Tayebi, H. (2011). A review on glucose and hydrogen peroxide biosensor based on modified electrode included silver nanoparticles. *Int. J. Electrochem. Sci.* 6(8), 3671-3683.

- ❖ Rahim, E. H., Kamounah, F. S., Frederiksen, J., & Christensen, J. B. (2001). Heck reactions catalyzed by PAMAM-dendrimer encapsulated Pd (0) nanoparticles. *Nano Lett.* 1(9), 499-501.
- ❖ Rajeev, R., Datta, R., Varghese, A., Sudhakar, Y. N., & George, L. (2021). Recent advances in bimetallic based nanostructures: Synthesis and electrochemical sensing applications. *Microchem. J.* 163, 105910.
- ❖ Rand, D. A. J., & Dell, R. M. (2008). Hydrogen energy challenges and prospects the royal society of chemistry. Thomas Graham House, Science Park, Milton Road, Cambridge CB4 0WF, UK ISBN, 978-0.
- ❖ Raut, S., Rich, R., Fudala, R., Butler, S., Kokate, R., Gryczynski, Z., ... & Gryczynski, I. (2014). Resonance energy transfer between fluorescent BSA protected Au nanoclusters and organic fluorophores. *Nanoscale*, 6(1), 385-391.
- ❖ Reguera, J., Langer, J., de Aberasturi, D. J., & Liz-Marzán, L. M. (2017). Anisotropic metal nanoparticles for surface enhanced Raman scattering. *Chem. Soc. Rev.* 46(13), 3866-3885.
- ❖ Rodriguez, P. C., Pereira, D. B., Borgkvist, A., Wong, M. Y., Barnard, C., Sonders, M. S.,& Sulzer, D. (2013). Fluorescent dopamine tracer resolves individual dopaminergic synapses and their activity in the brain. *Proc. Natl. Acad. Sci. U.S.A.* 110(3), 870-875.
- ❖ Roushani, M., & Zalpour, N. (2022). Impedimetric ultrasensitive detection of trypsin based on hybrid aptamer-2DMIP using a glassy carbon electrode modified by nickel oxide nanoparticle. *Microchem. J.* 172, 106955-106964

- ❖ Rypar, T., Adam, V., Vaculovicova, M., & Macka, M. (2021). Paperfluidic devices with a selective molecularly imprinted polymer surface for instrumentation-free distance-based detection of protein biomarkers. *Sens. Actuators B: Chem.* 341, 129999.
- ❖ Saeed, W., Abbasi, Z., Majeed, S., Shahzad, S. A., Khan, A. F., & Shaikh, A. J. (2021). An insight into the binding behavior of graphene oxide and noble metal nanoparticles. *J. Appl. Phys.* 129(12), 125302.
- ❖ Saha, J., Begum, A., Mukherjee, A., & Kumar, S. (2017). A novel green synthesis of silver nanoparticles and their catalytic action in reduction of Methylene Blue dye. *Sustain. Environ. Res.* 27(5), 245-250.
- ❖ Saha, k., Agasti, S. S., Kim, C., Li, X., & Rotelli, V. M. (2012). Gold nanoparticles in chemical and biological sensing. *Chem. rev.* 112(5), 2739-2779.
- ❖ Saldan, I., Semenyuk, Y., Marchuk, I., & Reshetnyak, O. (2015). Chemical synthesis and application of palladium nanoparticles. *J. Mater. Sci.* 50(6), 2337-2354.
- ❖ Salem, M. A., Bakr, E. A., & El-Attar, H. G. (2018). Pt@ Ag and Pd@ Ag core/shell nanoparticles for catalytic degradation of Congo red in aqueous solution. *Spectrochim. Acta A Mol. Biomol. Spectrosc.* 188, 155-163.
- ❖ Salve, M., Mandal, A., Amreen, K., Pattnaik, P. K., & Goel, S. (2020). Greenly synthesized silver nanoparticles for supercapacitor and electrochemical sensing applications in a 3D printed microfluidic platform. *Microchem. J.* 157, 104973.
- ❖ Samaei, S.M.; Gato-Trinidad, S.; Altaee, A. (2018). The application of pressure-driven ceramic membrane technology for the treatment of industrial wastewaters – A review. *Sep. Purif. Technol.* 200, 198–220.

- ❖ Samokhvalov, A. (2020). Analysis of various solid samples by synchronous fluorescence spectroscopy and related methods: A review. *Talanta*, 216, 120944-120962.
- ❖ Sardar, R., Funston, A. M., Mulvaney, P., & Murray, R. W. (2009). Gold nanoparticles: past, present, and future. *Langmuir*. 25(24), 13840-13851.
- ❖ Sau, T. K., Rogach, A. L., Jäckel, F., Klar, T. A., & Feldmann, J. (2010). Properties and applications of colloidal nonspherical noble metal nanoparticles. *Adv. Mater.* 22(16), 1805-1825.
- ❖ Schlapbach, L., Züttel, A., Gröning, P., Gröning, O., & Aebi, P. (2001). Hydrogen for novel materials and devices. *Appl. Phys. A*. 72(2), 245-253.
- ❖ Serrano, E., Rus, G., & Garcia-Martinez, J. (2009). Nanotechnology for sustainable energy. *Renew. Sust. Energ. Rev.* 13(9), 2373-2384.
- ❖ Shahzamani, Z., Ranjbar, M., Comini, E., Goodarzi, M. T., Salamat, H., & Sberveglieri, G. (2019). Palladium thin films on microfiber filtration paper as flexible substrate and its hydrogen gas sensing mechanism. *Int. J. Hydrog. Energy*. 44(31), 17185-17194.
- ❖ Shang, L., Bian, T., Zhang, B., Zhang, D., Wu, L. Z., Tung, C. H., ... & Zhang, T. (2014). Graphene-supported ultrafine metal nanoparticles encapsulated by mesoporous silica: robust catalysts for oxidation and reduction reactions. *Angew. Chem.* 126(1), 254-258.
- ❖ Sharma, C., Ansari, S., Ansari, M. S., Satsangee, S. P., & Srivastava, M. M. (2020). Single-step green route synthesis of Au/Ag bimetallic nanoparticles using clove buds extract: Enhancement in antioxidant bio-efficacy and catalytic activity. *Mater. Sci. Eng. C*. 116, 111153-111166.

- ❖ Sharma, G., Kumar, A., Sharma, S., Naushad, M., Dwivedi, R. P., ALOthman, Z. A., & Mola, G. T. (2019). Novel development of nanoparticles to bimetallic nanoparticles and their composites: A review. *J. King Saud Univ. Sci.* 31(2), 257-269.
- ❖ Sharma, G., Kumar, D., Kumar, A., Ala'a, H., Pathania, D., Naushad, M., & Mola, G. T. (2017). Revolution from monometallic to trimetallic nanoparticle composites, various synthesis methods and their applications: A review. *Mater. Sci. Eng. C.* 71, 1216-1230.
- ❖ Shen, D., Chen, L., Yang, J., Zhang, R., Wei, Y., Li, X., ... & Zhao, D. (2015). Ultradispersed palladium nanoparticles in three-dimensional dendritic mesoporous silica nanospheres: toward active and stable heterogeneous catalysts. *ACS Appl. Mater. Interfaces.* 7(31), 17450-17459.
- ❖ Shen, Q., Li, X., Li, R., & Wu, Y. (2020). Application of metal–organic framework materials and derived porous carbon materials in catalytic hydrogenation. *ACS Sustain. Chem. Eng.* 8(48), 17608-17621.
- ❖ Shenashen, M. A., El-Safty, S. A., & Elshehy, E. A. (2014). Synthesis, morphological control, and properties of silver nanoparticles in potential applications. *Particle & Particle Systems Characterization,* 31(3), 293-316.
- ❖ Shi, Y., Pang, Y., Huang, N., Sun, C., Pan, Y., Cheng, Y., ... & Zheng, H. (2019). Competitive method for fluorescent dopamine detection in cerebrospinal fluid based on the peroxidase-like activity of ficin. *Spectrochim. Acta - A: Mol. Bio.* 209, 8-13.
- ❖ Shibata, M., Kawata, N., Masumoto, T., & Kimura, H. (1988). Selective hydrogenation of unsaturated carbonyl compounds over an oxidized gold–zirconium alloy. *J. Chem. Soc. Chem. Commun.* (3), 154-156.

- ❖ Shoaei, N., Daneshpour, M., Azimzadeh, M., Mahshid, S., Khoshfetrat, S. M., Jahanpeyma, F., ... & Foruzandeh, M. (2019). Electrochemical sensors and biosensors based on the use of polyaniline and its nanocomposites: A review on recent advances. *Microchim. Acta.* 186(7), 1-29.
- ❖ Shukla, R., Bansal, V., Chaudhary, M., Basu, A., Bhonde, R. R., & Sastry, M. (2005). Biocompatibility of gold nanoparticles and their endocytotic fate inside the cellular compartment: a microscopic overview. *Langmuir.* 21(23), 10644-10654.
- ❖ Shylesh, S., Schuenemann, V., & Thiel, W. R. (2010). Magnetically separable nanocatalysts: bridges between homogeneous and heterogeneous catalysis. *Angew. Chem. Int. Ed.* 49(20), 3428-3459.
- ❖ Siamaki, A. R., Abd El Rahman, S. K., Abdelsayed, V., El-Shall, M. S., & Gupton, B. F. (2011). Microwave-assisted synthesis of palladium nanoparticles supported on graphene: A highly active and recyclable catalyst for carbon–carbon cross-coupling reactions. *J. Catal.* 279(1), 1-11.
- ❖ Siedlecka, E. M., & Stepnowski, P. (2009). The effect of alkyl chain length on the degradation of alkylimidazolium-and pyridinium-type ionic liquids in a Fenton-like system. *Environ. Sci. Pollut. Res.* 16(4), 453-458.
- ❖ Singh, L.; Rekha, P.; Chand, S. (2016). Cu-impregnated zeolite Y as highly active and stable heterogeneous Fenton-like catalyst for degradation of Congo red dye. *Sep. Purif. Technol.*, 170, 321–336.
- ❖ Singh, M., Singh, S., Prasad, S., & Gambhir, I. S. (2008). Nanotechnology in medicine and anti bacterial effect of silver nanoparticles. *Dig. J. Nanomater .Biostructures.* 3(3), 115-122.

- ❖ Singh, S. B., & Tandon, P. K. (2014). Catalysis: a brief review on nanocatalyst. *J. Energy Chem.* 2(3), 106-115.
- ❖ Singh, S. K., & Xu, Q. (2013). Nanocatalysts for hydrogen generation from hydrazine. *Catal. Sci. Technol.* 3(8), 1889-1900.
- ❖ Singh, S. K., Singh, A. K., Aranishi, K., & Xu, Q. (2011). Noble-metal-free bimetallic nanoparticle-catalyzed selective hydrogen generation from hydrous hydrazine for chemical hydrogen storage. *J. Am. Chem. Soc.* 133(49), 19638-19641.
- ❖ Singh, S. K., Zhang, X. B., & Xu, Q. (2009). Room-temperature hydrogen generation from hydrous hydrazine for chemical hydrogen storage. *J. Am. Chem. Soc.* 131(29), 9894-9895.
- ❖ Singh, S., Bharti, A., & Meena, V. K. (2014). Structural, thermal, zeta potential and electrical properties of disaccharide reduced silver nanoparticles. *J. Mater. Sci.: Mater. Electron.* 25(9), 3747-3752.
- ❖ Singh, S.; Pandey, P.C. 2020. Synthesis and application of functional Prussian blue nanoparticles for toxic dye degradation. *J. Environ. Chem. Eng.* 8(3), 103753.
- ❖ Somorjai, G. A., Frei, H., & Park, J. Y. (2009). Advancing the frontiers in nanocatalysis, biointerfaces, and renewable energy conversion by innovations of surface techniques. *J. Am. Chem. Soc.* 131(46), 16589-16605.
- ❖ Song, J., Ran, R., & Shao, Z. (2010). Hydrazine as efficient fuel for low-temperature SOFC through ex-situ catalytic decomposition with high selectivity toward hydrogen. *Int. J. Hydrog. Energy.* 35(15), 7919-7924.

- ❖ Song, Y., Liu, G., Wang, J., Dong, X., & Yu, W. (2014). Synthesis and luminescence resonance energy transfer based on noble metal nanoparticles and the NaYF₄: Tb³⁺ shell. *Phys. Chem. Chem. Phys.* 16(29), 15139-15145.
- ❖ Spata, V. A., & Carter, E. A. (2018). Mechanistic insights into photocatalyzed hydrogen desorption from palladium surfaces assisted by localized surface plasmon resonances. *ACS nano*, 12(4), 3512-3522.
- ❖ Sreekanth, T. V. M., Jung, M. J., & Eom, I. Y. (2016). Green synthesis of silver nanoparticles, decorated on graphene oxide nanosheets and their catalytic activity. *Appl. Surf. Sci.* 361, 102-106.
- ❖ Srinoi, P., Chen, Y. T., Vittur, V., Marquez, M. D., & Lee, T. R. (2018). Bimetallic nanoparticles: enhanced magnetic and optical properties for emerging biological applications. *Appl. Sci.* 8(7), 1106-1138.
- ❖ Su, B., Zhang, M., Shao, Y., & Girault, H. H. (2006). Solvent effect on redox properties of hexanethiolate monolayer-protected gold nanoclusters. *J. Phys. Chem. B.* 110(43), 21460-21466.
- ❖ Su, S., Wu, W., Gao, J., Lu, J., & Fan, C. (2012). Nanomaterials-based sensors for applications in environmental monitoring. *J. Mater. Chem.* 22(35), 18101-18110.
- ❖ Subramaniam, C., Pradeep, T., & Chakrabarti, J. (2005). Flow-induced transverse electrical potential across an assembly of gold nanoparticles. *Phys. Rev. Lett.* 95(16), 164501-164505.
- ❖ Sugunan, A., Thanachayanont, C., Dutta, J., & Hilborn, J. G. (2005). Heavy-metal ion sensors using chitosan-capped gold nanoparticles. *Sci. Technol. Adv. Mater.* 6(3-4), 335.

- ❖ Sugunan, A., Thanachayanont, C., Dutta, J., & Hilborn, J. G. (2005). Heavy-metal ion sensors using chitosan-capped gold nanoparticles. *Sci. Technol. Adv. Mater.* 6(3-4), 335.
- ❖ Sun, P., Hai, J., Sun, S., Lu, S., Liu, S., Liu, H., ... & Wang, B. (2020). Aqueous stable Pd nanoparticles/covalent organic framework nanocomposite: an efficient nanoenzyme for colorimetric detection and multicolor imaging of cancer cells. *Nanoscale*. 12(2), 825-831.
- ❖ Sun, Y., & Wang, H. H. (2007). High-performance, flexible hydrogen sensors that use carbon nanotubes decorated with palladium nanoparticles. *Adv. Mater.* 19(19), 2818-2823.
- ❖ Suwannarat, K., Thongthai, K., Ananta, S., & Srisombat, L. (2018). Synthesis of hollow trimetallic Ag/Au/Pd nanoparticles for reduction of 4-nitrophenol. *Colloids Surf. A: Physicochem. Eng. Asp.* 540, 73-80.
- ❖ Syafiuddin, A., Salim, M. R., Beng Hong Kueh, A., Hadibarata, T., & Nur, H. (2017). A review of silver nanoparticles: research trends, global consumption, synthesis, properties, and future challenges. *J. Chin. Chem. Soc.* 64(7), 732-756.
- ❖ Taleb, A., Petit, C., & Pilani, M. P. (1998). Optical properties of self-assembled 2D and 3D superlattices of silver nanoparticles. *The Journal of J. Phys. Chem. B.* 102(12), 2214-2220.
- ❖ Talekar, M., Kendall, J., Denny, W., & Garg, S. (2011). Targeting of nanoparticles in cancer: drug delivery and diagnostics. *AntiCancer Drugs*. 22(10), 949-962.

- ❖ Tang, J., Deng, L., Deng, H., Xiao, S., Zhang, X., & Hu, W. (2014). Surface segregation and chemical ordering patterns of Ag–Pd nanoalloys: Energetic factors, nanoscale effects, and catalytic implication. *J. Phys. Chem. C.* 118(48), 27850-27860.
- ❖ Tang, X.; Wang, Y.; Han, J.; Ni, L.; Zhang, H.; Li, C.; Li, J.; Qiu, Y. 2018. A novel fluorescent probe based on biphenyl and rhodamine for multi-metal ion recognition and its application. *Dalton Trans.*, 47(10), 3378–3387.
- ❖ Tang, Y., Chi, X., Zou, S., & Zeng, X. (2016). Facet effects of palladium nanocrystals for oxygen reduction in ionic liquids and for sensing applications. *Nanoscale*. 8(10), 5771-5779.
- ❖ Taniguchi, N. (1974) On the basic concept of nano-technology Proceedings of the International Conference on Production Engineering Tokyo Part II Japan Society of Precision Engineering.
- ❖ Tao, Y., Lin, Y., Ren, J., & Qu, X. (2013). A dual fluorometric and colorimetric sensor for dopamine based on BSA-stabilized Aunanoclusters. *Biosens. Bioelectron.* 42, 41-46.
- ❖ Tao, Y., Lin, Y., Ren, J., & Qu, X. (2013). A dual fluorometric and colorimetric sensor for dopamine based on BSA-stabilized Aunanoclusters. *Biosens. Bioelectron.* 42, 41-46.
- ❖ Teniou, A., Rhouati, A., & Catanante, G. (2022). A Simple Fluorescent Aptasensing Platform Based on Graphene Oxide for Dopamine Determination. *Appl. Biochem. Biotechnol.* 194, 1925–1937.
- ❖ Thanh, N. T., Maclean, N., & Mahiddine, S. (2014). Mechanisms of nucleation and growth of nanoparticles in solution. *Chem. Rev.* 114(15), 7610-7630.
- ❖ Thomas, S. W., Joly, G. D., & Swager, T. M. (2007). Chemical sensors based on amplifying fluorescent conjugated polymers. *Chem. Rev.* 107(4), 1339-1386.

- ❖ Ting, S. L., Ee, S. J., Ananthanarayanan, A., Leong, K. C., & Chen, P. (2015). Graphene quantum dots functionalized gold nanoparticles for sensitive electrochemical detection of heavy metal ions. *Electrochim. Acta.* 172, 7-11.
- ❖ Topuz, F., & Uyar, T. (2019). Atomic layer deposition of palladium nanoparticles on a functional electrospun poly-cyclodextrin nanoweb as a flexible and reusable heterogeneous nanocatalyst for the reduction of nitroaromatic compounds. *Nanoscale Adv.* 1(10), 4082-4089.
- ❖ Toshima, N., & Yonezawa, T. (1998). Bimetallic nanoparticles—novel materials for chemical and physical applications. *New. J. Chem.* 22(11), 1179-1201.
- ❖ Tripathi, V. S., Kandimalla, V. B., & Ju, H. (2006). Preparation of ormosil and its applications in the immobilizing biomolecules. *Sensor Actuat B: Chem.* 114(2), 1071-1082.
- ❖ Turkevich, J., Stevenson, P. C., Hiller, J. (1951). The nucleation and growth processes in the synthesis of colloidal Au. *Discus Faaday Soc.* 11, 55-75.
- ❖ Umamaheswari, C., Lakshmanan, A., & Nagarajan, N. S. (2018). Green synthesis, characterization and catalytic degradation studies of gold nanoparticles against congo red and methyl orange. *J. Photochem. Photobiol. B: Biol.* 178, 33-39.
- ❖ Uppal, M. A., Kafizas, A., Ewing, M. B., & Parkin, I. P. (2013). The room temperature formation of gold nanoparticles from the reaction of cyclohexanone and auric acid; a transition from dendritic particles to compact shapes and nanoplates. *J. Mater. Chem. A.* 1(25), 7351-7359.
- ❖ Veisi, H., Moradi, S. B., Saljooqi, A., & Safarimehr, P. (2019). Silver nanoparticle-decorated on tannic acid-modified magnetite nanoparticles ($\text{Fe}_3\text{O}_4 @ \text{TA/Ag}$) for highly

- active catalytic reduction of 4-nitrophenol, Rhodamine B and Methylene blue. *Mater. Sci. Eng. C.* 100, 445-452.
- ❖ Venkatesan, P., & Santhanalakshmi, J. (2012). Synthesis and characterization of surfactant stabilized trimetallic Au-Ag-Pd nanoparticles for heck coupling reaction. *Phys Chem.* 2, 12-15.
- ❖ Vijayaraj, K., Dinakaran, T., Lee, Y., Kim, S., Kim, H. S., Lee, J., & Chang, S. C. (2017). One-step construction of a molybdenum disulfide/multi-walled carbon nanotubes/polypyrrole nanocomposite biosensor for the ex-vivo detection of dopamine in mouse brain tissue. *Biochemi. Biophys. Research. Communi.* 494(1-2), 181-187.
- ❖ Waghmare, M., Khade, B., Chaudhari, P., & Dongre, P. (2018). Multiple layer formation of bovine serum albumin on silver nanoparticles revealed by dynamic light scattering and spectroscopic techniques. *J. Nanoparticle Res.* 20(7), 1-21.
- ❖ Wang, C., & Deng, L. (2014). Electrically controlled plasmonic lasing resonances with silver nanoparticles embedded in amplifying nematic liquid crystals. *Laser Phys. Lett.* 11(11), 115814.
- ❖ Wang, C., Yin, J., Han, S., Jiao, T., Bai, Z., Zhou, J., ... & Peng, Q. (2019). Preparation of palladium nanoparticles decorated polyethyleneimine/polycaprolactone composite fibers constructed by electrospinning with highly efficient and recyclable catalytic performances. *Catalysts*, 9(6), 559.
- ❖ Wang, Chengyan, Xingrong Tan, Shihong Chen, Ruo Yuan, Fangxin Hu, Dehua Yuan, and Yun Xiang. "Highly-sensitive cholesterol biosensor based on platinum–gold hybrid functionalized ZnO nanorods." *Talanta* 94 (2012): 263-270.

- ❖ Wang, H. B., Li, Y., Dong, G. L., Gan, T., & Liu, Y. M. (2017). A convenient and label-free colorimetric assay for dopamine detection based on the inhibition of the Cu (ii)-catalyzed oxidation of a 3, 3', 5, 5'-tetramethylbenzidine–H₂O₂ system. *New J. Chem.* 41(23), 14364-14369.
- ❖ Wang, J. (2012). Electrochemical biosensing based on noble metal nanoparticles. *Mikrochim. Acta*. 177(3), 245-270.
- ❖ Wang, J., Lin, W., Cao, E., Xu, X., Liang, W., & Zhang, X. (2017). Surface plasmon resonance sensors on Raman and fluorescence spectroscopy. *Sensors*, 17(12), 2719.
- ❖ Wang, L., & Yamauchi, Y. (2010). Autoprogrammed synthesis of triple-layered Au@Pd@ Pt core– shell nanoparticles consisting of a Au@ Pd bimetallic core and nanoporous Pt shell. *J. Am. Chem.Soc.* 132(39), 13636-13638.
- ❖ Wang, L., & Yamauchi, Y. (2011). Strategic synthesis of trimetallic Au@ Pd@ Pt core– shell nanoparticles from poly (vinylpyrrolidone)-based aqueous solution toward highly active electrocatalysts. *Chem. Mater.* 23(9), 2457-2465.
- ❖ Wang, L., Liu, Y., Li, W., Jiang, X., Ji, Y., Wu, X., ... & Chen, C. (2011). Selective targeting of gold nanorods at the mitochondria of cancer cells: implications for cancer therapy. *Nano lett.* 11(2), 772-780.
- ❖ Wang, Q., Cui, X., Chen, J., Zheng, X., Liu, C., Xue, T., ... & Zheng, W. (2012). Well-dispersed palladium nanoparticles on graphene oxide as a non-enzymatic glucose sensor. *RSC Adv.* 2(15), 6245-6249.
- ❖ Wang, W., Niu, Q., Zeng, G., Zhang, C., Huang, D., Shao, B., ... & Tang, X. (2020). 1D porous tubular g-C₃N₄ capture black phosphorus quantum dots as 1D/0D metal-free

- photocatalysts for oxytetracycline hydrochloride degradation and hexavalent chromium reduction. *Appl. Catal. B: Environ.* 273, 119051-119064.
- ❖ Wang, W., Wang, S., Ma, X., & Gong, J. (2011). Recent advances in catalytic hydrogenation of carbon dioxide. *Chem. Soc. Rev.* 40(7), 3703-3727.
- ❖ Wang, Y. C., DuChene, J. S., Huo, F., & Wei, W. D. (2014). An in situ approach for facile fabrication of robust and scalable SERS substrates. *Nanoscale*, 6(13), 7232-7236.
- ❖ Wang, Y. R., & Chu, W. (2011). Degradation of a xanthene dye by Fe (II)-mediated activation of Oxone process. *J. Hazard. Mater.* 186(2-3), 1455-1461.
- ❖ Wang, Y., Liu, J., Wang, P., Werth, C. J., & Strathmann, T. J. (2014). Palladium nanoparticles encapsulated in core–shell silica: A structured hydrogenation catalyst with enhanced activity for reduction of oxyanion water pollutants. *ACS Catal.* 4(10), 3551-3559.
- ❖ Wang, Y.R.; Chu, W. (2011). Degradation of a xanthene dye by Fe(II)-mediated activation of Oxone process. *J. Hazad. Mater.* 186(2–3), 1455–1461.
- ❖ Wang, Z. Y., Li, P., Cui, L., Qiu, J. G., Jiang, B., & Zhang, C. Y. (2020). Integration of nanomaterials with nucleic acid amplification approaches for biosensing. *Trends Anal. Chem.* 129, 115959.
- ❖ Wang, Z., Chen, X., Li, K., Bi, S., Wu, C., & Chen, L. (2015). Preparation and catalytic property of PVDF composite membrane with polymeric spheres decorated by Pd nanoparticles in membrane pores. *J. Membr. Sci.* 496, 95-107.
- ❖ Weber, M., Kim, J. Y., Lee, J. H., Kim, J. H., Iatsunskyi, I., Coy, E., ... & Kim, S. S. (2019). Highly efficient hydrogen sensors based on Pd nanoparticles supported on boron nitride coated ZnO nanowires. *J. Mater. Chem. A.* 7(14), 8107-8116.

- ❖ Wei, H., Rodriguez, K., Renneckar, S., & Vikesland, P. J. (2014). Environmental science and engineering applications of nanocellulose-based nanocomposites. *Environ. Sci. Nano.* 1(4), 302-316.
- ❖ Wei, X., Zhang, Z., & Wang, Z. (2019). A simple dopamine detection method based on fluorescence analysis and dopamine polymerization. *Microchem. J.* 145, 55-58.
- ❖ Winter, C. J. (2009). Hydrogen energy—Abundant, efficient, clean: A debate over the energy-system-of-change. *Int. j. hydrog. Energy.* 34(14), S1-S52.
- ❖ Wouters, D., & Schubert, U. S. (2004). Nanolithography and nanochemistry: probe-related patterning techniques and chemical modification for nanometer-sized devices. *Angew. Chem. Int. Ed.* 43(19), 2480-2495.
- ❖ Wu, L., Li, Z. W., Zhang, F., He, Y. M., & Fan, Q. H. (2008). Air-Stable and Highly Active Dendritic Phosphine Oxide-Stabilized Palladium Nanoparticles: Preparation, Characterization and Applications in the Carbon-Carbon Bond Formation and Hydrogenation Reactions. *Adv. Synth. Catal.* 350(6), 846-862.
- ❖ Wu, P., Ding, P., Ye, X., Li, L., He, X., & Wang, K. (2019). One-pot synthesized Cu/Au/Pt trimetallic nanoparticles as a novel enzyme mimic for biosensing applications. *RSC adv.* 9(26), 14982-14989.
- ❖ Wu, Z. S., Jiang, J. H., Fu, L., Shen, G. L., & Yu, R. Q. (2006). Optical detection of DNA hybridization based on fluorescence quenching of tagged oligonucleotide probes by gold nanoparticles. *Anal. Biochem.* 353(1), 22-29.
- ❖ Xia, Y., Xiong, Y., Lim, B., & Skrabalak, S. E. (2009). Shape-controlled synthesis of metal nanocrystals: simple chemistry meets complex physics. *Angew. Chem. Int. Ed.* 48(1), 60-103.

- ❖ Xu, H.M.; Sun, X.F.; Wang, S.Y.; Song, C.; Wang, S.G. (2018). Development of laccase/graphene oxide membrane for enhanced synthetic dyes separation and degradation. *Sep. Purif. Technol.* 204, 255–260.
- ❖ Xu, L., Zheng, G., Liu, Y., Su, J., Kuang, W., & Rao, W. (2017). The optical properties of Ag/ZnO nanocomposite thin films with different thickness. *Optik.* 147, 6-13.
- ❖ Yadav, M., & Xu, Q. (2012). Liquid-phase chemical hydrogen storage materials. *Energy Environ. Sci.* 5(12), 9698-9725.
- ❖ Yadav, N., Jaiswal, A. K., Dey, K. K., Yadav, V. B., Nath, G., Srivastava, A. K., & Yadav, R. R. (2018). Trimetallic Au/Pt/Ag based nanofluid for enhanced antibacterial response. *Mater. Chem. Phys.* 218, 10-17.
- ❖ Yamauchi, M., Kobayashi, H., & Kitagawa, H. (2009). Hydrogen storage mediated by Pd and Pt nanoparticles. *ChemPhysChem.* 10(15), 2566-2576.
- ❖ Yaneva, Z.; Georgieva, N. 2012. Insights into Congo red Adsorption on Agro-Industrial MaterialsSpectral, Equilibrium, Kinetic, Thermodynamic, Dynamic and Desorption Studies. A Review. *Int. J. Chem. Eng.* 4(2), 127–146.
- ❖ Yang, F., Kung, S. C., Cheng, M., Hemminger, J. C., & Penner, R. M. (2010). Smaller is faster and more sensitive: the effect of wire size on the detection of hydrogen by single palladium nanowires. *Acs Nano.* 4(9), 5233-5244.
- ❖ Yang, X., Chen, J., Hu, L., Wei, J., Shuai, M., Huang, D., ... & Zhao, P. (2019). Palladium separation by Pd-catalyzed gel formation via alkyne coupling. *Chem. Mater.* 31(18), 7386-7394.

- ❖ Yang, Y., & Kimura, K. (2006). Highly ordered superlattices from polydisperse Ag nanoparticles: A comparative study of fractionation and self-correction. *J. Phys. Chem.* 110(48), 24442-24449.
- ❖ Yang, Y., Wang, H., Wu, Y., & Yu, X. (2022). Dual recognition strategy for selective fluorescent detection of dopamine and antioxidants based on graphite carbon nitride in human blood serum. *Spectrochim. Acta - A: Mol. Bio.* 265, 120385-120393.
- ❖ Yang, Y., Zeng, G., Huang, D., Zhang, C., He, D., Zhou, C., ... & Zhou, Y. (2020). Molecular engineering of polymeric carbon nitride for highly efficient photocatalytic oxytetracycline degradation and H₂O₂ production. *Appl. Catal. B: Environ.* 272, 118970-118984.
- ❖ Yang, Y., Zhang, C., Huang, D., Zeng, G., Huang, J., Lai, C., ... & Xiong, W. (2019). Boron nitride quantum dots decorated ultrathin porous g-C₃N₄: intensified exciton dissociation and charge transfer for promoting visible-light-driven molecular oxygen activation. *Appl. Catal. B: Environ.* 245, 87-99.
- ❖ Yang, Z., Ma, F., Zhu, Y., Chen, S., Wang, C., & Lu, X. (2017). A facile synthesis of CuFe₂O₄/Cu₉S₈/PPy ternary nanotubes as peroxidase mimics for the sensitive colorimetric detection of H₂O₂ and dopamine. *Dalton Trans.* 46(34), 11171-11179.
- ❖ Yao, S., Liu, Z., & Li, L. (2021). Recent Progress in Nanoscale Covalent Organic Frameworks for Cancer Diagnosis and Therapy. *Nanomicro. Lett.* 13(1), 1-20.
- ❖ You, Y., Zou, J., Li, W. J., Chen, J., Jiang, X. Y., & Yu, J. G. (2022). Novel lanthanum vanadate-based nanocomposite for simultaneously electrochemical detection of dopamine and uric acid in fetal bovine serum. *Int. J. Biol. Macromol.* 195, 346-355.

- ❖ Yu, Y. H., Ma, C. C. M., Teng, C. C., Huang, Y. L., Lee, S. H., Wang, I., & Wei, M. H. (2012). Electrical, morphological, and electromagnetic interference shielding properties of silver nanowires and nanoparticles conductive composites. *Mater. Chem. Phys.* 136(2-3), 334-340.
- ❖ Zaleska-Medyńska, A., Marchelek, M., Diak, M., & Grabowska, E. (2016). Noble metal-based bimetallic nanoparticles: the effect of the structure on the optical, catalytic and photocatalytic properties. *Adv. Colloid Interface Sci.* 229, 80-107.
- ❖ Zhang, C., Chen, B. Q., Li, Z. Y., Xia, Y., & Chen, Y. G. (2015). Surface plasmon resonance in bimetallic core–shell nanoparticles. *J. Phys. Chem. A.* 119(29), 16836-16845.
- ❖ Zhang, F. Q., Peng, K. Q., Sun, R. N., Hu, Y., & Lee, S. T. (2015). Light trapping in randomly arranged silicon nanorocket arrays for photovoltaic applications. *Nanotechnology*, 26(37), 375401.
- ❖ Zhang, M. L., Fan, X., Zhou, H. W., Shao, M. W., Zapien, J. A., Wong, N. B., & Lee, S. T. (2010). A high-efficiency surface-enhanced Raman scattering substrate based on silicon nanowires array decorated with silver nanoparticles. *J. Phys. Chem. A.* 114(5), 1969-1975.
- ❖ Zhang, M., Liu, Y., Wang, J., & Tang, J. (2019). Photodeposition of palladium nanoparticles on a porous gallium nitride electrode for nonenzymatic electrochemical sensing of glucose. *Microchim. Acta.* 186(2), 1-8.
- ❖ Zhang, X. (2015). Gold nanoparticles: recent advances in the biomedical applications. *Cell Biochem. Biophys.* 72(3), 771-775.
- ❖ Zhang, X., Du, B., Wu, D., Ma, H., Zhang, Y., Li, H., & Wei, Q. (2015). Signal amplification strategy of triple-layered core–shell Au@ Pd@ Pt nanoparticles for

- ultrasensitive immunoassay detection of squamous cell carcinoma antigen. *J. Biomed. Nanotechnol.* 1(2), 245-252.
- ❖ Zhang, X., Qu, Y., Shen, W., You, S., Pei, X., Li, S., ... & Zhou, J. (2016). Colorimetric response of biogenetic gold nanoparticles to mercury (II) ions. *Colloids Surf. A: Physicochem. Eng. Asp.* 508, 360-365.
- ❖ Zhang, Z., Wang, H., Chen, Z., Wang, X., Choo, J., & Chen, L. (2018). Plasmonic colorimetric sensors based on etching and growth of noble metal nanoparticles: Strategies and applications. *Biosens. Bioelectron.* 114, 52-65.
- ❖ Zhao, C., Jiao, Y., Hua, J., Yang, J., & Yang, Y. (2018). Hydrothermal synthesis of nitrogen-doped carbon quantum dots as fluorescent probes for the detection of dopamine. *J. fluoresc.* 28(1), 269-276.
- ❖ Zhao, J., Pinchuk, A. O., McMahon, J. M., Li, S., Ausman, L. K., Atkinson, A. L., & Schatz, G. C. (2008). Methods for describing the electromagnetic properties of silver and gold nanoparticles. *Acc. Chem. Res.* 41(12), 1710-1720.
- ❖ Zhao, L., Zhang, Y., Huang, L. B., Liu, X. Z., Zhang, Q. H., He, C., ... & Wan, L. J. (2019). Cascade anchoring strategy for general mass production of high-loading single-atomic metal-nitrogen catalysts. *Nat. Commun.* 10(1), 1-11.
- ❖ Zhao, X., Zhao, H., Yan, L., Li, N., Shi, J., & Jiang, C. (2020). Recent developments in detection using noble metal nanoparticles. *Crit. Rev. Anal. Chem.* 50(2), 97-110.
- ❖ Zharov, V. P., Kim, J. W., Curiel, D. T., & Everts, M. (2005). Self-assembling nanoclusters in living systems: application for integrated photothermal nanodiagnostics and nanotherapy. *Nanomed.: Nanotechnol. Biol. Med.* 1(4), 326-345.

- ❖ Zheng, M., Cheng, R., Chen, X., Li, N., Li, L., Wang, X., & Zhang, T. (2005). A novel approach for CO-free H₂ production via catalytic decomposition of hydrazine. *Int. J. Hydrom. Energy.* 30(10), 1081-1089.
- ❖ Zhou, W. P., Lewera, A., Larsen, R., Masel, R. I., Bagus, P. S., & Wieckowski, A. (2006). Size effects in electronic and catalytic properties of unsupported palladium nanoparticles in electrooxidation of formic acid. *J. Phys. Chem. B.* 110(27), 13393-13398.
- ❖ Zhu, L., Xu, G., Song, Q., Tang, T., Wang, X., Wei, F., & Hu, Q. (2016). Highly sensitive determination of dopamine by a turn-on fluorescent biosensor based on aptamer labeled carbon dots and nano-graphite. *Sens. Actuators B. Chem.* 231, 506-512.
- ❖ Zhu, X., Li, J., He, H., Huang, M., Zhang, X., & Wang, S. (2015). Application of nanomaterials in the bioanalytical detection of disease-related genes. *Biosens. Bioelectron.* 74, 113-133.

List of publications

1. Pandey, P. C., **Mitra, M. D.**, Tiwari, A. K., & Singh, S. (2021). Synthetic incorporation of palladium-nickel bimetallic nanoparticles within mesoporous silica/silica nanoparticles as efficient and cheaper catalyst for both cationic and anionic dyes degradation. **J. Environ. Sci. Health Part A**. 56(4), 460-472.
2. Pandey, P. C., **Mitra, M. D.**, Shukla, S., & Narayan, R. J. (2021). Organotrialkoxysilane-Functionalized Noble Metal Monometallic, Bimetallic, and Trimetallic Nanoparticle Mediated Non-Enzymatic Sensing of Glucose by Resonance Rayleigh Scattering. **Biosensors**, 11(4), 122.
3. Pandey, P. C., **Mitra, M. D.**, Shukla, S., & Narayan, R. J. (2021). Organotrialkoxysilane-functionalized mesoporous Pd–Ni nanocatalyst for selective hydrazine decomposition and sensing. **MRS Comm.** 11(1), 78-85.
4. Pandey, P. C., **Mitra, M. D.**, Pandey, A. K., & Narayan, R. J. (2021). Organotrialkoxysilane mediated rapid and controlled synthesis metal nanoparticles in both homogeneous and heterogeneous phase and their catalytic applications. **MRS Adv.** 6(3), 43-53.
5. **Mitra, M. D.**, & Pandey, P. C. (2022). Functional trialkoxysilane mediated controlled synthesis of fluorescent gold nanoparticles and fluoremetric sensing of dopamine. **Optical Materials**, 132, 112810.

Conferences

Oral Presentations

- ❖ **Mitra, M. D.**, Pandey, P. C., Functional trialkoxysilane mediated synthesis of noble metal nanoparticles and their applications.“International Conference on Emerging Trends in Biotechnology (Biosangam-2022)” 10-12 Mar. 2022, organized by Department of Biotechnology, Motilal Nehru National Institute of Technology, Allahabad, Prayagraj, U.P, India.

Poster Presentations

- ❖ **Mitra, M. D.**, Pandey, P. C., Organo trialkoxysilane mediated synthesis of monometallic bimetallic and trimetallic nanoparticles mediated non-enzymatic sensing of glucose by resonance Rayleigh scattering, 3rd International conference on Environmental Agricultural and Chemical and Biological science (ICEACBS-2022) (Virtual mode) PG research department of zoology, The American College, Tamilnadu India. (**Best Poster presentation**).

- ❖ **Mitra, M. D.**, Pandey, P. C., epoxycyclohexyle ethyletrimethoxysilane mediated synthesis of Pd-Ni bimetallic nanoparticles and their use as a recyclable heterogeneous catalyst for dyes. Contemporary Trend And Future Prospect Of Fundamental Materials (CTFM) degradation. Department of chemistry, Banaras, Hindu University, U. P, India.