
LIST OF FIGURES

| | | Page No. |
|-----|--|----------|
| 1.1 | Classification of immunosuppressants available in the market | 2 |
| 1.2 | Therapeutic functions of rapamycin | 7 |
| 2.1 | Structure of Rapamycin | 15 |
| 2.2 | Biosynthetic pathway of polyketide antibiotics | 20 |
| 2.3 | Route of precursors biosynthesis for rapamycin production | 20 |
| 2.4 | Variation of performance ratio with oxygen-transfer rate in different types of ALRs versus an agitation tank | 31 |
| 2.5 | Steps of purification of rapamycin | 42 |
| 3.1 | Microorganisms grown on agar plates | 47 |
| 3.2 | Microscopic image of <i>S.hygroscopicus</i> | 53 |
| 3.3 | Schematic diagram of internal loop airlift reactor | 57 |
| 3.4 | Schematic diagram of set up of Internal Loop Air lift Reactor | 58 |
| 3.5 | Different carriers used for immobilization | 64 |
| 3.6 | Standard curve for reducing sugar estimation using DNS method | 70 |
| 3.7 | Standard HPLC plot | 70 |
| 4.1 | Result of training of data set with 619 epoch using ANN | 78 |
| 4.2 | Graph showing training epoch cycles vs. calculated mean square error of the supervised training for the designed ANN | 78 |
| 4.3 | Fitness plot showing the GA iterations based on fitness function | 80 |
| 4.4 | Production of rapamycin in 3L Stirred tank reactor | 82 |

| | | |
|--------|--|-----|
| 4.5A | Model fitting of experimental observations of dry cell mass, reducing sugar concentration and rapamycin production during batch production at initial mannose concentration 15 g/L | 87 |
| 4.5B | Model fitting of experimental observations of dry cell mass, reducing sugar concentration and rapamycin production during batch production at initial mannose concentration 20 g/L | 87 |
| 4.5 C | Model fitting of experimental observations of dry cell mass, reducing sugar concentration and rapamycin production during batch production at initial mannose concentration 25 g/L | 88 |
| 4.5D | Model fitting of experimental observations of dry cell mass, reducing sugar concentration and rapamycin production during batch production at initial mannose concentration 30 g/L | 88 |
| 4.6 | Plot for determination of maximum specific growth rate (μ_{\max}) at different initial mannose concentration | 90 |
| 4.7 | Graph between γ and θ at initial sugar concentrations | 94 |
| 4.8 | Variation of rapamycin production, dry cell mass, dissolved oxygen and reducing sugar concentration with time | 97 |
| 4.9 | Variation of dissolved oxygen concentration and volumetric oxygen transfer coefficient ($k_L a$) with respect to time | 97 |
| 4.10 A | Shear stress (τ) vs Shear rate (γ) for Newtonian Fluid Model | 99 |
| 4.10 B | Shear stress (τ) vs Shear rate (γ) for Power Law Model | 99 |
| 4.10 C | Shear stress (τ) vs Shear rate (γ) for Casson Fluid Model | 100 |
| 4.10 D | Shear stress (τ) vs Shear rate (γ) for Bingham Plastic Model | 100 |

| | | |
|------|--|-----|
| 4.11 | Variation of apparent viscosities with time at different shear rates | 104 |
| 4.12 | Comparison of flow behavior index and consistency index with variation in biomass concentration | 104 |
| 4.13 | Plot of Residual Shear Stress vs independent variable | 105 |
| 4.14 | Variation of dry cell weight of <i>S. hygroscopicus</i> per unit weight of different carriers | 109 |
| 4.15 | SEM image of different immobilized carriers | 109 |
| 4.16 | Cell Release Study in different immobilized systems | 111 |
| 4.17 | Production of rapamycin in different immobilized carriers | 111 |
| 4.18 | Rapamycin production and cell release during repeated batch fermentation using <i>S.hygroscopicus</i> immobilized on PUF | 112 |
| 4.19 | Fermentation set up of internal loop ALR | 114 |
| 4.20 | Variation of gas hold up at different gas velocities | 117 |
| 4.21 | Variation of rapamycin concentration with aeration rate in an internal loop airlift reactor | 118 |
| 4.22 | Variation of Rapamycin production, dry cell mass and reducing sugar concentration with time during batch production of rapamycin in air lift reactor | 118 |
| 4.23 | Variation of specific growth rate (μ) and specific product formation rate (q_p) in an ALR | 119 |
| 4.24 | Variation in dissolved oxygen concentration and k_{La} during batch production of rapamycin in airlift reactor | 121 |
| 4.25 | Co-cultured plates of <i>S.hygroscopicus</i> and <i>C.albicans</i> | 123 |
| 4.26 | Variation in rapamycin production with different inoculum size of <i>C.albicans</i> in the co-culture study | 125 |

| | | |
|------|--|-----|
| 4.27 | Effect of time of inoculation of <i>C.albicans</i> on rapamycin production in co-culture | 125 |
| 4.28 | Fed- batch profile for rapamycin production based on pulse-feeding | 128 |
| 4.29 | Variation of pulse feeding on growth rate (dx/dt) and product formation rate (dp/dt) during fed-batch fermentation | 130 |
| 4.30 | Comparison of flow behavior index (n) and consistency index (k) for fed-batch fermentation | 132 |
| 4.31 | Summary of steps followed for rapamycin purification | 134 |
| 4.32 | HPTLC chromatogram | 136 |
| 4.33 | Elution profile of rapamycin by silica gel column chromatography | 136 |
| 4.34 | Overlay diagram of FTIR spectra of standard rapamycin and the purified sample | 138 |
| 4.35 | HPLC chromatograms (A) Standard rapamycin (B) Ethyl acetate extract (C) Purified sample | 139 |