

References

- [1] K. v. Klitzing, G. Dorda, and M. Pepper, “New Method for High-Accuracy Determination of the Fine-Structure Constant Based on Quantized Hall Resistance,” *Phys. Rev. Lett.*, vol. 45, no. 6, pp. 494–497, Aug. 1980.
- [2] J. E. Moore, “The birth of topological insulators,” *Nature*, vol. 464, no. 7286, pp. 194–198, Mar. 2010.
- [3] C. L. Kane and E. J. Mele, “Z2 Topological Order and the Quantum Spin Hall Effect,” *Phys. Rev. Lett.*, vol. 95, no. 14, p. 146802, Sep. 2005.
- [4] L. Fu, C. L. Kane, and E. J. Mele, “Topological Insulators in Three Dimensions,” *Phys. Rev. Lett.*, vol. 98, no. 10, p. 106803, Mar. 2007.
- [5] B. A. Bernevig, T. L. Hughes, and S.-C. Zhang, “Quantum Spin Hall Effect and Topological Phase Transition in HgTe Quantum Wells,” *Science (80-.)*, vol. 314, no. 5806, pp. 1757–1761, Dec. 2006.
- [6] C. H. Li, O. M. J. van ‘t Erve, S. Rajput, L. Li, and B. T. Jonker, “Direct comparison of current-induced spin polarization in topological insulator Bi₂Se₃ and InAs Rashba states,” *Nat. Commun.*, vol. 7, p. 13518, Nov. 2016.
- [7] M. Z. Hasan, C. L. Kane, and J. Gabel, “Topological Insulators,” *Rev. Mod. Phys.*, vol. 82, no. 4, p. 23, 2010.
- [8] G. Gupta, M. B. A. Jalil, and G. Liang, “Evaluation of mobility in thin Bi₂Se₃ Topological Insulator for prospects of Local Electrical Interconnects,” *Sci. Rep.*, vol. 4, no. 1, p. 6838, May 2015.
- [9] D. Hsieh *et al.*, “Observation of Unconventional Quantum Spin Textures in Topological Insulators,” *Science (80-.)*, vol. 323, no. 5916, pp. 919–922, Feb. 2009.
- [10] J. E. Hirsch, “Spin Hall Effect,” *Phys. Rev. Lett.*, vol. 83, no. 9, pp. 1834–1837, Aug. 1999.
- [11] N. Nagaosa, J. Sinova, S. Onoda, A. H. MacDonald, and N. P. Ong, “Anomalous Hall effect,” *Rev. Mod. Phys.*, vol. 82, no. 2, pp. 1539–1592, May 2010.
- [12] F. D. M. Haldane, “Model for a Quantum Hall Effect without Landau Levels: Condensed-Matter Realization of the ‘Parity Anomaly,’” *Phys. Rev. Lett.*, vol. 61, no. 18, pp. 2015–2018, Oct. 1988.
- [13] X.-L. Qi and S.-C. Zhang, “Topological insulators and superconductors,” *Rev. Mod. Phys.*, vol. 83, no. 4, pp. 1057–1110, Oct. 2011.
- [14] M. Konig *et al.*, “Quantum Spin Hall Insulator State in HgTe Quantum Wells,” *Science (80-.)*, vol. 318, no. 5851, pp. 766–770, Nov. 2007.
- [15] D. Hsieh *et al.*, “A topological Dirac insulator in a quantum spin Hall phase,” *Nature*, vol. 452, no. 7190, pp. 970–974, Apr. 2008.
- [16] M. Z. Hasan and C. L. Kane, “Colloquium : Topological insulators,” *Rev. Mod. Phys.*, vol. 82, no. 4, pp. 3045–3067, Nov. 2010.
- [17] H. Zhang, C.-X. Liu, X.-L. Qi, X. Dai, Z. Fang, and S.-C. Zhang, “Topological insulators

- in Bi₂Se₃, Bi₂Te₃ and Sb₂Te₃ with a single Dirac cone on the surface,” *Nat. Phys.*, vol. 5, no. 6, pp. 438–442, Jun. 2009.
- [18] Y. Xia *et al.*, “Observation of a large-gap topological-insulator class with a single Dirac cone on the surface,” *Nat. Phys.*, vol. 5, no. 6, pp. 398–402, Jun. 2009.
 - [19] J. J. Cha *et al.*, “Magnetic Doping and Kondo Effect in Bi₂Se₃ Nanoribbons,” *Nano Lett.*, vol. 10, no. 3, pp. 1076–1081, Mar. 2010.
 - [20] H. Steinberg, D. R. Gardner, Y. S. Lee, and P. Jarillo-Herrero, “Surface State Transport and Ambipolar Electric Field Effect in Bi₂Se₃ Nanodevices,” *Nano Lett.*, vol. 10, no. 12, pp. 5032–5036, Dec. 2010.
 - [21] L. Wu *et al.*, “Tuning and stabilizing topological insulator Bi₂Se₃ in the intrinsic regime by charge extraction with organic overlayers,” *Appl. Phys. Lett.*, vol. 108, no. 22, p. 221603, May 2016.
 - [22] D. Kong *et al.*, “Rapid Surface Oxidation as a Source of Surface Degradation Factor for Bi₂Se₃,” *ACS Nano*, vol. 5, no. 6, pp. 4698–4703, Jun. 2011.
 - [23] N. Bansal, Y. S. Kim, M. Brahlek, E. Edrey, and S. Oh, “Thickness-Independent Transport Channels in Topological Insulator Bi₂Se₃ Thin Films,” *Phys. Rev. Lett.*, vol. 109, no. 11, p. 116804, Sep. 2012.
 - [24] D. O. Scanlon *et al.*, “Controlling Bulk Conductivity in Topological Insulators: Key Role of Anti-Site Defects,” *Adv. Mater.*, vol. 24, no. 16, pp. 2154–2158, Apr. 2012.
 - [25] J. J. Cha, D. Kong, S.-S. Hong, J. G. Analytis, K. Lai, and Y. Cui, “Weak Antilocalization in Bi₂(Se_xTe_{1-x})₃ Nanoribbons and Nanoplates,” *Nano Lett.*, vol. 12, no. 2, pp. 1107–1111, Feb. 2012.
 - [26] A. A. Taskin, Z. Ren, S. Sasaki, K. Segawa, and Y. Ando, “Observation of Dirac Holes and Electrons in a Topological Insulator,” *Phys. Rev. Lett.*, vol. 107, no. 1, p. 16801, Jun. 2011.
 - [27] Y. Ando *et al.*, “Electrical Detection of the Spin Polarization Due to Charge Flow in the Surface State of the Topological Insulator Bi_{1.5}Sb_{0.5}Te_{1.7}Se_{1.3},” *Nano Lett.*, vol. 14, no. 11, pp. 6226–6230, Nov. 2014.
 - [28] J. Tang *et al.*, “Electrical Detection of Spin-Polarized Surface States Conduction in (Bi_{0.53}Sb_{0.47})₂Te₃ Topological Insulator,” *Nano Lett.*, vol. 14, no. 9, pp. 5423–5429, Sep. 2014.
 - [29] A. R. Melnik *et al.*, “Spin-transfer torque generated by a topological insulator,” *Nature*, vol. 511, no. 7510, pp. 449–451, Jul. 2014.
 - [30] J. S. Lee, A. Richardella, D. R. Hickey, K. A. Mkhoyan, and N. Samarth, “Mapping the chemical potential dependence of current-induced spin polarization in a topological insulator,” *Phys. Rev. B*, vol. 92, no. 15, p. 155312, Oct. 2015.
 - [31] J. Tian, I. Miotkowski, S. Hong, and Y. P. Chen, “Electrical injection and detection of spin-polarized currents in topological insulator Bi₂Te₂Se,” *Sci. Rep.*, vol. 5, no. 1, p. 14293, Nov. 2015.
 - [32] Y. Wang *et al.*, “Topological Surface States Originated Spin-Orbit Torques,” *Phys. Rev.*

Lett., vol. 114, no. 25, p. 257202, Jun. 2015.

- [33] Y. Fan *et al.*, “Magnetization switching through giant spin–orbit torque in a magnetically doped topological insulator heterostructure,” *Nat. Mater.*, vol. 13, no. 7, pp. 699–704, Jul. 2014.
- [34] B. Xia, P. Ren, A. Sulaev, P. Liu, S.-Q. Shen, and L. Wang, “Indications of surface-dominated transport in single crystalline nanoflake devices of topological insulator Bi_{1.5}Sb_{0.5}Te_{1.8}Se_{0.2},” *Phys. Rev. B*, vol. 87, no. 8, p. 85442, Feb. 2013.
- [35] K. S. Novoselov, “Electric Field Effect in Atomically Thin Carbon Films,” *Science (80-.)*, vol. 306, no. 5696, pp. 666–669, Oct. 2004.
- [36] C. R. Dean *et al.*, “Boron nitride substrates for high-quality graphene electronics,” *Nat. Nanotechnol.*, vol. 5, no. 10, pp. 722–726, Oct. 2010.
- [37] B. Radisavljevic, A. Radenovic, J. Brivio, V. Giacometti, and A. Kis, “Single-layer MoS₂ transistors,” *Nat. Nanotechnol.*, vol. 6, no. 3, pp. 147–150, Mar. 2011.
- [38] D. Braga, I. Gutiérrez Lezama, H. Berger, and A. F. Morpurgo, “Quantitative Determination of the Band Gap of WS₂ with Ambipolar Ionic Liquid-Gated Transistors,” *Nano Lett.*, vol. 12, no. 10, pp. 5218–5223, Oct. 2012.
- [39] C. S. Fadley, “X-ray photoelectron spectroscopy: Progress and perspectives,” *J. Electron Spectros. Relat. Phenomena*, vol. 178–179, no. C, pp. 2–32, 2010.
- [40] M. Z. Hasan and C. L. Kane, “Colloquium- : Topological insulators,” *Rev. Mod. Phys.*, vol. 82, no. 4, pp. 3045–3067, Nov. 2010.
- [41] Y. L. Chen *et al.*, “Experimental Realization of a Three-Dimensional Topological Insulator, Bi₂Te₃,” *Science (80-.)*, vol. 325, no. 5937, pp. 178–181, Jul. 2009.
- [42] Y. L. Chen *et al.*, “Massive Dirac Fermion on the Surface of a Magnetically Doped Topological Insulator,” *Science (80-.)*, vol. 329, no. 5992, pp. 659–662, Aug. 2010.
- [43] Y.-Y. Li *et al.*, “Intrinsic Topological Insulator Bi₂Te₃ Thin Films on Si and Their Thickness Limit,” *Adv. Mater.*, vol. 22, no. 36, pp. 4002–4007, Sep. 2010.
- [44] F. Xiu *et al.*, “Manipulating surface states in topological insulator nanoribbons,” *Nat. Nanotechnol.*, vol. 6, no. 4, pp. 216–221, Apr. 2011.
- [45] J. G. Analytis, R. D. McDonald, S. C. Riggs, J.-H. Chu, G. S. Boebinger, and I. R. Fisher, “Two-dimensional surface state in the quantum limit of a topological insulator,” *Nat. Phys.*, vol. 6, no. 12, pp. 960–964, Dec. 2010.
- [46] D. Hsieh *et al.*, “A tunable topological insulator in the spin helical Dirac transport regime,” *Nature*, vol. 460, no. 7259, pp. 1101–1105, Aug. 2009.
- [47] X.-L. Qi and S.-C. Zhang, “The quantum spin Hall effect and topological insulators,” *Phys. Today*, vol. 63, no. 1, pp. 33–38, Jan. 2010.
- [48] P. Roushan *et al.*, “Topological surface states protected from backscattering by chiral spin texture,” *Nature*, vol. 460, no. 7259, pp. 1106–1109, Aug. 2009.
- [49] Y. H. Choi, N. H. Jo, K. J. Lee, J. B. Yoon, C. Y. You, and M. H. Jung, “Transport and magnetic properties of Cr-, Fe-, Cu-doped topological insulators,” *J. Appl. Phys.*, vol. 109, no. 7, p. 07E312, Apr. 2011.

- [50] P. P. J. Haazen *et al.*, “Ferromagnetism in thin-film Cr-doped topological insulator Bi₂Se₃,” *Appl. Phys. Lett.*, vol. 100, no. 8, p. 82404, Feb. 2012.
- [51] J. S. Dyck, P. Hájek, P. Lošt’ák, and C. Uher, “Diluted magnetic semiconductors based on Sb_{2-x}V_xTe₃(0.01<~x<~0.03),” *Phys. Rev. B*, vol. 65, no. 11, p. 115212, Mar. 2002.
- [52] J. S. Dyck, Č. Drašar, P. Lošt’ák, and C. Uher, “Low-temperature ferromagnetic properties of the diluted magnetic semiconductor Sb_{2-x}Cr_xTe₃,” *Phys. Rev. B*, vol. 71, no. 11, p. 115214, Mar. 2005.
- [53] Y. S. Hor *et al.*, “Development of ferromagnetism in the doped topological insulator Bi_{2-x}Mn_xTe₃,” *Phys. Rev. B*, vol. 81, no. 19, p. 195203, May 2010.
- [54] J. S. Dyck, P. Švanda, P. Lošt’ák, J. Horák, W. Chen, and C. Uher, “Magnetic and transport properties of the V_[sub 2]–VI_[sub 3] diluted magnetic semiconductor Sb_[sub 2-x]Mn_{[sub x}]Te_[sub 3],” *J. Appl. Phys.*, vol. 94, no. 12, p. 7631, 2003.
- [55] C.-Z. Chang *et al.*, “Thin Films of Magnetically Doped Topological Insulator with Carrier-Independent Long-Range Ferromagnetic Order,” *Adv. Mater.*, vol. 25, no. 7, pp. 1065–1070, Feb. 2013.
- [56] J. Cai, A. Retzker, F. Jelezko, and M. B. Plenio, “A large-scale quantum simulator on a diamond surface at room temperature,” *Nat. Phys.*, vol. 9, no. 3, pp. 168–173, Mar. 2013.
- [57] R. Sevak Singh, X. Wang, W. Chen, Ariando, and A. T. S. Wee, “Large room-temperature quantum linear magnetoresistance in multilayered epitaxial graphene: Evidence for two-dimensional magnetotransport,” *Appl. Phys. Lett.*, vol. 101, no. 18, p. 183105, Oct. 2012.
- [58] T. Arakane *et al.*, “Tunable Dirac cone in the topological insulator Bi_{2-x}Sb_xTe_{3-y}Sey,” *Nat. Commun.*, vol. 3, no. 1, p. 636, Jan. 2012.
- [59] H. Cao *et al.*, “Quantized Hall Effect and Shubnikov–de Haas Oscillations in Highly Doped Bi₂Se₃: Evidence for Layered Transport of Bulk Carriers,” *Phys. Rev. Lett.*, vol. 108, no. 21, p. 216803, May 2012.
- [60] H. Tang, D. Liang, R. L. J. Qiu, and X. P. A. Gao, “Two-Dimensional Transport-Induced Linear Magneto-Resistance in Topological Insulator Bi₂Se₃ Nanoribbons,” *ACS Nano*, vol. 5, no. 9, pp. 7510–7516, Sep. 2011.
- [61] H. He *et al.*, “High-field linear magneto-resistance in topological insulator Bi₂Se₃ thin films,” *Appl. Phys. Lett.*, vol. 100, no. 3, p. 32105, Jan. 2012.
- [62] P. Gehring, M. Burghard, and K. Kern, “Gate-controlled linear magnetoresistance in thin Bi₂Se₃ sheets,” *Appl. Phys. Lett.*, vol. 100, no. 21, p. 212402, May 2012.
- [63] Y. Yan, L.-X. Wang, D.-P. Yu, and Z.-M. Liao, “Large magnetoresistance in high mobility topological insulator Bi₂Se₃,” *Appl. Phys. Lett.*, vol. 103, no. 3, p. 33106, Jul. 2013.
- [64] Y. S. Hor *et al.*, “p-type Bi₂Se₃ for topological insulator and low-temperature thermoelectric applications,” *Phys. Rev. B*, vol. 79, no. 19, p. 195208, May 2009.
- [65] H.-T. He *et al.*, “Impurity Effect on Weak Antilocalization in the Topological Insulator Bi₂Te₃,” *Phys. Rev. Lett.*, vol. 106, no. 16, p. 166805, Apr. 2011.
- [66] J. Chen *et al.*, “Gate-Voltage Control of Chemical Potential and Weak Antilocalization in

- Bi_2Se_3 " *Phys. Rev. Lett.*, vol. 105, no. 17, p. 176602, Oct. 2010.
- [67] L. Bao *et al.*, "Weak Anti-localization and Quantum Oscillations of Surface States in Topological Insulator $\text{Bi}_2\text{Se}_2\text{Te}$," *Sci. Rep.*, vol. 2, no. 1, p. 726, Dec. 2012.
- [68] A. A. Abrikosov, "Quantum magnetoresistance," *Phys. Rev. B*, vol. 58, no. 5, pp. 2788–2794, Aug. 1998.
- [69] X. Wang, Y. Du, S. Dou, and C. Zhang, "Room Temperature Giant and Linear Magnetoresistance in Topological Insulator Bi_2Te_3 Nanosheets," *Phys. Rev. Lett.*, vol. 108, no. 26, p. 266806, Jun. 2012.
- [70] M. M. Parish and P. B. Littlewood, "Non-saturating magnetoresistance in heavily disordered semiconductors," *Nature*, vol. 426, no. 6963, pp. 162–165, Nov. 2003.
- [71] C. Herring, "Effect of Random Inhomogeneities on Electrical and Galvanomagnetic Measurements," *J. Appl. Phys.*, vol. 31, no. 11, pp. 1939–1953, Nov. 1960.
- [72] L. Bao *et al.*, "Quantum Corrections Crossover and Ferromagnetism in Magnetic Topological Insulators," *Sci. Rep.*, vol. 3, no. 1, p. 2391, Dec. 2013.
- [73] M. Liu *et al.*, "Crossover between Weak Antilocalization and Weak Localization in a Magnetically Doped Topological Insulator," *Phys. Rev. Lett.*, vol. 108, no. 3, p. 36805, Jan. 2012.
- [74] J. G. Checkelsky, J. Ye, Y. Onose, Y. Iwasa, and Y. Tokura, "Dirac-fermion-mediated ferromagnetism in a topological insulator," *Nat. Phys.*, vol. 8, no. 10, pp. 729–733, Oct. 2012.
- [75] H.-Z. Lu, J. Shi, and S.-Q. Shen, "Competition between Weak Localization and Antilocalization in Topological Surface States," *Phys. Rev. Lett.*, vol. 107, no. 7, p. 76801, Aug. 2011.
- [76] J. J. Cha *et al.*, "Effects of Magnetic Doping on Weak Antilocalization in Narrow Bi_2Se_3 Nanoribbons," *Nano Lett.*, vol. 12, no. 8, pp. 4355–4359, Aug. 2012.
- [77] R. Singh *et al.*, "Unusual negative magnetoresistance in $\text{Bi}_2\text{Se}_3\text{-y}\text{S}_y$ topological insulator under perpendicular magnetic field," *Appl. Phys. Lett.*, vol. 112, no. 10, p. 102401, Mar. 2018.
- [78] Z. J. Yue, X. L. Wang, and S. S. Yan, "Semimetal-semiconductor transition and giant linear magnetoresistances in three-dimensional Dirac semimetal $\text{Bi}_{0.96}\text{Sb}_{0.04}$ single crystals," *Appl. Phys. Lett.*, vol. 107, no. 11, p. 112101, Sep. 2015.
- [79] X. Guo *et al.*, "Single domain Bi_2Se_3 films grown on $\text{InP}(111)\text{A}$ by molecular-beam epitaxy," *Appl. Phys. Lett.*, vol. 102, no. 15, p. 151604, Apr. 2013.
- [80] D.-X. Qu, Y. S. Hor, J. Xiong, R. J. Cava, and N. P. Ong, "Quantum Oscillations and Hall Anomaly of Surface States in the Topological Insulator Bi_2Te_3 ," *Science (80-.)*, vol. 329, no. 5993, pp. 821–824, Aug. 2010.
- [81] M. Petrushevsky *et al.*, "Probing the surface states in Bi_2Se_3 using Shubnikov-de Hass effect," *Phys. Rev. B*, vol. 86, no. 4, p. 45131, Jul. 2012.
- [82] J. G. Checkelsky, Y. S. Hor, R. J. Cava, and N. P. Ong, "Bulk Band Gap and Surface State Conduction Observed in Voltage-Tuned Crystals of the Topological Insulator Bi_2Se_3 "

Phys. Rev. Lett., vol. 106, no. 19, p. 196801, May 2011.

- [83] Y. H. Liu *et al.*, “Gate-tunable coherent transport in Se-capped Bi₂Se₃ grown on amorphous SiO₂/Si,” *Appl. Phys. Lett.*, vol. 107, no. 1, p. 12106, Jul. 2015.
- [84] L. Ohnoutek *et al.*, “Strong interband Faraday rotation in 3D topological insulator Bi₂Se₃,” *Sci. Rep.*, vol. 6, no. 1, p. 19087, May 2016.
- [85] T. O. Wehling, A. M. Black-Schaffer, and A. V. Balatsky, “Dirac materials,” *Adv. Phys.*, vol. 63, no. 1, pp. 1–76, Jan. 2014.
- [86] F. Arnold *et al.*, “Negative magnetoresistance without well-defined chirality in the Weyl semimetal TaP,” *Nat. Commun.*, vol. 7, p. 11615, May 2016.
- [87] R. Singh *et al.*, “Large power factor and anomalous Hall effect and their correlation with observed linear magneto resistance in Co-doped Bi₂Se₃ 3D topological insulator,” *J. Phys. Condens. Matter*, vol. 28, no. 37, p. 376001, Sep. 2016.
- [88] C. Weyrich *et al.*, “Magnetoresistance oscillations in MBE-grown Sb₂Te₃ thin films,” *Appl. Phys. Lett.*, vol. 110, no. 9, p. 92104, Feb. 2017.
- [89] L. Zhang *et al.*, “Ferromagnetism in vanadium-doped Bi₂Se₃ topological insulator films,” *APL Mater.*, vol. 5, no. 7, p. 76106, Jul. 2017.
- [90] H. Li *et al.*, “Negative magnetoresistance in Dirac semimetal Cd₃As₂,” *Nat. Commun.*, vol. 7, p. 10301, Jan. 2016.
- [91] H. Keiter and G. Morandi, “Thermodynamic perturbation theory for systems with strong local correlations,” *Phys. Rep.*, vol. 109, no. 5, pp. 227–308, Jul. 1984.
- [92] X. F. Sun, K. Segawa, and Y. Ando, “Low-temperature nodal-quasiparticle transport in lightly doped YBa₂Cu₃O_y near the edge of the superconducting doping regime,” *Phys. Rev. B - Condens. Matter Mater. Phys.*, vol. 72, no. 10, pp. 1–4, 2005.
- [93] H. B. Nielsen and M. Ninomiya, “The Adler-Bell-Jackiw anomaly and Weyl fermions in a crystal,” *Phys. Lett. B*, vol. 130, no. 6, pp. 389–396, Nov. 1983.
- [94] V. Aji, “Adler-Bell-Jackiw anomaly in Weyl semimetals: Application to pyrochlore iridates,” *Phys. Rev. B*, vol. 85, no. 24, p. 241101, Jun. 2012.
- [95] D. T. Son and B. Z. Spivak, “Chiral anomaly and classical negative magnetoresistance of Weyl metals,” *Phys. Rev. B*, vol. 88, no. 10, p. 104412, Sep. 2013.
- [96] H. W. Jiang, I. Glazman, C. E. Johnson, and S. T. Hannahs, “Magnetic-field-induced transition from an Anderson insulator to a quantum Hall conductor,” *Surf. Sci.*, vol. 305, no. 1–3, pp. 120–123, Mar. 1994.
- [97] S. X. Zhang *et al.*, “Magneto-resistance up to 60 Tesla in topological insulator Bi₂Te₃ thin films,” *Appl. Phys. Lett.*, vol. 101, no. 20, p. 202403, Nov. 2012.
- [98] S. Zhao *et al.*, “Manipulating the quantum interference effect and magnetotransport of ZnO nanowires through interfacial doping,” *Nanoscale*, vol. 9, no. 44, pp. 17610–17616, 2017.
- [99] L.-F. Chen *et al.*, “Synthesis of Nitrogen-Doped Porous Carbon Nanofibers as an Efficient Electrode Material for Supercapacitors,” *ACS Nano*, vol. 6, no. 8, pp. 7092–7102, Aug. 2012.

- [100] O. Breunig, Z. Wang, A. A. Taskin, J. Lux, A. Rosch, and Y. Ando, “Gigantic negative magnetoresistance in the bulk of a disordered topological insulator,” *Nat. Commun.*, vol. 8, p. 15545, May 2017.
- [101] J. Dai *et al.*, “Toward the Intrinsic Limit of the Topological Bi₂Se₃,” *Phys. Rev. Lett.*, vol. 117, no. 10, p. 106401, Aug. 2016.
- [102] N. Borgwardt *et al.*, “Self-organized charge puddles in a three-dimensional topological material,” *Phys. Rev. B*, vol. 93, no. 24, p. 245149, Jun. 2016.
- [103] S. Hikami, A. I. Larkin, and Y. Nagaoka, “Spin-Orbit Interaction and Magnetoresistance in the Two Dimensional Random System,” *Prog. Theor. Phys.*, vol. 63, no. 2, pp. 707–710, Feb. 1980.
- [104] P. Goswami, J. H. Pixley, and S. Das Sarma, “Axial anomaly and longitudinal magnetoresistance of a generic three-dimensional metal,” *Phys. Rev. B*, vol. 92, no. 7, p. 75205, Aug. 2015.
- [105] J. Kim and S.-H. Jhi, “Magnetic phase transition in Fe-doped topological insulator Bi₂Se₃,” *Phys. Rev. B*, vol. 92, no. 10, p. 104405, Sep. 2015.
- [106] D. L. Sidebottom, “Universal Approach for Scaling the ac Conductivity in Ionic Glasses,” *Phys. Rev. Lett.*, vol. 82, no. 18, pp. 3653–3656, May 1999.
- [107] S.-Y. Xu *et al.*, “Hedgehog spin texture and Berry’s phase tuning in a magnetic topological insulator,” *Nat. Phys.*, vol. 8, no. 8, pp. 616–622, Aug. 2012.
- [108] L. A. Wray *et al.*, “A topological insulator surface under strong Coulomb, magnetic and disorder perturbations,” *Nat. Phys.*, vol. 7, no. 1, pp. 32–37, Jan. 2011.
- [109] D. Biswas, S. Thakur, G. Balakrishnan, and K. Maiti, “Exceptional surface and bulk electronic structures in a topological insulator, Bi₂Se₃,” *Sci. Rep.*, vol. 5, no. 1, p. 17351, Dec. 2015.
- [110] M. R. Scholz *et al.*, “Tolerance of Topological Surface States towards Magnetic Moments: Fe on Bi₂Te₃ and Bi₂Se₃,” *Phys. Rev. Lett.*, vol. 108, no. 25, p. 256810, Jun. 2012.
- [111] H.-J. Kim *et al.*, “Topological Phase Transitions Driven by Magnetic Phase Transitions in Fe_xBi₂Te₃(0 < x < 0.1) single crystal,” *Phys. Rev. Lett.*, vol. 110, no. 13, p. 136601, Mar. 2013.
- [112] T. Valla, Z.-H. Pan, D. Gardner, Y. S. Lee, and S. Chu, “Photoemission Spectroscopy of Magnetic and Nonmagnetic Impurities on the Surface of the Bi₂Se₃ Topological Insulator,” *Phys. Rev. Lett.*, vol. 108, no. 11, p. 117601, Mar. 2012.
- [113] D. Das, K. Malik, S. Bandyopadhyay, D. Das, S. Chatterjee, and A. Banerjee, “Magneto-resistive property study of direct and indirect band gap thermoelectric Bi-Sb alloys,” *Appl. Phys. Lett.*, vol. 105, no. 8, p. 82105, Aug. 2014.
- [114] B. Raquet, M. Viret, E. Sondergard, O. Cespedes, and R. Mamy, “Electron-magnon scattering and magnetic resistivity in 3d ferromagnets” *Phys. Rev. B*, vol. 66, no. 2, p. 24433, Jul. 2002.
- [115] C. H. Marrows and B. C. Dalton, “Spin Mixing and Spin-Current Asymmetry Measured by Domain Wall Magnetoresistance,” *Phys. Rev. Lett.*, vol. 92, no. 9, p. 97206, Mar. 2004.

