

List of figures

Figure No.	Figure Caption	Page No.
Figure 1.1	Schematic represents various component of a sensor involved in sensing.	3
Figure 1.2	Schematics of electrochemical sensors.	5
Figure 1.3	Au@Pd-RGO/GCE based sensors for the simultaneous determination of DA, AA and UA.	7
Figure 1.4	Optical colorimetric sensing of PANI-LBNF strips after 20 min incubation in aqueous solution of various concentrations of Hg ²⁺ .	10
Figure 1.5	Schematic representation of the composite of (a) partially reduced graphene oxide- gold nanorods and (b) MoS ₂ -gold nanoparticles.	16
Figure 1.6	Calixarene capped gold nanoparticles act as catalyst for the 4-nitrophenol reduction to 4-aminophenol.	21
Figure 1.7	A schematic overview of MOFs synthesis, applications and properties.	23
Figure 1.8	(a) Luminescence spectra of Tb(BTC)(H ₂ O) ₆ in Cu(NO ₃) ₂ in aqueous solution at various concentrations (excitation wavelength 300 nm). (b) Emission spectra of Tb(BTC)(H ₂ O) ₆ in aqueous suspension in presence of different acetone contents (excitation wavelength 300 nm)	24
Figure 1.9	Biosensing Mechanism and design of the MOCs enzyme Biocomposites (MEBCs)-Based Biosensor (with MEBC 3 As an Example)	25
Figure 1.10	Structure of 1,1'-2,2'-binaphthol based ligand.	27
Figure 1.11	Catalytic activity of {[Ti ₂ (μ-O) ₂ (binol)]} _n in asymmetric carbonyl-ene reaction.	28
Figure 1.12	Schematic illustration of oxidation reaction of TMB in the presence of H ₂ O ₂ catalyzed by QDs.	30
Figure 1.13	Schematic representation of immobilization methods.	32
Figure 2.1	Schematic representation of dual beam UV-Vis spectrophotometer.	41

List of figures

Figure 2.2	Block diagram for FT-IR.	42
Figure 2.3	Schematic representation of X-ray beam incident on a crystallographic material.	44
Figure 2.4	Physical process involved in XPS	46
Figure 3.1	Synthesis of MoS ₂ -QDs.	56
Figure 3.2	Absorption spectra of (a) MoS ₂ -QDs and (b) AuNPs@MoS ₂ -QDs, Inset of (b) shows the enlarged view of AuNPs@MoS ₂ -QDs peak. (524 nm)	59
Figure 3.3	X-ray Diffraction of (a) MoS ₂ QDs (b) AuNPs@MoS ₂ -QDs obtained from JCPDS (CAS no. 77-1716 and 7440-57-5) file, (c) and (d) are the zeta potential graph of MoS ₂ QDs and AuNPs@MoS ₂ -QDs respectively.	60
Figure 3.4	(a) TEM images of MoS ₂ -QDs and corresponding SAED pattern (inset), (b) particle size distribution of MoS ₂ -QDs.	60
Figure 3.5	(a), (b) and (c) TEM image of AuNPs@MoS ₂ -QDs composite (d) HR TEM image of AuNPs@MoS ₂ -QDs showing lattice fringe spacing of AuNPs and MoS ₂ -QDs Inset of figure (a) shows corresponding SAED pattern.	61
Figure 3.6	EDS mapping of AuNPs@MoS ₂ -QDs composite with overlapped image.	62
Figure 3.7	AFM image of MoS ₂ QDs and AuNPs@MoS ₂ -QDs composites.	63
Figure 3.8	The absorbance spectra of TMB oxidation products for different reaction systems: (A) (a)AuNPs@MoS ₂ -QDs composite, (b) AuNPs@MoS ₂ -QDs composite +TMB, (Inset image shows enlarge view). (c) AuNPs@MoS ₂ -QDs composite +TMB + H ₂ O ₂ and (B) time dependent study of (a) AuNPs@MoS ₂ -QDs composite, (b) TMB + H ₂ O ₂ and (c)AuNPs@MoS ₂ -QDs composite +TMB + H ₂ O ₂ . (Inset image shows corresponding color image).The reaction system contains AuNPs@MoS ₂ -QDs composite (0.2 µg/ml), TMB (1mM) and H ₂ O ₂ (0.15mM).	65
Figure 3.9	UV-Vis spectra of different catalytic system (MoS ₂ -QDs,	65

List of figures

- AuNPs, simple mixture of AuNPs-MoS₂-QDs and AuNPs@MoS₂-QDs composite)
- Figure 3.10** The catalytic reaction at the presence of different radical scavengers. 66
- Figure 3.11** Dependency of AuNPs@MoS₂-QDs composite catalytic activity on (a) Temperature, (b) pH and (c) H₂O₂ concentration. Optimize reaction conditions are AuNPs@MoS₂-QDs composite(0.2 μg/ml), TMB (1mM),H₂O₂(0.15 mM), 1ml of acetate buffer (0.2 M, pH 4.0) at 40°C. 68
- Figure 3.12** The enzyme kinetic parameter of AuNPs@MoS₂-QDs composite system toward substrates (a and b). Optimized reaction condition of AuNPs@MoS₂-QDs composite (0.2μg/ml) in 1ml of 0.2 M sodium acetate buffer at 40°C (pH 4.0). (a) Kinetics of AuNPs@MoS₂-QDs composite for H₂O₂ in fix TMB (0.1 mM) (b) kinetic for TMB in fix H₂O₂ concentration (1mM). 70
- Figure 3.13** (a) Colorimetric response, (b) corresponding calibration plot, (c) Electrochemical response and (d) corresponding calibration plot of AuNPs@MoS₂-QDs modified electrode catalysis of H₂O₂ in 0.1 M PBS, (pH, 7.0) 73
- Figure 3.14** The catalytic mechanism of TMB oxidation in presence H₂O₂catalyzed by AuNPs@MoS₂-QDs composite. 74
- Figure 3.15** (a) UV-Visible spectra of colorimetric sensing of glucose based on AuNPs@MoS₂-QDs composite+ TMB + GOx in buffer. Inset shows visible color in presence of glucose (1, 2, 5, 10, 15, 20, 30, 50, 100, 150, 200, 300 and 400 μM) (b) calibration curve. 74
- Figure 3.16** (a) Interference study and (b) Selectivity of developed sensor system towards glucose. Reaction condition are 0.5 ml of sodium acetate buffer (0.2 M, pH 4.0) with TMB (1 mM), 0.2 μg /ml, AuNPs@MoS₂-QDs composite and 1 mg/ml, glucose oxidase with different interference. 76

List of figures

Figure 3.17	The reproducibility test of developed sensing method (a) Intra-day and (b) Inter-day repeatability for same setup condition for glucose (1 μ M to 400 μ M) detection.	77
Figure 3.18	(a) UV-Vis spectra of test of glucose level in serum by portable test kit (based on AuNPs@MoS ₂ -QDs composite) and inset change the color of wells hydrogel with presence glucose level, with typical colorimetric chart with level glucose (in order to 0, 2, 4, 5, 8, 10, 11, 12 mM), (top) and (b) calibration curve.	77
Figure 3.19	Absorption spectra of glucose level test in tear by portable test kit and inset change the color of wells hydrogel with presence glucose level (50 μ M, 100, 250, 400, 600, 800 and 1mM) in tear (b) corresponding calibration curve.	79
Figure 3.20	Absorption spectra of glucose level test in saliva by portable test kit and inset shows change in the color of wells hydrogel with presence glucose level (5 μ M, 150, 250, 350, 450, 550, 700 and 800 μ M) in saliva (b) corresponding calibration curve.	79
Figure 4.1	Reaction scheme for synthesis of nano porous co-ordination polymer AHMT-Pd.	87
Figure 4.2	Portable test kit procedure for the detection of glucose in serum samples. (A) Opening the snap cap of the test kit (B) adding diluted serum sample solution to the test kit, (C) closing the snap cap and turning the test kit upside down before incubation at 40 °C for 30 min, and (D) turning the test kit upside down again and opening the snap cap to observe the color change of hydrogel (blue color in the hydrogel).	88
Figure 4.3	FT-IR spectra of AHMT (a) and AHMT-Pd (b).	91
Figure 4.4	X-ray diffraction of AHMT-Pd (a) AHMT (b) and Pd (0) (c) obtained from JCPDS (CAS no. 65-6174.) file.	92
Figure 4.5	Absorption spectra of (a) AHMT and (b) AHMT –Pd.	92
Figure 4.6	XPS spectra of AHMT–Pd for Pd (3d), N (1s), S (2p) and C (1s) regions.	94

List of figures

Figure 4.7	Structural morphology of AHMT-Pd (a) and (b) HR SEM, (c) and (d) TEM, inset of (d) corresponding SAED pattern.	95
Figure 4.8	EDX of AHMT-Pd provided by HR TEM graph.	96
Figure 4.9	EDS Mapping of AHMT-Pd.	96
Figure 4.10	Structural network of AHMT-Pd.	97
Figure 4.11	(a) TGA (left ordinate) and (b) DTA (right ordinate) plot of AHMT-Pd.	98
Figure 4.12	Plot of the Thermal activation energy for AHMT-Pd.	99
Figure 4.13	(A) Absorption spectra of various combinations of reaction systems (B) A typical absorbance peak changes with of TMB oxidation at 652 nm in different systems: AHMT-Pd, AHMT-Pd + TMB, TMB + H ₂ O ₂ , TMB + AHMT-Pd + H ₂ O ₂ . (Inset of B shows corresponding photograph of TMB oxidation products). AHMT-Pd (0.5 μg mL ⁻¹), TMB (1mM) and H ₂ O ₂ 0.5 mM.	100
Figure 4.14	Possible mechanism for the electron transfer investigation and OH radical's formation for production of colored TMB oxidation product in catalysis reaction.	101
Figure 4.15	(a) The Normalized UV-Vis spectrum of AHMT-Pd, (b) $(\alpha h\nu)^2$ versus optical band gap achieved from equation.	102
Figure 4.16	The peroxidase-like activity of AHMT-Pd and dependency on a) temperature, (b) pH, (c) H ₂ O ₂ concentration and (d) AHMT-Pd concentration. Experiments were carried out at 0.5 μg·mL ⁻¹ AHMT-Pd in 1mL of 0.2 M acetate buffer with TMB (5 mM) as substrate. The H ₂ O ₂ concentration was 0.5 mM at pH 4.0, the temperature was 40 °C	104
Figure 4.17	The enzyme kinetic K _m and V _{max} of AHMT-Pd toward TMB substrates (a and b). The analysis of reaction rate in presence of AHMT-Pd, 0.5 μg·mL ⁻¹ in 1 mL of 0.2 M sodium acetate buffer at pH 4.0, 40 °C. (a) The H ₂ O ₂ concentration (10 mM) was fixed for AHMT-Pd and various TMB concentrations (b) Concentration of reactant TMB (0.1 mM) was fixed and the H ₂ O ₂ was varied for AHMT-Pd.	107

List of figures

- Figure 4.18** Au@Pd-RGO/GCE based sensors for the simultaneous determination of DA, AA and UA. **107**
- Figure 4.19** Electrochemical response of AHMT-Pd modified electrode catalysis of H₂O₂ in 0.1 M PBS, (pH, 7.0) **108**
- Figure 4.20** Schematic representation of the composite of (a) partially reduced graphene oxide- gold nanorods and (b) MoS₂-gold nanoparticles. **109**
- Figure 4.21** (a) UV-Visible spectra of a mixture of AHMT-Pd + TMB + GOx with different concentration of glucose (1 μM, 2, 10, 15, 20, 30, 50, 100, 150, 200 and 300 μM) (b) corresponding calibration curve. **110**
- Figure 4.22** (a) Intra-day and (b) inter-day repeatability test result of AHMT-Pd (0.5 μgmL⁻¹) mediated detection of glucose (concentration range 1 μM - 300 μM). The error bar represents the standard deviation of triplicate measurement. **111**
- Figure 4.23** The catalytic reaction at the presence of different interferences. **112**
- Figure 4.24** (a) Specificity of method for glucose detection. Specificity checked in 0.5 mL of 0.2 M sodium acetate buffer (pH 4.0) with TMB (1 mM), 0.5 μgmL⁻¹, AHMT-Pd and 1 mgmL⁻¹, glucose oxidase. (b) Ten times repeated experiments for glucose detection with AHMT-Pd (0.5 μgmL⁻¹), (1 mM) TMB solution and measurement in same set of reaction condition with GOx (1 mgmL⁻¹) and (0.1 mM) glucose. The error bars represent the standard deviation of three-time measurements. **113**
- Figure 4.25** Human serum glucose level and corresponding typical colorimetric chart (from left to right: 0, 2, 4, 5, 8, 10, 11, 12 mM) (top); the hydrogel color change in presence of various concentrations of glucose in human complex system (opening the snap cap). **114**
- Figure 4.26** UV-Visible spectra of AHMT-Pd + TMB + GOx with different concentration of glucose (10 μM, 100, 200, 400, **115**

List of figures

	600, 800 and 1000 μM) in tear (b) corresponding calibration curve (c) UV-Visible spectra of AHMT-Pd + TMB + GOx with different concentration of glucose (5 μM , 50, 100, 200, 400, 600 and μM) in saliva (b) corresponding calibration curve.	
Figure 5.1	Synthesis of tungsten disulfide-quantum dots (WS_2 -QDs).	121
Figure 5.2	Absorption spectra of (a) WS_2 -QDs and (b) AuNPs@ WS_2 -QDs, Inset of (b) shows the enlarged view of AuNPs@ WS_2 -QDs peak. (600 nm).	123
Figure 5.3	X-ray Diffraction of (a) WS_2 -QDs (b) AuNPs@ WS_2 -QDs obtained from JCPDS (CAS no. 84-1398 and 65-2870) file.	123
Figure 5.4	(a) and (b) are the zeta potential graph of WS_2 -QDs and AuNPs@ WS_2 -QDs respectively.	124
Figure 5.5	(a) TEM image of WS_2 -QDs and inset shows SEAD pattern (b) particle size distribution graph of WS_2 -QDs (c,d,e) TEM image of AuNPs@ WS_2 -QDs composite (f) HRTEM image of AuNPs@ WS_2 -QDs with lattice fringe spacing of AuNPs ~ 0.23 nm and WS_2 -QDs ~ 0.21 nm and inset of (c) shows SAED pattern of AuNPs@ WS_2 -QDs to confirm crystalline nature.	125
Figure 5.6	EDS mapping of AuNPs@ WS_2 -QDs composite with overlapped image.	126
Figure 5.7	AFM image of WS_2 -QDs and AuNPs@ WS_2 -QDs composites	127
Figure 5.8	(a) UV-visible absorption spectra of reaction system containing AuNPs@ WS_2 -QDs composite+TMB+ H_2O_2 (B) Time-dependent absorbance changes at 652 nm of reaction mixture containing AuNPs@ WS_2 -QDs composite, H_2O_2 , and TMB are $1.0 \mu\text{g mL}^{-1}$, 0.5 mM, and 2.0 mM respectively.	128
Figure 5.9	The catalytic reaction system in presence of various radical scavengers.	129
Figure 5.10	Optimizations of various parameters for catalytic activity of AuNPs@ WS_2 -QDs composite: (a) H_2O_2 , (b) temperature and (c) pH. Experiments were carried out mixture of	131

List of figures

- AuNPs@WS₂-QDs (1 μg.mL⁻¹), TMB (1mM) and H₂O₂ (0.5 mM) in pH 4.0 maintained by acetate buffer at 40°C.
- Figure 5.11** Steady state kinetic measurement of AuNPs@WS₂-QDs composite through (a) variation of H₂O₂ concentration and fix TMB and (b) variation of TMB concentration and fix H₂O₂. **132**
- Figure 5.12** (a) Concentration dependent UV-Visible graph for H₂O₂ detection (1 to 200 μM) and (b) linear calibration plot for H₂O₂. Inset of (a) shows the images of TMB oxidation colored product for different concentration of H₂O₂. **133**
- Figure 5.13** Mechanism of choline detection based on TMB oxidation catalyzed by AuNPs@WS₂-QDs composite. **134**
- Figure 5.14** (a) UV-Visible spectra of colorimetric sensing of choline based on AuNPs@WS₂-QDs composite + TMB + ChOx in buffer, inset shows visible color of various concentration of choline and (b) corresponding calibration curve at 652 nm. **135**
- Figure 5.15** (a) Selectivity analysis AuNPs@WS₂-QDs composite system with interfering substances by recording absorbance at 652 nm, including 10.0 mM of ascorbic acid, cysteine, cholesterol, uric acid, urea and (b) reproducibility of six samples prepared in same set of conditions. **135**
- Figure 5.16** The manufacture of AuNPs@WS₂-QDs composite based choline detection test strips for quick detection of choline level by using naked eyes. **137**
- Figure 5.17** Demonstration of the utility of paper strip in the analysis of choline in real sample. Visualize the choline in real sample by paper strip (a) control,(b) 120 μM. **137**
- Figure 5.18** (a) UV-Visible spectra of colorimetric sensing of choline based on AuNPs@WS₂-QDs composite + TMB + ChOx in milk, inset shows visible color of various concentration of choline and (b) corresponding calibration curve at 652 nm. **138**
- Figure 5.19** (a) UV-Visible spectra of colorimetric sensing of choline based on AuNPs@WS₂-QDs composite + TMB + ChOx in serum, inset shows visible color of various concentration of **138**

List of figures

	choline and (b) corresponding calibration curve at 652 nm.	
Figure 6.1	Schematics of Ag@AHMT synthesis	145
Figure 6.2	Time dependent study for the formation of Ag@AHMT nanoparticles by UV-Vis spectrum, (at 0 min, 5 min, 15 min, 25 min, 35 min, 45 min, 55 min)	146
Figure 6.3	(a) UV-Vis spectra (b) FT-IR Spectra of AHMT and Ag@AHMT nanoparticles.	146
Figure 6.4	(a) X-ray Diffraction of Silver nanoparticles indexed from JCPDS(CAS no. 7440-22-4) file and (b) XPS spectrum of Ag (0)	147
Figure 6.5	HR SEM of Ag@AHMT nanoparticles.	148
Figure 6.6	TEM images of Ag@AHMT nanoparticles (a,b,c) and corresponding SAED pattern (d).	150
Figure 6.7	Particle size distribution of Ag@AHMT nanoparticles.	150
Figure 6.8	EDAX of Ag@AHMT nanoparticles.	151
Figure 6.9	EDS Mapping of Ag@AHMT silver nanoparticles.	151
Figure 6.10	Surface Zeta potential graph of Ag@AHMT nanoparticles (a) and naked silver nanoparticles (b)	152
Figure 6.11	Stability of Ag@AHMT Nps checked by UV-Vis.	153
Figure 6.12	(a) Colorimetric sensing with different concentration of picric acid (1, 2, 3, 4, 5, 6, 7, 8 nM). (b) Calibration curve of colorimetric sensing with different concentration of picric acid (1, 2, 3, 4, 5, 6, 7, 8 nM)	154
Figure 6.13	Photographs showing the color-change during colorimetric sensing of Picric acid.	155
Figure 6.14	TEM images of Ag@AHMT+Picric acid.	155
Figure 6.15	The schematic presentation of formation of Ag@AHMT nanoparticles and interaction with picric acid.	156
Figure 6.16	Intra-day (a) and inter-day (b) repeatability test for the detection of picric acid (concentration ranges from 1nM to 8 nM).	157
Figure 7.1	Possible mechanism for formation of AHMT-Ag.	161

List of figures

Figure 7.2	Reaction scheme for synthesis of nanocrystalline coordination polymer AHMT-Ag.	164
Figure 7.3	Schematic illustration of controlled experiments for the formation of AHMT–Ag using TEMPO and BHT.	165
Figure 7.4	(I) FT-IR of spectra (a) AHMT-Ag (b) AHMT and (II) Absorption spectra of (a) AHMT and (b) AHMT –Ag.	166
Figure 7.5	(I) X-ray Diffraction of (a) AHMT-Ag (b) AHMT and (c) Ag (0) obtained from JCPDS (CAS no. 87-0720.) file. (II) Le-bail fit of AHMT-Ag.	167
Figure 7.6	XPS spectra of the AHMT–Ag for Ag (3d), N (1s), S (2p) and C (1s) regions.	169
Figure 7.7	¹ H NMR of (a) AHMT and (b) AHMT –Ag.	170
Figure 7.8	HR SEM of AHMT-Ag.	171
Figure 7.9	HR TEM of AHMT-Ag (a, b), size distribution graph from TEM image (c) and corresponding SAED pattern (d).	172
Figure 7.10	EDX of AHMT–Ag provided by HR-TEM graph.	172
Figure 7.11	EDS Mapping of AHMT-Ag.	173
Figure 7.12	Ball-stick model structure of AMT–Ag in the (1) <i>anti</i> and (2) <i>syn</i> orientation generated by CS 3D Chem bio–draw with structural energy minimization. Here blue, cyan, light grey, golden, dark grey (large size) and pink balls represent N, H, C, S, Ag atoms and lone pairs.	175
Figure 7.13	Three possible structures of the AHMT dimer coordinated with Ag atom obtained at the B3LYP/6-31G**+LanL2DZ level of theory in water solvent.	176
Figure 7.14	Proposed Structural network of AHMT-Ag.	176
Figure 7.15	Zeta potential of AHMT-Ag.	177
Figure 7.16	TGA plot of AHMT-Ag.	178
Figure 7.17	Thermal activation energy plot of AHMT–Ag.	179
Figure 7.18	The CV of the electrodes in 5mM Fe(III) in PBS pH 7 unmodified CPE (a) and AHMT-Ag modified CPE (b).	180
Figure 7.19	Cyclic voltammetric (CV) response by the serial addition of	182

List of figures

6-MP to AHMT–Ag/CPE in PBS at different pH.

- Figure 7.20** (a) CV of AHMT–Ag/CPE in 18 μM 6-MP at scan rates 10, 20, 30, 50, 70 and 100 mV, (b) plots of the 6-MP anodic peak current against $v^{1/2}$ and (c) anodic peak potentials against $\log v$. **183**
- Figure 7.21** Effect of Matrix on the anodic peak current and potential, (A) anodic current vs. dilution times, (B) peak potential vs. dilution times. The spiked concentration of 6-MP in serum is 60 μM (mean \pm SD for $n = 5$). **184**
- Figure 7.22** (a) Differential pulse voltammogram recorded on AHMT–Ag/CPE in 0.1 M PBS (pH 7) by successive addition of 6-MP, inset of (a) cyclic voltammogram recorded by successive addition of 6-MP, (b) corresponding calibration plot. **185**
- Figure 7.23** Mechanism for electro–oxidation of 6-MP. **186**
- Figure 7.24** (a) Differential pulse voltammogram recorded on AHMT–Ag/CPE in 0.1 M PBS (pH 7) by successive addition of 6-MP Tablet, inset of (a) cyclic voltammogram recorded by successive addition of 6-MP, (b) corresponding calibration plot. **187**
- Figure 7.25** (a) Differential pulse voltammogram recorded on AHMT–Ag/CPE in urine sample by successive addition of 6-MP, inset of (a) cyclic voltammogram recorded by successive addition of 6-MP, (b) corresponding calibration plot. **188**
- Figure 7.26** Differential pulse voltammogram recorded on AHMT–Ag/CPE in serum sample by successive addition of 6-MP, inset of (a) cyclic voltammogram recorded by successive addition of 6-MP, (b) corresponding calibration plot. **188**
- Figure 7.27** Analytical figures for the comparison of present work with other methods. **189**
- Figure 7.28** Differential pulse voltammetric (DPV) responses of electrochemical sensor to 55 μM 6-MP, 55 μM 6-MP+ 100 μM ethanol, 55 μM 6-MP + 100 μM glucose, 55 μM 6-MP+ **190**

List of figures

100 μM starch, 55 μM 6-MP+ 100 μM Ascorbic acid, 55 μM 6-MP+ 100 μM uric acid and 55 μM 6-MP+ 100 μM Dopamine. Error bar = RSD, ($n = 5$).